

Tobacco in Australia

Facts & Issues

Relevant news and research

9.9 Are there inequalities in access to and use of treatment for dependence on tobacco-delivered nicotine?

Last updated October 2020

Research:

Zyambo, CM, Hendricks, PS, Cropsey, KL, Willig, JH, Wilson, CM, Gakumo, CA et al (2020). Racial disparities and factors associated with prescription for smoking cessation medications among smokers receiving routine clinical care for HIV. *AIDS Care*, 1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32530307>

Kock, L, & Brown, J. (2019). Commentary on Smith et al. (2019): Equity positive impact of English Stop Smoking Services underlines need for comprehensive approach to public health interventions. *Addiction*. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31677198>

Young-Wolff KC, Adams SR, Tan ASL, Adams AS, Klebaner D, et al. Disparities in knowledge and use of tobacco treatment among smokers in California following healthcare reform. *Preventive Medicine Reports*, 2019; 14:100847. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31024786>

van Wijk EC, Landais LL, and Harting J. Understanding the multitude of barriers that prevent smokers in lower socioeconomic groups from accessing smoking cessation support: A literature review. *Preventive Medicine*, 2019; 123:143-51. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30902700>

Smith CE, Hill SE, and Amos A. Impact of specialist and primary care stop smoking support on socioeconomic inequalities in cessation in the UK: A systematic review and national equity analysis. *Addiction*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31357250>

McQueen A, Roberts C, Garg R, Caburnay C, Fu Q, et al. Specialized tobacco quitline and basic needs navigation interventions to increase cessation among low income smokers: Study protocol for a randomized controlled trial. *Contemporary Clinical Trials*, 2019; 80:40-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30904596>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Smith C, Hill S, and Amos A. Stop smoking inequalities: A systematic review of socioeconomic inequalities in experiences of smoking cessation interventions in the UK. Cancer Research UK, 2018. Available from:

https://www.cancerresearchuk.org/sites/default/files/stop_smoking_inequalities_2018.pdf.

Landrine H, Corral I, and Campbell KM. Racial disparities in healthcare provider advice to quit smoking. Preventive Medicine Reports, 2018; 10:172-5. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29868363>

Danan ER, Fu SS, Clothier BA, Noorbaloochi S, Hammett PJ, et al. The equity impact of proactive outreach to smokers: Analysis of a randomized trial. American Journal of Preventive Medicine, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30139707>

Bailey SR, Heintzman J, Jacob RL, Puro J, and Marino M. Disparities in smoking cessation assistance in US primary care clinics. American Journal of Public Health, 2018:e1-e9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29927641>

Soulakova JN and Crockett LJ. Level of cigarette consumption and duration of smoking abstinence during failed Quit attempts among long-term daily smokers: The role of race/ethnicity and cessation aids. J Racial Ethn Health Disparities, 2017. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28444627>

Hooks-Anderson DR, Salas J, Secret S, Skiold-Hanlin S, and Scherrer JF. Association between race and receipt of counselling or medication for smoking cessation in primary care. Family Practice, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29045650>

Edwards SA, Callaghan RC, Mann RE, and Bondy SJ. Association between socioeconomic status and access to care and quitting smoking with and without assistance. Nicotine & Tobacco Research, 2017; 20(1):40–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28340126>

Dahne J, Wahlquist AE, Garrett-Mayer E, Heckman BW, Michael Cummings K, et al. The differential impact of state tobacco control policies on cessation treatment utilization across established tobacco disparities groups. Preventive Medicine, 2017; 105:319–25. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28987337>

Brose LS and McEwen A. Neighbourhood deprivation and outcomes of stop smoking support - an observational study. PLoS ONE, 2016; 11(1):e0148194. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26824352>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Wang L, Jin Y, Lu B, and Ferketich AK. A cross-country study of smoking cessation assistance utilization in 16 low and middle income countries: Data from the global adult Tobacco survey (2008-2012). *Nicotine & Tobacco Research*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26116084>

Paul C, Wolfenden L, Tzelepis F, Yoong S, Bowman J, et al. Nicotine replacement therapy as a smoking cessation aid among disadvantaged smokers: What answers do we need? *Drug and Alcohol Review*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26661119>

Brathwaite R, Addo J, Smeeth L, and Lock K. A systematic review of Tobacco smoking prevalence and description of Tobacco control strategies in sub-saharan african countries; 2007 to 2014. *PLoS ONE*, 2015; 10(7):e0132401. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26162085>

Bosdriesz JR, Willemsen MC, Stronks K, and Kunst AE. Socioeconomic inequalities in smoking cessation in 11 European countries from 1987 to 2012. *Journal of Epidemiology and Community Health*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25841241>

Kotz D. Rigorous methodology is needed to analyze and interpret observational data on the use and effectiveness of smoking cessation aids. *Nicotine & Tobacco Research*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25320111>

Clare P, Slade T, Courtney RJ, Martire KA, and Mattick RP. The use of smoking cessation and Quit support services by socioeconomic status over 10 years of the National Drug Strategy Household Survey. *Nicotine & Tobacco Research*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25320110>

Aspinall PJ and Mitton L. Smoking prevalence and the changing risk profiles in the UK ethnic and migrant minority populations: Implications for stop smoking services. *Public Health*, 2014; 128(3):297–306. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24612958>

Wang X and Abughosh S. Racial and ethnic disparity in smoking cessation medication use among adult smokers in the United States. *Value in Health*, 2013; 16(3):A241. Available from: [http://www.valueinhealthjournal.com/article/S1098-3015\(13\)01297-7/fulltext](http://www.valueinhealthjournal.com/article/S1098-3015(13)01297-7/fulltext)

Huang Y, Britton J, Hubbard R, and Lewis S. Who receives prescriptions for smoking cessation medications? An association rule mining analysis using a large primary care database. *Tobacco Control*, 2012; [Epub ahead of print]. Available from: <http://tobaccocontrol.bmj.com/content/early/2012/01/13/tobaccocontrol-2011-050124.full>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Christiansen B, Reeder K, Hill M, Baker TB, and Fiore MC. Barriers to effective tobacco-dependence treatment for the very poor. *J Stud Alcohol Drugs*, 2012; 73(6):874-84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23036204>

Christiansen B, Reeder K, Hill M, Baker T, and Fiore M. Barriers to effective tobacco-dependence treatment for the very poor. *Journal of Studies on Alcohol and Drugs*, 2012; 73(6):874–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23036204>

Zhu S, Wong S, Stevens C, Nakashima D, and Gamst A. Use of a smokers' Quitline by Asian language speakers: Results from 15 years of operation in California. *American Journal of Public Health*, 2010; [Epub ahead of print]. Available from: <http://ajph.aphapublications.org/cgi/reprint/AJPH.2009.168385v1?view=long&pmid=20299658>

Thornley S, Jackson G, McRobbie H, Sinclair S, and Smith J. Few smokers in South Auckland access subsidised nicotine replacement therapy. *The New Zealand Medical Journal*, 2010; 123(1308):16–27. Available from: <http://www.nzma.org.nz/journal/123-1308/3943/>

Willett J, Hood N, Burns E, Swetlick J, Wilson S, et al. Clinical faxed referrals to a tobacco Quitline reach, enrollment, and participant characteristics. *American Journal of Preventive Medicine*, 2009; 36(4):337–40. Available from: <http://www.ajpm-online.net/article/S0749-3797%2808%2901010-6/abstract>

Stuber J and Galea S. Who conceals their smoking status from their health care provider? *Nicotine & Tobacco Research*, 2009; 11(3):303–7. Available from: <http://ntr.oxfordjournals.org/content/11/3/303.long>

Solberg L, Parker E, Foldes S, and Walker P. Disparities in tobacco cessation medication orders and fills among special populations. *Nicotine & Tobacco Research*, 2009; 12(2):144–51. Available from: <http://ntr.oxfordjournals.org/content/12/2/144.long>

Miller CL and Sedivy V. Using a quitline plus low-cost nicotine replacement therapy to help disadvantaged smokers to quit. *Tobacco Control*, 2009; 18(2):144–9. Available from: <http://tobaccocontrol.bmj.com/cgi/content/abstract/18/2/144>

Knudsen HK. Smoking cessation services in adolescent substance abuse treatment: Opportunities missed? *Journal of Drug Issues*, 2009; 39(2). Available from: <http://www2.criminology.fsu.edu/~jdi/39n2.htm>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Kaestle C and Wiles B. Targeting high-risk neighborhoods for tobacco prevention education in schools. *American Journal of Public Health*, 2009; 100(9):1708–13. Available from: <http://ajph.aphapublications.org/cgi/content/full/100/9/1708?view=long&pmid=20019323>

Chang F, Hu T, Lo S, Yu P, Chao K, et al. Quit smoking advice from health professionals in taiwan: The role of funding policy and smokers' socioeconomic status. *Tobacco Control*, 2009; 19:44–9. Available from: <http://tobaccocontrol.bmj.com/content/19/1/44.full?sid=27a3840d-26fa-4c5c-acab-1ae919062be6>

Boles M, Rohde K, He H, Maher J, Stark M, et al. Effectiveness of a tobacco quitline in an indigenous population: A comparison between alaska native people and other first-time quitline callers who set a quit date. *International Journal of Circumpolar Health*, 2009; 68(2):170–81. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19517876

Sood A, Andoh J, Verhulst SJ, Rajoli N, and Hopkins-Price P. Characteristics of smokers calling a national reactive telephone helpline. *American Journal of Health Promotion*, 2008; 22(13):176–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18251117>

Sonnenfeld N, Schappert S, and Lin S. Racial and ethnic differences in delivery of tobacco-cessation services. *American Journal of Preventive Medicine*, 2008; 36(1):21–8. Available from: <http://www.ajpm-online.net/article/S0749-3797%2808%2900844-1/fulltext>

Reed M and Burns D. A population-based examination of racial and ethnic differences in receiving physicians' advice to quit smoking. *Nicotine & Tobacco Research*, 2008; 10(9):1487–94. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802323217>

Fu S, Kodl M, Joseph A, Hatsukami D, Johnson E, et al. Racial/ethnic disparities in the use of nicotine replacement therapy and quit ratios in lifetime smokers ages 25 to 44 years. *Cancer Epidemiology Biomarkers & Prevention*, 2008; 17(7):1640–7. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2593846/>

Burns E and Levinson A. Discontinuation of nicotine replacement therapy among smoking-cessation attempters. *American Journal of Preventive Medicine*, 2008; 34(3):212–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18312809>

Browning K, Ferketich A, Salsberry P, and Wewers M. Socioeconomic disparity in provider-delivered assistance to quit smoking. *Nicotine & Tobacco Research*, 2008; 10(1):55–61. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200701704905>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Andoh J, Verhulst S, Ganesh M, Hopkins-Price P, Edson B, et al. Sex- and race-related differences among smokers using a national helpline are not explained by socioeconomic status. *Journal of the National Medical Association*, 2008; 100(2):200–7. Available from: www.nmanet.org/images/uploads/Journal/OC200.pdf

Siahpush M, Wakefield M, Spittal M, and Durkin S. Anti-smoking television advertising and socioeconomic variations in calls to Quitline. *Journal of Epidemiology and Community Health*, 2007; 61(4):298–301. Available from: <http://jech.bmj.com/cgi/content/full/61/4/298>

Lillard DR, Plassmann V, Kenkel D, and Mathios A. Who kicks the habit and how they do it: Socioeconomic differences across methods of quitting smoking in the USA. *Social Science & Medicine*, 2007; 64(12):2504–19. Available from: http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6VBF-4NDVHDN-1&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=9c995358ca89698ee32e3f83db6ee0be

Elin R, Nick R, Alison C, Sarah L, and John B. Use of nicotine replacement therapy in socioeconomically deprived young smokers: A community-based pilot randomised controlled trial. *Tobacco Control*, 2006; 15(5):373–6. Available from: <http://ejournals.ebsco.com/direct.asp?ArticleID=47B3970FE1458C036B27>

Chapman S and Liberman J. Ensuring smokers are adequately informed: Reflections on consumer rights, manufacturer responsibilities, and policy implications. *Tobacco Control*, 2005; 14(suppl. 2):ii8–ii13. Available from: http://tc.bmjournals.com/cgi/content/abstract/14/suppl_2/ii8

Burns D and Warner K. Smokers who have not Quit: Is cessation more difficult and should we change our strategies, in those who continue to smoke. *Smoking and Tobacco Control Monograph no.15*. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute, 2003. Available from: <http://cancercontrol.cancer.gov/tcrb/monographs/15/monograph15-chapter1.pdf>.

Liberman J and Clough J. Corporations that kill: The criminal liability of tobacco manufacturers. *Criminal Law Journal*, 2002; 26(4):223. Available from: http://www.vctc.org.au/tc-res/criminal_liability_paper.pdf

Borland R, Pigott R, Rintoul D, Shore S, and Young S. Barriers to access of smoking cessation programs, nicotine replacement therapy and other pharmacotherapies for the general Australian population and at risk population groups. Literature review and volumes 1 and 2. Canberra: Final Report to Commonwealth Department of Health and Ageing, 2002.

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Chapman S and Wakefield M. Tobacco control advocacy in Australia: Reflections on 30 years of progress. *Health Education & Behavior*, 2001; 28(3):274–89. Available from: <http://heb.sagepub.com/cgi/content/abstract/28/3/274>

Carter S, Borland R, and Chapman C. Finding the strength to kill your best friend—smokers talk about smoking and quitting. Sydney: Australian Smoking Cessation Consortium and GlaxoSmithKline Consumer Healthcare, 2001. Available from: <http://tobacco.health.usyd.edu.au/site/supersite/resources/pdfs/killbestfriend.pdf>.

Royal College of Physicians of London. Nicotine addiction in Britain. A report of the Tobacco advisory group of the royal college of physicians. London: Royal College of Physicians of London, 2000. Available from: <http://bookshop.rcplondon.ac.uk/details.aspx?e=131>.

Chapman S, Wilson D, and Wakefield M. Smokers' understanding of cigarette yield labels. *Medical Journal of Australia*, 1986; 145(8):376–9.

9.9.1 Quitlines

Griffin JL, Segal KS, and Nahvi S. Barriers to telephone Quitline use among methadone-maintained smokers. *Nicotine & Tobacco Research*, 2015; 17(8):931–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26180217>

Varghese M, Sheffer C, Stitzer M, Landes R, Brackman SL, et al. Socioeconomic disparities in telephone-based treatment of tobacco dependence. *American Journal of Public Health*, 2014; 104(8):e76–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24922165>

Campbell HS, Baskerville NB, Hayward LM, Brown KS, and Ossip DJ. The reach ratio—a new indicator for comparing quitline reach into smoking subgroups. *Nicotine & Tobacco Research*, 2014; 16(4):491–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24311698>

9.9.2 Disparities in use of treatments for tobacco dependence

Dahne, J, Wahlquist, AE, Smith, TT, & Carpenter, MJ. (2020). The differential impact of nicotine replacement therapy sampling on cessation outcomes across established tobacco disparities groups. *Prev Med*, 136, 106096. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32320705>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

Goodin, A, Talbert, J, Freeman, PR, Hahn, EJ, & Fallin-Bennett, A. (2020). Appalachian disparities in tobacco cessation treatment utilization in Medicaid. *Substance Abuse Treatment, Prevention and Policy*, 15(1), 5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31959212>

Fillon M. Study evaluates mobile telephone interventions for socioeconomically disadvantaged smokers. *CA: A Cancer Journal for Clinicians*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30856675>

Zhu SH, Anderson CM, Zhuang YL, Gamst AC, and Kohatsu ND. Smoking prevalence in medicaid has been declining at a negligible rate. *PLoS ONE*, 2017; 12(5):e0178279. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28542637>

Wilson A, Guillaumier A, George J, Denham A, and Bonevski B. A systematic narrative review of the effectiveness of behavioural smoking cessation interventions in selected disadvantaged groups (2010-2017). *Expert Rev Respir Med*, 2017; 11(8):617-30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28608758>

Sorensen G, Pednekar M, Cordeira LS, Pawar P, Nagler EM, et al. Effects of a worksite tobacco control intervention in India: The mumbai worksite tobacco control study, a cluster-randomised trial. *Tobacco Control*, 2017; 26(2):210-6. Available from: <http://tobaccocontrol.bmj.com/content/tobaccocontrol/26/2/210.full.pdf>

Hammett P, Fu SS, Nelson D, Clothier B, Saul JE, et al. A proactive smoking cessation intervention for socioeconomically disadvantaged smokers: The role of smoking-related stigma. *Nicotine & Tobacco Research*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28398492>

Flocke SA, Hoffman R, Eberth JM, Park H, Birkby G, et al. The prevalence of Tobacco use at federally qualified health centers in the United States, 2013. *Preventing Chronic Disease*, 2017; 14:E29. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28384096>

Feinberg A, Lopez PM, Wyka K, Islam N, Seidl L, et al. Prevalence and correlates of smoking among low-income adults residing in New York city public housing developments-2015. *J Urban Health*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28656541>

Kostandinov O. Community confronting high smoking rates among disadvantaged Quit Resource Centre, 2015. Available from: <http://www.quit.org.au/resource-centre/media-advertising/media-releases/community-confronting-high-smoking-rates-among-disadvantaged.html>

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

O'Brien J, Salmon AM, and Penman A. What has fairness got to do with it? Tackling tobacco among Australia's disadvantaged. *Drug and Alcohol Review*, 2012; 31(5):723–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22524309>

Bobo JK, Anderson JR, and Bowman A. Training chemical dependency counselors to treat nicotine dependence. *Addictive Behaviors*, 1997; 22(1):23–30. Available from: http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6VC9-3SX4Y19-3&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=22b8a6d61b8ca86e042c714e626abd7c

9.9.2.1 Disparities in use of treatments

9.9.2.2 Disparities in compliance with treatment

9.9.3 Disparities in provision of quit smoking advice and referral by general practitioners

Sipe, CJ, Koopmeiners, JS, Donny, EC, Hatsukami, DK, & Murphy, SE. (2020). UGT2B10 Genotype Influences Serum Cotinine Levels and Is a Primary Determinant of Higher Cotinine in African American Smokers. *Cancer Epidemiol Biomarkers Prev*. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32532831>

Angus C, Brown J, Beard E, Gillespie D, Buykx P, et al. Socioeconomic inequalities in the delivery of brief interventions for smoking and excessive drinking: Findings from a cross-sectional household survey in England. *BMJ Open*, 2019; 9(4):e023448. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31048422>

9.9.4 Harm reduction: a strategy of benefit for disadvantaged groups?

Lucherini, M, Hill, S, & Smith, K. (2019). Potential for non-combustible nicotine products to reduce socioeconomic inequalities in smoking: a systematic review and synthesis of best available evidence. [S]. *BMC Public Health*, 19(1), 1469. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31694602>

News reports:

tobaccoinaustralia.org.au

Tobacco in Australia

Facts & Issues

No authors listed. Medicaid tobacco cessation: Big gaps remain in efforts to get smokers to quit. Medical News Today, 2016. Available from: <http://www.medicalnewstoday.com/releases/304728.php?tw>

Medicare Australia. Pbs statistics. Canberra: Medicare Australia, 2008. Last update: Viewed Available from: <http://www.medicareaustralia.gov.au/provider/pbs/stats.shtml>.

9.9.1 Quitlines

9.9.2 Disparities in use of treatments for tobacco dependence

Butt C. Quit Victoria to help tongans give up smoking. The Age, 2015. Available from: <http://www.theage.com.au/victoria/quit-victoria-to-help-tongans-give-up-smoking-20150329-1ma7fr.html>

Partos T and Borland R. Data file of responses to eighth wave of the international Tobacco control four-country survey, in *Provided to Merryn Pearce of the Tobacco Control Unit2012*, Cancer Council Victoria.

9.9.2.1 Disparities in use of treatments

9.9.2.2 Disparities in compliance with treatment

9.9.3 Disparities in provision of quit smoking advice and referral by general practitioners

9.9.4 Harm reduction: a strategy of benefit for disadvantaged groups?