

# Tobacco in Australia

## Facts & Issues

---

### Relevant news and research

#### 9.9 Are there inequalities in access to and use of treatment for dependence on tobacco-delivered nicotine?

*Last updated December 2024*

Research:	2
9.9.1 Quitlines	7
9.9.2 Disparities in use of treatments for tobacco dependence	8
9.9.2.1 Disparities in use of treatments	9
9.9.2.2 Disparities in compliance with treatment	9
9.9.3 Disparities in provision of quit smoking advice and referral by general practitioners	9
9.9.4 Harm reduction: a strategy of benefit for disadvantaged groups?	10
News reports:	10
9.9.1 Quitlines	10
9.9.2 Disparities in use of treatments for tobacco dependence	10
9.9.2.1 Disparities in use of treatments	11
9.9.2.2 Disparities in compliance with treatment	11
9.9.3 Disparities in provision of quit smoking advice and referral by general practitioners	11
9.9.4 Harm reduction: a strategy of benefit for disadvantaged groups?	11

## Research:

**Chandra, M, Talluri, R, Fokom Domgue, J, & Shete, S. (2024). Prevalence and disparities in receiving medical advice to quit tobacco use in the US adult population. *Front Public Health*, 12, 1383060. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39376998>**

Klaiman, T, Farrell, N, Sheu, D, Belk, A, Silvestri, J, Kim, J et al. (2024). Use of tobacco during COVID-19: A qualitative study among medically underserved individuals. *PLoS One*, 19(8), e0308966. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39159172>

Vander Woude, CA, Buszkiewicz, JH, Travis, N, James, LN, & Fleischer, NL. (2024). Tobacco Control Policies and Sociodemographic Disparities in Cigarette Smoking Behaviors in the U.S.: A Systematic Review Protocol. *AJPM Focus*, 3(5), 100256. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39161402>

Long, SE, Lama, Y, & D'Angelo, H. (2023). Digital Communication Inequalities Among U.S. Adults Reporting Current Cigarette Use. *Am J Prev Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37793558>

Perez, GK, Rabin, JT, Tandon, M, Strauss, NM, Irwin, K, Philpotts, L et al. (2023). Do Tobacco Treatment Trials Address Disparities in Smoking Outcomes Among Black and Hispanic Cancer Patients? A Systematic Review of Smoking Cessation Interventions for Black and Hispanic Patients Diagnosed with Cancer. *J Racial Ethn Health Disparities*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37468742>

Teferra, AA, Wing, JJ, Lu, B, Xu, W, Roberts, ME, & Ferketich, AK. (2023). Examining trends in health care access measures among low-income adult smokers in Ohio: 2012-2019. *Prev Med Rep*, 31, 102106. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36820365>

Evins, E, & Cather, C. (2022). Addressing Barriers to Provision of First-Line Pharmacotherapy for Tobacco Use Disorder. [MS Top Pick]. *Psychiatr Serv*, appips20220467. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36349495>

Pisinger, C, Toxvaerd, CG, & Rasmussen, M. (2022). Smoking Cessation Programs Are Less Effective in Smokers with Low Socioeconomic Status Even When Financial Incentives for Quitting Smoking Are Offered-A Community-Randomized Smoking Cessation Trial in Denmark. *Int J Environ Res Public Health*, 19(17). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36078595>

Liu, J, Brighton, E, Tam, A, Godino, J, Brouwer, KC, Smoot, CB et al. (2021). Understanding health disparities affecting utilization of tobacco treatment in low-income patients in an urban health center in Southern California. *Prev Med Rep*, 24, 101541. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/34976615>

Zyambo, CM, Hendricks, PS, Cropsey, KL, Willig, JH, Wilson, CM, Gakumo, CA et al (2020). Racial disparities and factors associated with prescription for smoking cessation medications among smokers receiving routine clinical care for HIV. *AIDS Care*, 1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32530307>

Kock, L, & Brown, J. (2019). Commentary on Smith et al. (2019): Equity positive impact of English Stop Smoking Services underlines need for comprehensive approach to public health interventions. *Addiction*. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31677198>

Young-Wolff KC, Adams SR, Tan ASL, Adams AS, Klebaner D, et al. Disparities in knowledge and use of tobacco treatment among smokers in California following healthcare reform. *Preventive Medicine Reports*, 2019; 14:100847. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31024786>

van Wijk EC, Landais LL, and Harting J. Understanding the multitude of barriers that prevent smokers in lower socioeconomic groups from accessing smoking cessation support: A literature review. *Preventive Medicine*, 2019; 123:143-51. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30902700>

Smith CE, Hill SE, and Amos A. Impact of specialist and primary care stop smoking support on socioeconomic inequalities in cessation in the UK: A systematic review and national equity analysis. *Addiction*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31357250>

McQueen A, Roberts C, Garg R, Caburnay C, Fu Q, et al. Specialized tobacco quitline and basic needs navigation interventions to increase cessation among low income smokers: Study protocol for a randomized controlled trial. *Contemporary Clinical Trials*, 2019; 80:40-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30904596>

Smith C, Hill S, and Amos A. Stop smoking inequalities: A systematic review of socioeconomic inequalities in experiences of smoking cessation interventions in the UK. Cancer Research UK, 2018. Available from: [https://www.cancerresearchuk.org/sites/default/files/stop\\_smoking\\_inequalities\\_2018.pdf](https://www.cancerresearchuk.org/sites/default/files/stop_smoking_inequalities_2018.pdf).

Landrine H, Corral I, and Campbell KM. Racial disparities in healthcare provider advice to quit smoking. *Preventive Medicine Reports*, 2018; 10:172-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29868363>

Danan ER, Fu SS, Clothier BA, Noorbaloochi S, Hammett PJ, et al. The equity impact of proactive outreach to smokers: Analysis of a randomized trial. *American Journal of Preventive Medicine*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30139707>

Bailey SR, Heintzman J, Jacob RL, Puro J, and Marino M. Disparities in smoking cessation assistance in US primary care clinics. *American Journal of Public Health*, 2018:e1-e9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29927641>

Soulakova JN and Crockett LJ. Level of cigarette consumption and duration of smoking abstinence during failed quit attempts among long-term daily smokers: The role of race/ethnicity and cessation aids. *J Racial Ethn Health Disparities*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28444627>

Hooks-Anderson DR, Salas J, Secrest S, Skiodl-Hanlin S, and Scherrer JF. Association between race and receipt of counselling or medication for smoking cessation in primary care. *Family Practice*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29045650>

Edwards SA, Callaghan RC, Mann RE, and Bondy SJ. Association between socioeconomic status and access to care and quitting smoking with and without assistance. *Nicotine & Tobacco Research*, 2017; 20(1):40–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28340126>

Dahne J, Wahlquist AE, Garrett-Mayer E, Heckman BW, Michael Cummings K, et al. The differential impact of state tobacco control policies on cessation treatment utilization across established tobacco disparities groups. *Preventive Medicine*, 2017; 105:319–25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28987337>

Brose LS and McEwen A. Neighbourhood deprivation and outcomes of stop smoking support - an observational study. *PLoS ONE*, 2016; 11(1):e0148194. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26824352>

Wang L, Jin Y, Lu B, and Ferketich AK. A cross-country study of smoking cessation assistance utilization in 16 low and middle income countries: Data from the global adult Tobacco survey (2008-2012). *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26116084>

Paul C, Wolfenden L, Tzelepis F, Yoong S, Bowman J, et al. Nicotine replacement therapy as a smoking cessation aid among disadvantaged smokers: What answers do we need? *Drug and Alcohol Review*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26661119>

Brathwaite R, Addo J, Smeeth L, and Lock K. A systematic review of Tobacco smoking prevalence and description of Tobacco control strategies in sub-saharan african countries; 2007 to 2014. *PLoS ONE*, 2015; 10(7):e0132401. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26162085>

Bosdriesz JR, Willemsen MC, Stronks K, and Kunst AE. Socioeconomic inequalities in smoking cessation in 11 European countries from 1987 to 2012. *Journal of Epidemiology and Community Health*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25841241>

Kotz D. Rigorous methodoloy is needed to analyze and interpret observational data on the use and effectiveness of smoking cessation aids. *Nicotine & Tobacco Research*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25320111>

Clare P, Slade T, Courtney RJ, Martire KA, and Mattick RP. The use of smoking cessation and Quit support services by socioeconomic status over 10 years of the National Drug Strategy Household Survey. *Nicotine & Tobacco Research*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25320110>

Aspinall PJ and Mitton L. Smoking prevalence and the changing risk profiles in the UK ethnic and migrant minority populations: Implications for stop smoking services. *Public Health*, 2014; 128(3):297–306. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24612958>

Wang X and Abughosh S. Racial and ethnic disparity in smoking cessation medication use among adult smokers in the United States. *Value in Health*, 2013; 16(3):A241. Available from: [http://www.valueinhealthjournal.com/article/S1098-3015\(13\)01297-7/fulltext](http://www.valueinhealthjournal.com/article/S1098-3015(13)01297-7/fulltext)

Huang Y, Britton J, Hubbard R, and Lewis S. Who receives prescriptions for smoking cessation medications? An association rule mining analysis using a large primary care database. *Tobacco Control*, 2012; [Epub ahead of print]. Available from: <http://tobaccocontrol.bmjjournals.org/content/early/2012/01/13/tobaccocontrol-2011-050124.full>

Christiansen B, Reeder K, Hill M, Baker TB, and Fiore MC. Barriers to effective tobacco-dependence treatment for the very poor. *J Stud Alcohol Drugs*, 2012; 73(6):874-84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23036204>

Christiansen B, Reeder K, Hill M, Baker T, and Fiore M. Barriers to effective tobacco-dependence treatment for the very poor. *Journal of Studies on Alcohol and Drugs*, 2012; 73(6):874–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23036204>

Zhu S, Wong S, Stevens C, Nakashima D, and Gamst A. Use of a smokers' Quitline by Asian language speakers: Results from 15 years of operation in California. *American Journal of Public Health*, 2010; [Epub ahead of print]. Available from:

<http://ajph.aphapublications.org/cgi/reprint/AJPH.2009.168385v1?view=long&pmid=20299658>

Thornley S, Jackson G, McRobbie H, Sinclair S, and Smith J. Few smokers in South Auckland access subsidised nicotine replacement therapy. *The New Zealand Medical Journal*, 2010; 123(1308):16–27. Available from: <http://www.nzma.org.nz/journal/123-1308/3943/>

Willett J, Hood N, Burns E, Swetlick J, Wilson S, et al. Clinical faxed referrals to a tobacco Quitline reach, enrollment, and participant characteristics. *American Journal of Preventive Medicine*, 2009; 36(4):337–40. Available from: <http://www.ajpm-online.net/article/S0749-3797%2808%2901010-6/abstract>

Stuber J and Galea S. Who conceals their smoking status from their health care provider? *Nicotine & Tobacco Research*, 2009; 11(3):303–7. Available from: <http://ntr.oxfordjournals.org/content/11/3/303.long>

Solberg L, Parker E, Foldes S, and Walker P. Disparities in tobacco cessation medication orders and fills among special populations. *Nicotine & Tobacco Research*, 2009; 12(2):144–51. Available from: <http://ntr.oxfordjournals.org/content/12/2/144.long>

Miller CL and Sedivy V. Using a quitline plus low-cost nicotine replacement therapy to help disadvantaged smokers to quit. *Tobacco Control*, 2009; 18(2):144–9. Available from: <http://tobaccocontrol.bmjjournals.org/cgi/content/abstract/18/2/144>

Knudsen HK. Smoking cessation services in adolescent substance abuse treatment: Opportunities missed? *Journal of Drug Issues*, 2009; 39(2). Available from: <http://www2.criminology.fsu.edu/~jdi/39n2.htm>

Kaestle C and Wiles B. Targeting high-risk neighborhoods for tobacco prevention education in schools. *American Journal of Public Health*, 2009; 100(9):1708–13. Available from: <http://ajph.aphapublications.org/cgi/content/full/100/9/1708?view=long&pmid=20019323>

Chang F, Hu T, Lo S, Yu P, Chao K, et al. Quit smoking advice from health professionals in Taiwan: The role of funding policy and smokers{inverted exclamation}| socioeconomic status. *Tobacco Control*, 2009; 19:44–9. Available from: <http://tobaccocontrol.bmjjournals.org/content/19/1/44.full?sid=27a3840d-26fa-4c5c-acab-1ae919062be6>

Boles M, Rohde K, He H, Maher J, Stark M, et al. Effectiveness of a tobacco quitline in an indigenous population: A comparison between Alaska Native people and other first-time quitline callers who set a quit date. *International Journal of Circumpolar Health*, 2009; 68(2):170–81. Available from:

[http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list\\_uids=19517876](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19517876)

Sood A, Andoh J, Verhulst SJ, Rajoli N, and Hopkins-Price P. Characteristics of smokers calling a national reactive telephone helpline. *American Journal of Health Promotion*, 2008; 22(13):176–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18251117>

Sonnenfeld N, Schappert S, and Lin S. Racial and ethnic differences in delivery of tobacco-cessation services. *American Journal of Preventive Medicine*, 2008; 36(1):21–8. Available from: <http://www.ajpm-online.net/article/S0749-3797%2808%2900844-1/fulltext>

Reed M and Burns D. A population-based examination of racial and ethnic differences in receiving physicians' advice to quit smoking. *Nicotine & Tobacco Research*, 2008; 10(9):1487–94. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802323217>

Fu S, Kodl M, Joseph A, Hatsukami D, Johnson E, et al. Racial/ethnic disparities in the use of nicotine replacement therapy and quit ratios in lifetime smokers ages 25 to 44 years. *Cancer Epidemiology Biomarkers & Prevention*, 2008; 17(7):1640–7. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2593846/>

Burns E and Levinson A. Discontinuation of nicotine replacement therapy among smoking-cessation attempters. *American Journal of Preventive Medicine*, 2008; 34(3):212–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18312809>

Browning K, Ferketich A, Salsberry P, and Wewers M. Socioeconomic disparity in provider-delivered assistance to quit smoking. *Nicotine & Tobacco Research*, 2008; 10(1):55–61. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200701704905>

Andoh J, Verhulst S, Ganesh M, Hopkins-Price P, Edson B, et al. Sex- and race-related differences among smokers using a national helpline are not explained by socioeconomic status. *Journal of the National Medical Association*, 2008; 100(2):200–7. Available from: [www.nmanet.org/images/uploads/Journal/OC200.pdf](http://www.nmanet.org/images/uploads/Journal/OC200.pdf)

Siahpush M, Wakefield M, Spittal M, and Durkin S. Anti-smoking television advertising and socio-economic variations in calls to Quitline. *Journal of Epidemiology and Community Health*, 2007; 61(4):298–301. Available from: <http://jech.bmjjournals.org/cgi/content/full/61/4/298>

Lillard DR, Plassmann V, Kenkel D, and Mathios A. Who kicks the habit and how they do it: Socioeconomic differences across methods of quitting smoking in the USA. *Social Science & Medicine*, 2007; 64(12):2504–19. Available from: [http://www.sciencedirect.com/science?\\_ob=ArticleURL&\\_udi=B6VBF-4NDVHDN-1&\\_user=10&\\_rdoc=1&\\_fmt=&\\_orig=search&\\_sort=d&view=c&\\_acct=C000050221&\\_version=1&\\_urIVersion=0&\\_userid=10&md5=9c995358ca89698ee32e3f83db6ee0be](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6VBF-4NDVHDN-1&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&view=c&_acct=C000050221&_version=1&_urIVersion=0&_userid=10&md5=9c995358ca89698ee32e3f83db6ee0be)

Elin R, Nick R, Alison C, Sarah L, and John B. Use of nicotine replacement therapy in socioeconomically deprived young smokers: A community-based pilot randomised controlled trial. *Tobacco Control*, 2006; 15(5):373–6. Available from: <http://ejournals.ebsco.com/direct.asp?ArticleID=47B3970FE1458C036B27>

Chapman S and Liberman J. Ensuring smokers are adequately informed: Reflections on consumer rights, manufacturer responsibilities, and policy implications. *Tobacco Control*, 2005; 14(suppl. 2):ii8–ii13. Available from: [http://tc.bmjjournals.com/cgi/content/abstract/14/suppl\\_2/ii8](http://tc.bmjjournals.com/cgi/content/abstract/14/suppl_2/ii8)

Burns D and Warner K. Smokers who have not Quit: Is cessation more difficult and should we change our strategies, in those who continue to smoke. *Smoking and Tobacco Control Monograph no.15*. Bethesda, MD: US Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute, 2003. Available from: <http://cancercontrol.cancer.gov/tcrb/monographs/15/monograph15-chapter1.pdf>.

Liberman J and Clough J. Corporations that kill: The criminal liability of tobacco manufacturers. *Criminal Law Journal*, 2002; 26(4):223. Available from: [http://www.vctc.org.au/tc-res/criminal\\_liability\\_paper.pdf](http://www.vctc.org.au/tc-res/criminal_liability_paper.pdf)

Borland R, Pigott R, Rintoul D, Shore S, and Young S. Barriers to access of smoking cessation programs, nicotine replacement therapy and other pharmacotherapies for the general Australian population and at risk population groups. Literature review and volumes 1 and 2. Canberra: Final Report to Commonwealth Department of Health and Ageing, 2002.

Chapman S and Wakefield M. Tobacco control advocacy in Australia: Reflections on 30 years of progress. *Health Education & Behavior*, 2001; 28(3):274–89. Available from: <http://heb.sagepub.com/cgi/content/abstract/28/3/274>

Carter S, Borland R, and Chapman C. Finding the strength to kill your best friend—smokers talk about smoking and quitting. Sydney: Australian Smoking Cessation Consortium and GlaxoSmithKline Consumer Healthcare, 2001. Available from: <http://tobacco.health.usyd.edu.au/site/supersite/resources/pdfs/killbestfriend.pdf>.

Royal College of Physicians of London. Nicotine addiction in britain. A report of the Tobacco advisory group of the royal college of physicians. London: Royal College of Physicians of London, 2000. Available from: <http://bookshop.rcplondon.ac.uk/details.aspx?e=131>.

Chapman S, Wilson D, and Wakefield M. Smokers' understanding of cigarette yield labels. *Medical Journal of Australia*, 1986; 145(8):376–9.

### 9.9.1 *Quitlines*

**Monshi, SS, Alqahtani, MM, Alangari, AS, Sinky, TH, Al-Zalabani, AH, & Alanazi, AM. (2023). The association between cultural and social occasions and smoking cessation: The case of Saudi Arabia. *Tob Induc Dis*, 21, 153. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38026497>**

Kreuter, MW, Garg, R, Fu, Q, Caburnay, C, Thompson, T, Roberts, C et al (2023). Helping low-income smokers quit: findings from a randomized controlled trial comparing specialized quitline services with and without social needs navigation. *Lancet Reg Health Am*, 23, 100529. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37408953>

Griffin JL, Segal KS, and Nahvi S. Barriers to telephone Quitline use among methadone-maintained smokers. *Nicotine & Tobacco Research*, 2015; 17(8):931–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26180217>

Varghese M, Sheffer C, Stitzer M, Landes R, Brackman SL, et al. Socioeconomic disparities in telephone-based treatment of tobacco dependence. *American Journal of Public Health*, 2014; 104(8):e76–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24922165>

Campbell HS, Baskerville NB, Hayward LM, Brown KS, and Ossip DJ. The reach ratio--a new indicator for comparing quitline reach into smoking subgroups. *Nicotine & Tobacco Research*, 2014; 16(4):491–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24311698>

### 9.9.2 Disparities in use of treatments for tobacco dependence

**Rosenthal, SR, Wensley, IA, & Noel, JK. (2022). Eliminating Disparities in Young Adult Tobacco Use: The Need for Integrated Behavioral Healthcare. *R I Med J (2013)*, 105(2), 38-42. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35211709>**

Coughlin, SS, Vernon, M, Majeed, B, Clary, C Moore, J., Islam, KM, & Tingen, MS. (2020). Tobacco Cessation, Rural Residence, and Lung Cancer. *J Environ Health Sci*, 6(1), 1-4. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/33163626>

van Straaten, B, Meerkerk, GJ, van den Brand, FA, Lucas, P, de Wit, N, & Nagelhout, GE. (2020). How can vulnerable groups be recruited to participate in a community-based smoking cessation program and perceptions of effective elements: A qualitative study among participants and professionals. *Tob Prev Cessat*, 6, 64. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/33241164>

Dahne, J, Wahlquist, AE, Smith, TT, & Carpenter, MJ. (2020). The differential impact of nicotine replacement therapy sampling on cessation outcomes across established tobacco disparities groups. *Prev Med*, 136, 106096. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32320705>

Goodin, A, Talbert, J, Freeman, PR, Hahn, EJ, & Fallin-Bennett, A. (2020). Appalachian disparities in tobacco cessation treatment utilization in Medicaid. *Substance Abuse Treatment, Prevention and Policy*, 15(1), 5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31959212>

Fillon M. Study evaluates mobile telephone interventions for socioeconomically disadvantaged smokers. *CA: A Cancer Journal for Clinicians*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30856675>

Zhu SH, Anderson CM, Zhuang YL, Gamst AC, and Kohatsu ND. Smoking prevalence in medicaid has been declining at a negligible rate. *PLoS ONE*, 2017; 12(5):e0178279. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28542637>

Wilson A, Guillaumier A, George J, Denham A, and Bonevski B. A systematic narrative review of the effectiveness of behavioural smoking cessation interventions in selected disadvantaged groups (2010-2017). *Expert Rev Respir Med*, 2017; 11(8):617-30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28608758>

Sorensen G, Pednekar M, Cordeira LS, Pawar P, Nagler EM, et al. Effects of a worksite tobacco control intervention in India: The mumbai worksite tobacco control study, a cluster-randomised trial. *Tobacco Control*, 2017; 26(2):210-6. Available from: <http://tobaccocontrol.bmjjournals.org/content/tobaccocontrol/26/2/210.full.pdf>

Hammett P, Fu SS, Nelson D, Clothier B, Saul JE, et al. A proactive smoking cessation intervention for socioeconomically disadvantaged smokers: The role of smoking-related stigma. Nicotine & Tobacco Research, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28398492>

Flocke SA, Hoffman R, Eberth JM, Park H, Birkby G, et al. The prevalence of Tobacco use at federally qualified health centers in the United States, 2013. Preventing Chronic Disease, 2017; 14:E29. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28384096>

Feinberg A, Lopez PM, Wyka K, Islam N, Seidl L, et al. Prevalence and correlates of smoking among low-income adults residing in New York city public housing developments-2015. J Urban Health, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28656541>

Kostandinov O. Community confronting high smoking rates among disadvantaged Quit Resource Centre, 2015. Available from: <http://www.quit.org.au/resource-centre/media-advertising/media-releases/community-confronting-high-smoking-rates-among-disadvantaged.html>.

O'Brien J, Salmon AM, and Penman A. What has fairness got to do with it? Tackling tobacco among Australia's disadvantaged. Drug and Alcohol Review, 2012; 31(5):723–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22524309>

Bobo JK, Anderson JR, and Bowman A. Training chemical dependency counselors to treat nicotine dependence. Addictive Behaviors, 1997; 22(1):23–30. Available from: [http://www.sciencedirect.com/science?\\_ob=ArticleURL&\\_udi=B6VC9-3SX4Y19-3&\\_user=10&\\_rdoc=1&\\_fmt=&\\_orig=search&\\_sort=d&view=c&\\_acct=C000050221&\\_version=1&\\_urIVersion=0&\\_userid=10&md5=22b8a6d61b8ca86e042c714e626abd7c](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6VC9-3SX4Y19-3&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&view=c&_acct=C000050221&_version=1&_urIVersion=0&_userid=10&md5=22b8a6d61b8ca86e042c714e626abd7c)

#### *9.9.2.1 Disparities in use of treatments*

Tseng, TS, Celestin, MD, Jr, Yu, Q, Li, M, Luo, T, & Moody-Thomas, S. (2021). Use of Geographic Information System Technology to Evaluate Health Disparities in Smoking Cessation Class Accessibility for Patients in Louisiana Public Hospitals. *Front Public Health*, 9, 712635. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/34476230>

#### *9.9.2.2 Disparities in compliance with treatment*

Gajos, JM, Hawes, ES, Chana, SM, Mrug, S, Wolford-Clevenger, C, Businelle, MS et al. (2022). Daily adherence to nicotine replacement therapy in low-income smokers: The role of gender, negative mood, motivation, and self-efficacy. [MS Top Pick]. *Addict Behav*, 138, 107543. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36423571>

#### *9.9.3 Disparities in provision of quit smoking advice and referral by general practitioners*

**Upadhyay, S, & Jones, J. (2024). Patient-provider communication quality: Socioeconomic disparities in smoking outcomes. *Tob Prev Cessat*, 10.** Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38469412>

Manns, A, Mahdjoub, S, Ibanez, G, Jarrier, E, Daeipour, A, Melchior, M, & El-Khoury, F. (2023). Health professional's perception of a smoking cessation intervention among disadvantaged patients

participating in a pragmatic randomized trial. *BMC Health Serv Res*, 23(1), 993. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37710246>

Braciszewski, JM, Sala-Hamrick, KJ, Zelenak, L, Gootee, J, Elsiss, F, Ottolini, J et al. (2022). Reducing Smoking Cessation Disparities: Capacity for a Primary Care- and Technology-Based Approach Among Medicaid Recipients. [MS Top Pick]. *J Clin Psychol Med Settings*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36400987>

Maki, KG., & Volk, RJ. (2022). Disparities in Receipt of Smoking Cessation Assistance Within the US. *JAMA Netw Open*, 5(6), e2215681. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35648402>

Sipe, CJ, Koopmeiners, JS, Donny, EC, Hatsukami, DK, & Murphy, SE. (2020). UGT2B10 Genotype Influences Serum Cotinine Levels and Is a Primary Determinant of Higher Cotinine in African American Smokers. *Cancer Epidemiol Biomarkers Prev*. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32532831>

Angus C, Brown J, Beard E, Gillespie D, Buykx P, et al. Socioeconomic inequalities in the delivery of brief interventions for smoking and excessive drinking: Findings from a cross-sectional household survey in England. *BMJ Open*, 2019; 9(4):e023448. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31048422>

#### *9.9.4 Harm reduction: a strategy of benefit for disadvantaged groups?*

Lucherini, M, Hill, S, & Smith, K. (2019). Potential for non-combustible nicotine products to reduce socioeconomic inequalities in smoking: a systematic review and synthesis of best available evidence. [S ]. *BMC Public Health*, 19(1), 1469. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31694602>

## **News reports:**

No authors listed. Medicaid tobacco cessation: Big gaps remain in efforts to get smokers to quit. Medical News Today, 2016. Available from: <http://www.medicalnewstoday.com/releases/304728.php?tw>

Medicare Australia. Pbs statistics. Canberra: Medicare Australia, 2008. Last update: Viewed Available from: <http://www.medicareaustralia.gov.au/provider/pbs/stats.shtml>.

#### *9.9.1 Quitlines*

#### *9.9.2 Disparities in use of treatments for tobacco dependence*

Butt C. Quit Victoria to help tongans give up smoking. The Age, 2015. Available from:  
<http://www.theage.com.au/victoria/quit-victoria-to-help-tongans-give-up-smoking-20150329-1ma7fr.html>

Partos T and Borland R. Data file of responses to eighth wave of the international Tobacco control four-country survey, in *Provided to Merryn Pearce of the Tobacco Control Unit* 2012, Cancer Council Victoria.

*9.9.2.1 Disparities in use of treatments*

*9.9.2.2 Disparities in compliance with treatment*

*9.9.3 Disparities in provision of quit smoking advice and referral by general practitioners*

*9.9.4 Harm reduction: a strategy of benefit for disadvantaged groups?*