

# Tobacco in Australia

## Facts & Issues

---

### Relevant news and research

#### 9A.1 People living in regional and remote areas of Australia

*Last updated November 2022*

Research:.....	1
9A.1 People living in regional and remote areas of Australia.....	1
9A.1.1 Trends in the prevalence of smoking .....	6
9A.1.2 Contribution of smoking to health outcomes and social inequality .....	6
9A.1.3 Interventions for reducing smoking .....	6
News reports:.....	8
9A.1 People living in regional and remote areas of Australia.....	8
9A.1.1 Trends in the prevalence of smoking .....	8
9A.1.2 Contribution of smoking to health outcomes and social inequality .....	8
9A.1.3 Interventions for reducing smoking .....	8

#### Research:

##### *9A.1 People living in regional and remote areas of Australia*

Azagba, S, Ebling, T, & Shan, L. (2022). A double disparity: Rural sexual minorities and tobacco use among U.S. adults. *Addict Behav*, 137, 107527. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/36308838>

tobaccoinaustralia.org.au

Dorilas E, Hill SC, and Pesko MF. Tobacco surcharges associated with reduced aca marketplace enrollment. *Health Aff (Millwood)*, 2022; 41(3):398-405. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35254928>

Consultation draft national tobacco strategy 2022-2030. Canberra: Commonwealth of Australia, 2022. Available from: [https://consultations.health.gov.au/atodb/national-tobacco-strategy-2022-2030/supporting\\_documents/Draft%20NTS%2020222030%20for%20consultaion%20hub.pdf](https://consultations.health.gov.au/atodb/national-tobacco-strategy-2022-2030/supporting_documents/Draft%20NTS%2020222030%20for%20consultaion%20hub.pdf).

Darker C, Loudon K, O'Connell N, Castello S, Burke E, et al. An application of precis-2 to evaluate trial design in a pilot cluster randomised controlled trial of a community-based smoking cessation intervention for women living in disadvantaged areas of ireland. *Pilot Feasibility Stud*, 2022; 8(1):19. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35078530>

Yuan L, Zhao Z, Wang J, Du M, Xiao Y, et al. Decomposition and comparative analysis of the prevalence of and factors associated with smoking between the rural and urban elderly population in China: A national cross-sectional study. *Front Public Health*, 2022; 10:824587. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35372230>

Khan A, Green K, Smoll N, Khandaker G, Gartner C, et al. Roles, experiences, and perspectives of the stakeholders of "10,000 lives" smoking cessation initiative in central queensland: Findings from an online survey during COVID-19 situation. *Health Promotion Journal of Australia*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35322498>

Mitchell SA, Woods-Giscombe C, Kneipp SM, Beeber LS, and Kulbok PA. Social determinants of smoking in women from low-income rural backgrounds: Findings from a photovoice study. *Archives of Psychiatric Nursing*, 2021; 35(1):56-65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33593516>

Ozga JE, Romm KF, Turiano NA, Douglas A, Dino G, et al. Cumulative disadvantage as a framework for understanding rural tobacco use disparities. *Experimental and Clinical Psychopharmacology*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34014742>

Australian Institute of Health and Welfare. Australian burden of disease study: Impact and causes of illness and death in Australia 2018. Canberra: AIHW, Australian Government 2021. Available from: <https://www.aihw.gov.au/reports/burden-of-disease/abds-impact-and-causes-of-illness-and-death-in-aus/summary>.

Khan A, Green K, Khandaker G, Lawler S, and Gartner C. How can a coordinated regional smoking cessation initiative be developed and implemented? A programme logic model to evaluate the '10,000 lives' health promotion initiative in central queensland, Australia. *BMJ Open*, 2021; 11(3):e044649. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33771827>

Talbot JA, Ziller EC, and Milkowski CM. Use of electronic health records to manage tobacco screening and treatment in rural primary care. *Journal of Rural Health*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34468036>

Yau MTK and Okoli CTC. Associations of tobacco use and consumption with rurality among patients with psychiatric disorders: Does smoke-free policy matter? *Journal of Rural Health*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34339080>

Australian Bureau of Statistics. Pandemic insights into Australian smokers, 2020-21. 2021. Available from: <https://www.abs.gov.au/articles/pandemic-insights-australian-smokers-2020-21>.

Dong X, Ding M, Chen W, Liu Z, and Yi X. Relationship between smoking, physical activity, screen time, and quality of life among adolescents. *International Journal of Environmental Research and Public Health*, 2020; 17(21). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33142847>

Roberts ME, Doogan NJ, Tanenbaum E, Stillman FA, Mumford EA, et al. How should we define "rural" when investigating rural tobacco use in the United States? *Substance Abuse*, 2020:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33320797>

Australian Institute for Health and Welfare. Australia's health 2020. 2020. Available from: <https://www.aihw.gov.au/reports-data/australias-health>

Australian Institute for Health and Welfare. Rural and remote health. 2020. Available from: <https://www.aihw.gov.au/reports/australias-health/rural-and-remote-health>.

Byaruhanga J, Paul CL, Wiggers J, Byrnes E, Mitchell A, et al. Connectivity of real-time video counselling versus telephone counselling for smoking cessation in rural and remote areas: An exploratory study. *International Journal of Environmental Research and Public Health*, 2020; 17(8). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32331356>

Lum A, Skelton E, McCarter KL, Handley T, Judd L, et al. Smoking cessation interventions for people living in rural and remote areas: A systematic review protocol. *BMJ Open*, 2020; 10(11):e041011. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33208333>

Ramsey AT, Baker TB, Pham G, Stoneking F, Smock N, et al. Low burden strategies are needed to reduce smoking in rural healthcare settings: A lesson from cancer clinics. *International Journal of Environmental Research and Public Health*, 2020; 17(5). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32155775>

Smith PM, Seamark LD, and Beck K. Integration of an evidence-based tobacco cessation program into a substance use disorders program to enhance equity of treatment access for northern, rural, and remote communities. *Transl Behav Med*, 2020; 10(3):555-64. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32766869>

US Department of Health and Human Services. Smoking cessation. A report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, 2020. Available from: <https://www.hhs.gov/sites/default/files/2020-cessation-sgr-full-report.pdf>.

Byaruhanga J, Wiggers J, Paul CL, Byrnes E, Mitchell A, et al. Acceptability of real-time video counselling compared to other behavioural interventions for smoking cessation in rural and remote areas. *Drug and Alcohol Dependence*, 2020; 217:108296. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32980788>

Australian Institute of Health and Welfare. Alcohol and other drug use in regional and remote Australia: Consumption, harms and access to treatment 2016–17. Cat. no. HSE 212. Canberra: AIHW, 2019. Available from: <https://www.aihw.gov.au/getmedia/78ea0b3d-4478-4a1f-a02a-3e3b5175e5d8/aihw-hse-212.pdf.aspx?inline=true>.

Australian Institute of Health and Welfare. Rural & remote health. Canberra: AIHW, 2019. Available from: <https://www.aihw.gov.au/reports/rural-remote-australians/rural-remote-health>.

Wakerman J and Humphreys JS. “Better health in the bush”: Why we urgently need a national rural and remote health strategy. *Medical Journal of Australia*, 2019; 210(5):202-3.e1. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.5694/mja2.50041>

Buettner-Schmidt K, Miller DR, and Maack B. Disparities in rural tobacco use, smoke-free policies, and tobacco taxes. *West J Nurs Res*, 2019:193945919828061. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30774036>

Byaruhanga J, Tzelepis F, Paul C, Wiggers J, Byrnes E, et al. Cost per participant recruited from rural and remote areas into a smoking cessation trial via online or traditional strategies: Observational study. *Journal of Medical Internet Research*, 2019; 21(11):e14911. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31714253>

Poltyn-Zaradna K, Zatonska K, Basiak A, Sozanska B, Gawel-Dabrowska D, et al. Sociodemographic characteristic of changes in smoking patterns in rural and urban population of pure Poland study: Findings from 6-year follow up. *BMC Public Health*, 2019; 19(1):6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30606160>

Gay C, Clements-Nolle K, Packham J, Ackerman G, Lensch T, et al. Community-level exposure to the rural mining industry: The potential influence on early adolescent alcohol and tobacco use. *Journal of Rural Health*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29388274>

Hinton A, Nagaraja HN, Cooper S, and Wewers ME. Tobacco product transition patterns in rural and urban cohorts: Where do dual users go? *Preventive Medicine Reports*, 2018; 12:241-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30377574>

Hall MB and Vos P. Comparison of cancer fatalism among rural smokers and nonsmokers. *Journal of Community Health*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30267351>

Nighbor TD, Doogan NJ, Roberts ME, Cepeda-Benito A, Kurti AN, et al. Smoking prevalence and trends among a u.S. National sample of women of reproductive age in rural versus urban settings. *PLoS One*, 2018; 13(11):e0207818. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30485376>

Noonan D, Silva S, Njuru J, Bishop T, Fish LJ, et al. Feasibility of a text-based smoking cessation intervention in rural older adults. *Health Education Research*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29309599>

Tzelepis F, Wiggers J, Paul CL, Byaruhanga J, Byrnes E, et al. A randomised trial of real-time video counselling for smoking cessation in regional and remote locations: Study protocol. *Contemporary Clinical Trials*, 2018; 74:70-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30290277>

[https://www.contemporaryclinicaltrials.com/article/S1551-7144\(18\)30401-4/fulltext](https://www.contemporaryclinicaltrials.com/article/S1551-7144(18)30401-4/fulltext)

Pesko MF and Roberts AMT. Adolescent tobacco use in urban versus rural areas of the United States: The influence of Tobacco Control policy environments. *The Journal of Adolescent Health*, 2017; 61(1):70-6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28363718>

Zhi K, Wang L, Han Y, Vaughn MG, Qian Z, et al. Trends in cigarette smoking among older male adults in China: An urban-rural comparison. *J Appl Gerontol*, 2017;733464817716967. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28653557>

Haines HM, Cynthia O, Pierce D, and Bourke L. Notwithstanding high prevalence of overweight and obesity, smoking remains the most important factor in poor self-rated health and hospital use in an Australian regional community. *AIMS Public Health*, 2017; 4(4):402-17. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29546226>

Skinner A, Havard A, Tran DT, and Jorm LR. Access to subsidized smoking cessation medications by Australian smokers aged 45 years and older: A population-based cohort study. *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27613898>

Yang L, Liu Y, Wang J, and Jia C. Prevalence and risk factors associated with hardcore smoking among rural daily male smokers. *American Journal on Addictions*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27706867>

Tall JA, Brew BK, Saurman E, and Jones TC. Implementing an anti-smoking program in rural-remote communities: Challenges and strategies. *Rural Remote Health*, 2015; 15(4):3516. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26530272>

Standing Council on Health, National strategic framework for rural and remote health Pandora electronic collection., ed., Australia R and Regional Health A. [Canberra]: [Rural and Regional Health Australia]; 2012. Available from: <http://nla.gov.au/nla.arc-149260>.

Johnston V and Thomas DP. What works in indigenous tobacco control? The perceptions of remote indigenous community members and health staff. *Health Promotion Journal of Australia*, 2010; 21(1):45–50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20406152>

Bailey B and Cole L. Rurality and birth outcomes: Findings from southern appalachia and the potential role of pregnancy smoking. *The Journal of Rural Health*, 2009; 25(2):141–9. Available from: <http://www3.interscience.wiley.com/user/accessdenied?ID=122295777&Act=2138&Code=4719&Page=/cgi-bin/fulltext/122295777/HTMLSTART>

Li FX, Robson PJ, Ashbury FD, Hatcher J, and Bryant HE. Smoking frequency, prevalence and trends, and their socio-demographic associations in alberta, canada. *Canadian Journal of Public Health*, 2009; 100(6):453-8. Available from: <https://pubmed.ncbi.nlm.nih.gov/20209740>

Chapman A, Bunker S, Dunbar J, Philpot B, McNamara K, et al. Rural smokers—a prevention opportunity. *Australian Family Physician*, 2009; 38(5):352–6. Available from: <https://www.racgp.org.au/afp/2009/may/rural-smokers>

Hutcheson TD, Greiner KA, Ellerbeck EF, Jeffries SK, Mussulman LM, et al. Understanding smoking cessation in rural communities. *Journal of Rural Health*, 2008; 24(2):116-24. Available from: <https://pubmed.ncbi.nlm.nih.gov/18397444/>

Tracey E, Baker D, Chen W, Stavrou E, and Bishop J. Cancer in new south wales: Incidence, mortality and prevalence report 2005. Sydney: The Cancer Institute New South Wales, 2007. Available from: <http://www.cancercouncil.com.au/editorial.asp?pageid=9>.

Ministerial Council on Drug Strategy. Australian national tobacco strategy 2004-2009. Canberra: Department of Health and Ageing, 2005. Available from: [https://untobaccocontrol.org/impldb/wp-content/uploads/reports/Australia\\_annex7\\_national\\_tobacco\\_strategy2004\\_2009.pdf](https://untobaccocontrol.org/impldb/wp-content/uploads/reports/Australia_annex7_national_tobacco_strategy2004_2009.pdf).

Ministerial Council on Drug Strategy. Meeting the challenges of the next five years-5&6: Ideas and resources for addressing inequity in smoking-related harms by addressing social, cultural and economic determinants, and by tailoring to ensure access for disadvantaged groups National Tobacco Strategy, supporting documents, Canberra: Department of Health and Ageing, 2005. Available from: <http://www.health.gov.au/internet/wcms/publishing.nsf/Content/phd-pub-tobacco-tobccstrat2-cnt.htm/>.

Australian Institute of Health and Welfare. Health in rural and remote Australia. AIHW cat. no. PHE 6. Canberra: AIHW, 1998. Available from: <http://www.aihw.gov.au/publications/health/hrra/index.html>.

#### *9A.1.1 Trends in the prevalence of smoking*

**Parker, MA, Weinberger, AH, Eggers, EM, Parker, ES, & Villanti, AC. (2022). Trends in Rural and Urban Cigarette Smoking Quit Ratios in the US From 2010 to 2020. *JAMA Netw Open*, 5(8), e2225326. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35921112>**

National Preventative Health Taskforce. Australia: The healthiest country by 2020. National Preventative Health Strategy, Canberra: Commonwealth of Australia 2009. Available from: [https://www.health.qld.gov.au/\\_data/assets/pdf\\_file/0022/424426/nphs-overview.pdf](https://www.health.qld.gov.au/_data/assets/pdf_file/0022/424426/nphs-overview.pdf).

#### *9A.1.2 Contribution of smoking to health outcomes and social inequality*

**Thomson, B, Emberson, J, Lacey, B, Lewington, S, Peto, R, Jemal, A, & Islami, F. (2022). Association Between Smoking, Smoking Cessation, and Mortality by Race, Ethnicity, and Sex Among US Adults. *JAMA Netw Open*, 5(10), e2231480. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36279139>**

Hendi, AS, & Ho, JY. (2022). Smoking and the widening inequality in life expectancy between metropolitan and nonmetropolitan areas of the United States. *Front Public Health*, 10, 942842. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36159248>

#### *9A.1.3 Interventions for reducing smoking*

**Lee, SH, Yi, YH, Lee, YI, Lee, HY, & Lim, KM. (2022). Factors associated with long-term smoking relapse in those who succeeded in smoking cessation using regional smoking cessation programs. *Medicine (Baltimore)*, 101(31), e29595. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35945709>**

Little, MA, Porter, KJ, Pebley, K, Wiseman, KP, Cohn, W, Anderson, RT, & Krukowski, RA. (2022). Evaluating the feasibility of pharmacist-facilitated tobacco cessation interventions in independent community pharmacies in rural Appalachia. *J Am Pharm Assoc (2003)*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35953377>

Vance, L, Glanville, B, Ramkumar, K, Chambers, J, & Tzelepis, F. (2022). The effectiveness of smoking cessation interventions in rural and remote populations: Systematic review and meta-analyses. *Int J Drug Policy*, 106, 103775. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35772266>

Patten, C, Brockman, T, Kelpin, S, Sinicrope, P, Boehmer, K, St Sauver, J et al. (2022). Interventions for Increasing Digital Equity and Access (IDEA) among rural patients who smoke: Study protocol for a pragmatic randomized pilot trial. *Contemp Clin Trials*, 106838. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35760340>

## News reports:

*9A.1 People living in regional and remote areas of Australia*

*9A.1.1 Trends in the prevalence of smoking*

*9A.1.2 Contribution of smoking to health outcomes and social inequality*

*9A.1.3 Interventions for reducing smoking*