

Tobacco in Australia

Facts & Issues

Relevant news and research

9A.3 People with substance use and mental disorders

Last updated December 2024

Research:	2
9A.3 People with substance use and mental disorders	2
9A.3.1 Prevalence of smoking among people with mental health problems	50
9A.3.1.2 Smoking prevalence over time	58
9A.3.2 Contribution of smoking to health outcomes and social inequality	60
9A.3.3 Explanations for higher smoking prevalence	67
9A.3.4 Barriers to be overcome when quitting	77
9A.3.5 Interventions for reducing smoking	84
9A.3.5.1 Depression	103
9A.3.5.2 Anxiety	117
9A.3.5.3 Suicidal ideation	121
9A.3.5.4 Attention-deficit/hyperactivity disorder (ADHD)	121
9A.3.5.5 Substance use disorders	123
9A.3.5.6 Post-traumatic stress disorder (PTSD)	155
9A.3.5.7 Severe mental illness, including bipolar disorder and schizophrenia	159
9A.3.5.7.1 Bipolar disorder	164
9A.3.5.7.1 Schizophrenia	166
9A.3.6 The role of health professionals and health settings	178
News:	194
9A.3 People with substance use and mental disorders	194
9A.3.1 Prevalence of smoking among people with mental health problems	194
9A.3.1.2 Smoking prevalence over time	194
9A.3.2 Contribution of smoking to health outcomes and social inequality	194

9A.3.3 Explanations for higher smoking prevalence.....	194
9A.3.4 Barriers to be overcome when quitting.....	194
9A.3.5 Interventions for reducing smoking	194
9A.3.5.1 Depression	195
9A.3.5.2 Anxiety	195
9A.3.5.3 Suicidal ideation.....	195
9A.3.5.4 Attention-deficit/hyperactivity disorder (ADHD)	195
9A.3.5.5 Substance use disorders	195
9A.3.5.6 Post-traumatic stress disorder (PTSD).....	195
9A.3.5.7 Severe mental illness, including bipolar disorder and schizophrenia	195
9A.3.6 The role of health professionals and health settings	195

Research:

9A.3 People with substance use and mental disorders

Szerman, N, Parro, C, Vega, P, Basurte-Villamor, I, & Ruiz-Veguilla, M. (2024). Tobacco use disorder in patients with other mental disorders: a dual disorder perspective from clinical neuroscience. *Front Psychiatry*, 15, 1427561. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39465048>

Qiu, Y, & Ma, X. (2024). Using machine learning models to identify the risk of depression in middle-aged and older adults with frequent and infrequent nicotine use: A cross-sectional study. *J Affect Disord*, 367, 554-561. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39222853>

Nagawa, CS, Rigotti, NA, Chang, Y, Levy, DE, Streck, JM, Ylioja, T et al. (2024). Association between Smoking Abstinence and Depression and Anxiety Symptoms After Hospital Discharge: The Helping HAND 4 Trial. *J Addict Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39115217>

Parul, P, Joseph, B, Datta, S, & Rahman, MA. (2024). Correlates of Tobacco Use Among People with Mental Illness Within Asia: A Scoping Review. *Community Ment Health J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39126555>

Encinosa, W, Bernard, D, & Valdez, RB. (2024). Heavy vs Light Smoking: Its Association with Opioid Use, Chronic Pain, and Mental Health. *Am J Prev Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39025246>

Guo, Y, & Yan, J. (2024). Association between tobacco smoke exposure and depression: the NHANES 2005-2018 and Mendelian randomization study. *Arch Public Health*, 82(1), 100. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38961510>

Parro-Torres, C, Casas, M, Martinez-Raga, J, Pinet Ogue, MC, Saiz Martinez, PA, & Szerman, N. (2024). Tobacco use disorder and other mental disorders: The neglected dual disorder. *Span J Psychiatry Ment Health*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38740329>

Bover Manderski, MT, Ganz, O, Chen-Sankey, J, Villanti, AC, & Delnevo, CD. (2023). Evaluating Correlations Between Premium Cigar Smoking and Mental Health and Substance Use Dependence Conditions Among U.S. Adults, 2010-2019. *Nicotine Tob Res*, 25(Suppl_1), S94-S101. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37506237>

Campion, J, Johnston, G, Shiers, D, & Chew-Graham, C. (2023). Why should we prioritise smoking cessation for people with mental health conditions? *Br J Gen Pract*, 73(731), 251-253. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37230792>

White, AM, Barnes, AJ, & Garner, W. (2023). Racial/ethnic differences in the association between menthol cigarette use and mental illness among adults who smoke in the United States. *J Ethn Subst Abuse*, 1-21. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37052141>

Lee, B, Levy, D, & Seo, DC. (2023). Underlying patterns of the co-occurrence of tobacco use and mental health among youth. *J Behav Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36637734>

Vong, V, Simpson-Yap, S, Phaiju, S, Davenport, RA, Neate, SL, Pisano, MI, & Reece, JC. (2023). The association between tobacco smoking and depression and anxiety in people with multiple sclerosis: A systematic review. *Mult Scler Relat Disord*, 70, 104501. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36621161>

Mason, TB, Dolgon-Krutolow, A, Smith, KE, & Leventhal, AM. (2022). Body Dissatisfaction and Binge Eating: The Moderating Roles of Sweet Taste Reward Sensitivity and Dietary Restraint among Tobacco Product Users. *Int J Environ Res Public Health*, 19(23). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36497598>

Rosen, RL, Ramasubramani, RS, Benowitz, NL, Gandhi, KK, & Williams, JM. (2023). Caffeine levels and dietary intake in smokers with schizophrenia and bipolar disorder. *Psychiatry Res*, 319, 114989. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36470161>

Wang, DM, Xia, LY, Zhou, HX, Tian, Y, Dai, QL, Xiu, MH et al. (2023). Smoking affects the association between cognitive impairment and P50 inhibition defects in patients with chronic schizophrenia: A case-control study. *Asian J Psychiatry*, 79, 103391. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36516649>

Xi, Q., Meza, R., Leventhal, A., & Tam, J. (2022). Modeling cigarette smoking disparities between people with and without serious psychological distress in the US, 1997-2100. *Prev Med*, 166, 107385. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36495925>

Browning, KO, DeSarno, MJ, Davis, DR, Streck, JM, Bergeria, C L, Harfmann, RF et al. (2022). Relating individual differences in the reinforcing value of smoking and dependence severity to nicotine exposure levels in vulnerable populations. *Prev Med*, 107312. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36272516>

Taylor, GMJ, & Treur, JL. (2023). An application of the stress-diathesis model: A review about the association between smoking tobacco, smoking cessation, and mental health. *Int J Clin Health Psychol*, 23(1), 100335. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36247407>

van der Heijden HS, Schirmbeck F, Kempton MJ, van der Gaag M, Allott K, et al. Impact of smoking behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study - corrigendum. *European Psychiatry*, 2022; 65(1):e17. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35225210>

Robinson JD, Cui Y, Karam-Hage M, Kypriotakis G, Versace F, et al. Topiramate decreases the salience of motivationally relevant visual cues among smokers with alcohol use disorder. *Alcohol Clin Exp Res*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35037278>

Portilla A, Meza MF, and Lizana PA. Association between emotional exhaustion and tobacco consumption in teachers. *International Journal of Environmental Research and Public Health*, 2022; 19(5). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35270298>

https://mdpi-res.com/d_attachment/ijerph/ijerph-19-02606/article_deploy/ijerph-19-02606.pdf

Ohi K, Takai K, Kuramitsu A, Sugiyama S, and Shioiri T. Common brain cortical abnormality in smoking behavior and bipolar disorder: Discriminant analysis using cortical thickness and surface area. *Cereb Cortex*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35040937>

Lopes de Oliveira T, Oliveira RVC, Griep RH, Moreno AB, Melo ECP, et al. The brazilian longitudinal study of adult health (elsa-brasil): Factors related to smoking cessation. *International Journal of Behavioral Medicine*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34984649>

Lin SC, Gathua N, Thompson C, Sripipatana A, and Makaroff L. Disparities in smoking prevalence and associations with mental health and substance use disorders in underserved communities across the United States. *Cancer*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35253202>

Kim GE, Kim MH, Lim WJ, and Kim SI. The effects of smoking habit change on the risk of depression: Analysis of data from the Korean national health insurance service. *J Affect Disord*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35085672>

Kendall AD, Hedeker D, Diviak KR, and Mermelstein RJ. The mood boost from tobacco cigarettes is more erratic with the additions of cannabis and alcohol. *Nicotine & Tobacco Research*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34999839>

Evans-Polce RJ, Kcomt L, Veliz PT, Boyd CJ, and McCabe SE. Mental health-related quality of life by sex and sexual identity among u.S. Adults with alcohol and tobacco use disorder. *Journal of Psychiatric Research*, 2022; 149:136-44. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35276630>

Dugas EN, Wellman RJ, Sylvestre MP, Belanger RE, and O'Loughlin J. Who mixes tobacco with cannabis and does mixing relate to nicotine dependence? *Addictive Behaviors*, 2022; 128:107254. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35085951>

De Oliveira RM, Santos JLF, and Furegato ARF. Nurses' beliefs, attitudes, and opinions regarding a smoking ban in a psychiatric hospital. *Nursing*, 2022; 52(4):55-61. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35358995>

Australian Institute of Health Welfare. National opioid pharmacotherapy statistics annual data collection. Canberra: AIHW, 2022. Available from: <https://www.aihw.gov.au/reports/alcohol-other-drug-treatment-services/national-opioid-pharmacotherapy-statistics>.

Australian Institute of Health and Welfare. Mental health services in Australia. AIHW, 2022. Available from: <https://www.aihw.gov.au/reports/mental-health-services/mental-health-services-in-australia/report-contents/summary-of-mental-health-services-in-australia>.

Wycoff AM, Motschman CA, Griffin SA, Helle AC, Piasecki TM, et al. Simultaneous use of alcohol and cigarettes in a mixed psychiatric sample: Daily-life associations with smoking motives, craving, stimulation, sedation, and affect. *Psychol Addict Behav*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34726421>

Welfare AloHa. Mental health services in Australia. Canberra: AIHW, 2021. Available from: <https://www.aihw.gov.au/reports/mental-health-services/mental-health-services-in-australia>.

Wang N, Yao T, Sung HY, and Max W. The association of cannabis use and cigarette smoking with psychological distress among adults in california. *Substance Use and Misuse*, 2021:1-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34753379>

van der Heijden S, Schirmbeck F, Kempton MJ, van der Gaag M, Allot K, et al. Impact of smoking behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study. *European Psychiatry*, 2021:1-20. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34544507>

Taylor GM, Lindson N, Farley A, Leinberger-Jabari A, Sawyer K, et al. Smoking cessation for improving mental health. *Cochrane Database of Systematic Reviews*, 2021; 3(3):CD013522. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33687070>

Sher L. Smoking enhances suicide risk-a significant role in the COVID-19 pandemic? *QJM*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34694396>

Shepherd JM, Rogers AH, Garey L, Ditre JW, Smit T, et al. Tobacco use severity in relation to opioid misuse and dependence among adult tobacco users with chronic pain: The moderating role of pain-related anxiety. *Addictive Behaviors*, 2021; 117:106844. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33545622>

Ray LA, Green R, Enders C, Leventhal AM, Grodin EN, et al. Efficacy of combining varenicline and naltrexone for smoking cessation and drinking reduction: A randomized clinical trial. *American Journal of Psychiatry*, 2021; 178(9):818–28. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34080890>

McRobbie H, Boland VC, and Courtney RJ. Assessing cannabis and tobacco co-use: The pros and cons of additional data collection. *Addiction*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33417256>

Hasselblad Lundstrom N, Holgersen NK, and Haastrup MB. The effect of smoking on the plasma concentration of tricyclic antidepressants - a systematic review. *Acta Neuropsychiatr*, 2021;1-24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34497000>

Echeverria I, Cotaina M, Jovani A, Mora R, Haro G, et al. Proposal for the inclusion of tobacco use in suicide risk scales: Results of a meta-analysis. *International Journal of Environmental Research and Public Health*, 2021; 18(11). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34198855>

Driezen P, Gravely S, Wadsworth E, Smith DM, Loewen R, et al. Increasing cannabis use is associated with poorer cigarette smoking cessation outcomes: Findings from the itc four country smoking and vaping surveys, 2016-2018. *Nicotine & Tobacco Research*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34111281>

Domi A, Barbier E, Adermark L, and Domi E. Targeting the opioid receptors: A promising therapeutic avenue for treatment in "heavy drinking smokers". *Alcohol Alcohol*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33479741>

Churchill V, Duong HT, Massey ZB, and Popova L. Serious psychological distress is associated with higher intentions to quit among smokers during the COVID-19 pandemic. *J Psychoactive Drugs*, 2021;1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34930099>

Chesney E, Robson D, Patel R, Shetty H, Richardson S, et al. The impact of cigarette smoking on life expectancy in schizophrenia, schizoaffective disorder and bipolar affective disorder: An electronic case register cohort study. *Schizophr Res*, 2021; 238:29–35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34563995>

Badanapurkar A, Nelson D, Varghese S, Singh R, and Haddad PM. Support and attitudes of qatar mental health professionals to a proposed mental health inpatient smoking ban: Results of a cross-sectional survey. *J Psychiatr Ment Health Nurs*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34143910>

Australian Institute of Health and Welfare. Australian burden of disease study 2018: Key findings. Australian Burden of Disease Study series 24. Cat. no. BOD 30, Canberra: AIHW 2021. Available from: <https://www.aihw.gov.au/reports/burden-of-disease/burden-of-disease-study-2018-key-findings/contents/key-findings>.

Asharani PV, Lau JH, Ai Ling Seet V, Devi F, Wang P, et al. Smoking-related health beliefs in a sample of psychiatric patients: Factors associated with the health beliefs and validation of the health belief questionnaire. *International Journal of Environmental Research and Public Health*, 2021; 18(4). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33562257>

West JC, Peasley-Miklus C, Graham AL, Mays D, Mermelstein R, et al. Impact of alcohol and drug use on smoking and cessation in socioeconomically disadvantaged young adults. *Addictive Behaviors*, 2020; 110:106486. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32688226>

Smith CL, Jenkins G, Burduli E, Tham P, Miguel A, et al. Crossover associations of alcohol and smoking, craving and biochemically verified alcohol and nicotine use in heavy drinking smokers. *Behav Pharmacol*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32516270>

Segura AG, Mitjans M, Fatjo-Vilas M, Garcia-Portilla MP, Garcia-Alvarez L, et al. Smoking cessation improves clinical outcome in severe mental disorders and is modulated by genetic variability at chrna5 gene. *Schizophr Res*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32553632>

Salt V and Osborne C. Mental health, smoking and poverty: Benefits of supporting smokers to quit. *BJPsych Bull*, 2020; 44(5):213–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32847647>

Piercy H, Garfield JBB, Lubman DI, Lam T, and Manning V. Improved rates of treatment success following alcohol and other drug treatment among clients who quit or reduce their tobacco smoking. *Drug and Alcohol Review*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32869419>

Parker MA and Weinberger AH. Opioid use disorder trends from 2002 to 2017 by cigarette smoking status in the United States. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32960266>

Nguyen LC, Durazzo TC, Dwyer CL, Rauch AA, Humphreys K, et al. Predicting relapse after alcohol use disorder treatment in a high-risk cohort: The roles of anhedonia and smoking. *Journal of Psychiatric Research*, 2020; 126:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32403028>

Miller SC, Hemmy Asamsama O, Silvestri MM, Bonanno C, and Krondilou K. Commentary on a study of tobacco-related violence in a mental health setting. *International Journal of Mental Health Nursing*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32510782>

Kwan Y, Kim HS, Kang DR, and Kim ATH. Trend in the prevalence of non-daily smoking and their relationship with mental health using the korea health and nutrition examination survey. *International Journal of Environmental Research and Public Health*, 2020; 17(10). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32414082>

Hayhurst KP, Jones A, Cairns D, Jahr S, Williams E, et al. Tobacco smoking rates in a national cohort of people with substance use disorder receiving treatment. *Eur Addict Res*, 2020:1-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32720918>

Hammett PJ, Taylor BC, Lando HA, Widome R, Erickson DJ, et al. Serious mental illness and smoking cessation treatment utilization: The role of healthcare providers. *J Behav Health Serv Res*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32378032>

Guydish J, Kapiteni K, Le T, Campbell B, Pinsker E, et al. Tobacco use and tobacco services in california substance use treatment programs. *Drug and Alcohol Dependence*, 2020; 214:108173. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32693199>

Gravelly S, Driezen P, Smith DM, Borland R, Lindblom EN, et al. International differences in patterns of cannabis use among adult cigarette smokers: Findings from the 2018 itc four country smoking and vaping survey. *Int J Drug Policy*, 2020; 79:102754. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32305827>

Goodwin RD. Impact of cannabis use on nicotine and tobacco use outcomes. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32480403>

Ferguson TF, Beauchamp A, Rosen EM, Ray AN, Theall KP, et al. Pilot study of the adaptation of an alcohol, tobacco, and illicit drug use intervention for vulnerable urban young adults. *Front Public Health*, 2020; 8:314. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32766200>

de Oliveira RM, Santos JLF, and Furegato ARF. Prevalence and epidemiological profile of smokers in the psychiatric population and general population: Smokers in psychiatric population. *Journal of Addictions Nursing*, 2020; 31(3):E13-E24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32868617>

Das-Munshi J, Semrau M, Barbui C, Chowdhary N, Gronholm PC, et al. Correction to: Gaps and challenges: WHO treatment recommendations for tobacco cessation and management of substance use disorders in people with severe mental illness. *BMC Psychiatry*, 2020; 20(1):326. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32576169>

Das-Munshi J, Semrau M, Barbui C, Chowdhary N, Gronholm PC, et al. Gaps and challenges: WHO treatment recommendations for tobacco cessation and management of substance use disorders in people with severe mental illness. *BMC Psychiatry*, 2020; 20(1):237. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32410598>

Dahal R, Bhattarai A, and Adhikari K. Variation in characteristics of people with mental disorders across smoking status in the canadian general population. *Tob Prev Cessat*, 2020; 6:61. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33163707>

Cruvinel E, Liebman E, Leite I, Hu J, and Richter KP. Prevalence of smoking, quit attempts and access to cessation treatment among adults with mental illness in Brazil: A cross-sectional analysis of a national health survey. *BMJ Open*, 2020; 10(5):e033959. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32461292>

Carpenter KM, Torres AJ, Salmon EE, Carlini BH, Vickerman KA, et al. Marijuana use and adherence to smoking cessation treatment among callers to tobacco quitlines. *Preventing Chronic Disease*, 2020; 17:E102. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32915131>

Caponnetto P, Polosa R, Robson D, and Bauld L. Tobacco smoking, related harm and motivation to quit smoking in people with schizophrenia spectrum disorders. *Health Psychol Res*, 2020; 8(1):9042. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32510003>

Britton M, Haddad S, and Derrick JL. Prospective and daily effects of cannabis use on smoking outcomes during a self-guided quit attempt. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32396637>

Beard E, Brown J, West R, and Michie S. Are population trends in high-risk alcohol consumption in smokers associated with trends in quit attempts and quit success? A time-series analysis. *BMJ Open*, 2020; 10(7):e034262. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32690733>

Australian Institute for Health and Welfare. Mental health services in Australia. AIHW, 2020. Available from: <https://www.aihw.gov.au/reports/mental-health-services/mental-health-services-in-australia/report-contents/summary-of-mental-health-services-in-australia>.

Asharani PV, Ling Seet VA, Abdin E, Siva Kumar FD, Wang P, et al. Smoking and mental illness: Prevalence, patterns and correlates of smoking and smoking cessation among psychiatric patients. *International Journal of Environmental Research and Public Health*, 2020; 17(15). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32752263>

Andersson A, Hegvik TA, Chen Q, Rosenqvist MA, Kvalvik LG, et al. Attention-deficit/hyperactivity disorder and smoking habits in pregnant women. *PLoS One*, 2020; 15(6):e0234561. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32555596>

Weinberger AH, Pacek LR, Wall MM, Gbedemah M, Lee J, et al. Cigarette smoking quit ratios among adults in the USA with cannabis use and cannabis use disorders, 2002-2016. *Tobacco Control*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30952691>

Roche DJO, Bujarski S, Green R, Hartwell EE, Leventhal AM, et al. Alcohol, tobacco, and marijuana consumption is associated with increased odds of same-day substance co- and tri-use. *Drug and Alcohol Dependence*, 2019; 200:40-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31085377>

Oluwoye O, Monroe-DeVita M, Burduli E, Chwastiak L, McPherson S, et al. Impact of tobacco, alcohol and cannabis use on treatment outcomes among patients experiencing first episode psychosis: Data from the national raise-etp study. *Early Interv Psychiatry*, 2019; 13(1):142–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29356438>

Kathuria H, Seibert RG, Cobb V, Herbst N, Weinstein ZM, et al. Perceived barriers to quitting cigarettes among hospitalized smokers with substance use disorders: A mixed methods study. *Addictive Behaviors*, 2019; 95:41-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30836208>

Bold KW, Zweben A, Fucito LM, Piepmeier ME, Muvvala S, et al. Longitudinal findings from a randomized clinical trial of varenicline for alcohol use disorder with comorbid cigarette smoking. *Alcohol Clin Exp Res*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30817018>

Wilhelm K, Handley T, and Reddy P. A case for identifying smoking in presentations to the emergency department with suicidality. *Australas Psychiatry*, 2018; 26(2):176–80. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29417825>

Pacek LR, Copeland J, Dierker L, Cunningham CO, Martins SS, et al. Among whom is cigarette smoking declining in the United States? The impact of cannabis use status, 2002-2015. *Drug and Alcohol Dependence*, 2018; 191:355-60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30179761>

Okoli CTC and Seng S. Associations between secondhand tobacco smoke exposure and nicotine dependence and smoking cessation attempts among adult tobacco users with a psychiatric disorder. *Biol Res Nurs*, 2018:1099800418781914. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29902928>

Nakano H, Ohira T, Maeda M, Yabe H, Ohtsuru A, et al. Associations of disaster-related and psychosocial factors with changes in smoking status after a disaster: A cross-sectional survey after

the great east Japan earthquake. *BMJ Open*, 2018; 8(6):e018943. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29961000>

Monteschi M, de Padua AI, Perez CR, Castellano MV, Cedano S, et al. A simple test to assess the emotional status of smokers. *Clin Respir J*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30256526>

Moeller SJ, Fink DS, Gbedemah M, Hasin DS, Galea S, et al. Trends in illicit drug use among smokers and nonsmokers in the United States, 2002-2014. *Journal of Clinical Psychiatry*, 2018; 79(3).

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29894597>

Lappan S, Thorne CB, Long D, and Hendricks PS. Longitudinal and reciprocal relationships between psychological well-being and smoking. *Nicotine & Tobacco Research*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30219867>

Klemperer EM, Hughes JR, and Naud S. Study characteristics influence the efficacy of substance abuse treatments: A meta-analysis of medications for smoking cessation. *Nicotine & Tobacco Research*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30380134>

Holma KM, Holma I, Ketokivi M, Oquendo MA, and Isometsa E. The relationship between smoking and suicidal behavior in psychiatric patients with major depressive disorder. *Arch Suicide Res*, 2018;1-23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29883280>

Gulsen A and Uygur B. Psychological features of smokers. *Respir Care*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30065079>

Goodwin R, Pacek L, Copeland J, Moeller S, Dierker L, et al. Trends in daily cannabis use among cigarette smokers: United States, 2002-2014. *American Journal of Public Health*, 2018; 108(1):137-42. Available from: <http://ajph.aphapublications.org/doi/abs/10.2105/AJPH.2017.304050>

<https://www.ncbi.nlm.nih.gov/pubmed/29161058>

Ergun D, Ergun R, Ergun B, and Kurt OK. Occupational risk factors and the relationship of smoking with anxiety and depression. *Turk Thorac J*, 2018; 19(2):77-83. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29755811>

Cohn AM, Johnson AL, Rose SW, Pearson JL, Villanti AC, et al. Population-level patterns and mental health and substance use correlates of alcohol, marijuana, and tobacco use and co-use in US young adults and adults: Results from the population assessment for tobacco and health. *American Journal on Addictions*, 2018; 27(6):491-500. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30152111>

Australian Institute of Health and Welfare. Alcohol and other drug treatment services in Australia 2016–17: Key findings. Canberra, AIHW 2018. Available from:

<https://www.aihw.gov.au/reports/alcohol-other-drug-treatment-services/aodts-2016-17-key-findings/contents/summary>.

Australian Bureau of Statistics. National health survey 2017–18: First results. 2018. Available from: <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/national-health-survey-first-results/2017-18>.

Ashare RL and Wetherill RR. The intersection of sex differences, tobacco use, and inflammation: Implications for psychiatric disorders. *Curr Psychiatry Rep*, 2018; 20(9):75. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30094593>

Wang S, Ungvari GS, Forester BP, Chiu HF, Wu Y, et al. Gender differences in general mental health, smoking, drinking and chronic diseases in older adults in jilin province, China. *Psychiatry Res*, 2017; 251:58-62. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28189080>

Waaktaar T, Kan KJ, and Torgersen S. The genetic and environmental architecture of substance use development from early adolescence into young adulthood: A longitudinal twin study of comorbidity of alcohol, tobacco and illicit drugs use. *Addiction*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29057620>

Shu BC, Lung FW, Guo SE, Chen KC, Fang SY, et al. Ineffectiveness of the tobacco hazards prevention act in patients with severe and persistent mental illness. *Public Health*, 2017; 153:118-27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29032168>

Oliveira RM, Santos JLF, and Furegato ARF. Tobacco addiction in the psychiatric population and in the general population. *Rev Lat Am Enfermagem*, 2017; 25:e2945. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29211192>

Okoli CTC, Wiggins A, Fallin-Bennett A, and Rayens MK. A retrospective analysis of the comparative effectiveness of smoking cessation medication among individuals with mental illness in community-based mental health and addictions treatment settings. *J Psychiatr Ment Health Nurs*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28635015>

No authors listed. Mental health and smoking partnership statement on electronic cigarettes why smoking and mental health matters. Smokefree Action UK 2017. Available from: <http://smokefreeaction.org.uk/wp-content/uploads/2017/11/FINAL-Partnership-Statement-on-E-cigarettes.pdf>.

Molla Z, Dube L, Krah W, and Soboka M. Tobacco dependence among people with mental illness: A facility-based cross sectional study from southwest ethiopia. *BMC Res Notes*, 2017; 10(1):289. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28716128>

Martinez U, Fernandez Del Rio E, Lopez-Duran A, Martinez-Vispo C, and Becona E. Types of smokers who seek smoking cessation treatment according to psychopathology. *Journal of Dual Diagnosis*, 2017:0. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29111906>

Loreto AR, Carvalho CFC, Frallonardo FP, Ismael F, Andrade AG, et al. Smoking cessation treatment for patients with mental disorders using cbt and combined pharmacotherapy. *Journal of Dual Diagnosis*, 2017:1-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28489501>

Lo CC, Yang F, Ash-Houchen W, and Cheng TC. Racial/ethnic differences in cigarette use: The roles of mental illness and health-care access/utilization. *Substance Use and Misuse*, 2017;1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29172859>

Kulik MC and Glantz SA. Softening among US smokers with psychological distress: More quit attempts and lower consumption as smoking drops. *American Journal of Preventive Medicine*, 2017; 53(6):810–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29029966>

Kourakos M, Saridi M, Kafkia T, Rekleiti M, Souliotis K, et al. Factors affecting mental patients' behaviors and attitudes regarding smoking. *Asian Pac J Cancer Prev*, 2017; 18(7):1991-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28749641>

King JL, Reboussin BA, Spangler J, Cornacchione Ross J, and Sutfin EL. Tobacco product use and mental health status among young adults. *Addictive Behaviors*, 2017; 77:67-72. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28965069>

Kilibarda B, Mravcik V, Oechsler H, and Martens MS. Association of smoking status with substance use and psychological distress in serbia. *Public Health*, 2017; 152:95-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28869853>

Japuntich SJ, Sherman SE, Joseph AM, Clothier B, Noorbaloochi S, et al. Proactive tobacco treatment for individuals with and without a mental health diagnosis: Secondary analysis of a pragmatic randomized controlled trial. *Addictive Behaviors*, 2017; 76:15-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28735036>

Gupta R, Qaiser B, He L, Hiekkalinna TS, Zheutlin AB, et al. Neuregulin signaling pathway in smoking behavior. *Transl Psychiatry*, 2017; 7(8):e1212. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28892072>

Crowley R, Kirschner N, Dunn A, and Bornstein S. Health and public policy to facilitate effective prevention and treatment of substance use disorders involving illicit and prescription drugs: An american college of physicians position paper. *Annals of Internal Medicine*, 2017; 166(10):733-6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28346947>

Cooney JL, Cooper S, Grant C, Sevarino K, Krishnan-Sarin S, et al. A randomized trial of contingency management for smoking cessation during intensive outpatient alcohol treatment. *Journal of Substance Abuse Treatment*, 2017; 72:89–96. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27542442>

Cohn A, Elmasry H, and Niaura R. Facility-level, state, and financial factors associated with changes in the provision of smoking cessation services in US substance abuse treatment facilities: Results from the national survey of substance abuse treatment services 2006 to 2012. *Journal of Substance Abuse Treatment*, 2017; 77:107-14. Available from:

[http://www.journalofsubstanceabusetreatment.com/article/S0740-5472\(16\)30254-9/fulltext](http://www.journalofsubstanceabusetreatment.com/article/S0740-5472(16)30254-9/fulltext)

<http://www.ncbi.nlm.nih.gov/pubmed/28476262>

Clyde M, Pipe A, Els C, Reid R, and Tulloch H. Factor structure of the smoking cessation self-efficacy questionnaire among smokers with and without a psychiatric diagnosis. *Psychol Addict Behav*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28182446>

Braillon A. Smoking is the first cause of morbidity and death in psychiatric settings. *CMAJ*, 2017; 189(49):E1534. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29229718>

Bradizza CM, Stasiewicz PR, Zhuo Y, Rusczyk M, Maisto SA, et al. Smoking cessation for pregnant smokers: Development and pilot test of an emotion regulation treatment supplement to standard smoking cessation for negative affect smokers. *Nicotine & Tobacco Research*, 2017; 19(5):578-84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28403472>

Baker AL. Commentary on mathew et al. (2017): Improving the effectiveness of interventions to reduce smoking among people living with severe mental ill-health. *Addiction*, 2017; 112(3):413-4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28168789>

Aschbrenner KA, Naslund JA, Gill L, Hughes T, O'Malley AJ, et al. Qualitative analysis of social network influences on quitting smoking among individuals with serious mental illness. *J Ment Health*, 2017;1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28675331>

Aditya A, Kaluskar A, Pandya S, Vinay V, and Sharma A. Risk of psychiatric morbidity in patients with tobacco habits: A cross-sectional study. *J Clin Diagn Res*, 2017; 11(7):ZC33-ZC5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28893039>

Smoking cessation services can also help improve mental health. *Nursing Standard*, 2017; 31(24):16. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28176559>

"Psychiatric disorders in smokers seeking treatment for tobacco dependence: Relations with tobacco dependence and cessation": Correction to piper et al. (2010). *Journal of Consulting and Clinical Psychology*, 2017; 85(9):853. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28857593>

Wang JB and Cataldo JK. Medical marijuana legalization and co-use in adult cigarette smokers. *American Journal of Health Behavior*, 2016; 40(2):205-14. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26931752>

US Food and Drug Administration (FDA). Joint meeting of the psychopharmacologic drugs advisory committee and drug safety and risk management advisory committee. 2016. Available from: <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/PsychopharmacologicDrugsAdvisoryCommittee/UCM520103.pdf>

Tatreau JR, Harris S, Sheitman B, and Steiner BD. Cardiometabolic assessment, diagnosis, and treatment of chronic medical illnesses during an inpatient psychiatric hospitalization: Colocated medical care versus treatment as usual. *Prim Care Companion CNS Disord*, 2016; 18(6). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28006091>

Tam J, Warner KE, and Meza R. Smoking and the reduced life expectancy of individuals with serious mental illness. *American Journal of Preventive Medicine*, 2016; 51(6):958–66. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27522471>

- Subramaniam P, McGlade E, and Yurgelun-Todd D. Comorbid cannabis and tobacco use in adolescents and adults. *Curr Addict Rep*, 2016; 3(2):182-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27175326>
- Slyepchenko A, Brunoni AR, McIntyre RS, Quevedo J, and Carvalho AF. The adverse effects of smoking on health outcomes in bipolar disorder: A review and synthesis of biological mechanisms. *Curr Mol Med*, 2016; 16(2):187–205. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26812916>
- Schauer GL, Hall CD, Berg CJ, Donovan DM, Windle M, et al. Differences in the relationship of marijuana and tobacco by frequency of use: A qualitative study with adults aged 18-34 years. *Psychol Addict Behav*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27099958>
- Sankaranarayanan A and Castle D. Burden associated with smoking as a suicidal risk factor in an Australian sample of patients with psychosis. *Australas Psychiatry*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27130725>
- Ruggles KV, Fang Y, Tate J, Mentor SM, Bryant KJ, et al. What are the patterns between depression, smoking, unhealthy alcohol use, and other substance use among individuals receiving medical care? A longitudinal study of 5479 participants. *AIDS Behav*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27475945>
- Poorolajal J and Darvishi N. Smoking and suicide: A meta-analysis. *PLoS One*, 2016; 11(7):e0156348. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27391330>
- Mould DR and Meibohm B. Drug development of therapeutic monoclonal antibodies. *BioDrugs*, 2016; 30(4):275–93. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27342605>
- Meurk C, Ford P, Sharma R, Fitzgerald L, and Gartner C. Views and preferences for nicotine products as an alternative to smoking: A focus group study of people living with mental disorders. *International Journal of Environmental Research and Public Health*, 2016; 13(11). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27886046>
- Lo J, Patel P, and Roberts B. A systematic review on tobacco use among civilian populations affected by armed conflict. *Tobacco Control*, 2016; 25(2):129–40. Available from: <http://tobaccocontrol.bmj.com/content/25/2/129.abstract>
- Keskitalo-Vuokko K, Korhonen T, and Kaprio J. Gene-environment interactions between depressive symptoms and smoking quantity. *Twin Res Hum Genet*, 2016:1–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27161145>
- Garcia-Portilla MP and Bobes J. Smoking cessation programs for persons with schizophrenia: An urgent unmet need. *Rev Psiquiatr Salud Ment*, 2016; 9(4):181-4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27969002>
- Foster DW, Buckner JD, Schmidt NB, and Zvolensky MJ. Multisubstance use among treatment-seeking smokers: Synergistic effects of coping motives for cannabis and alcohol use and social anxiety/depressive symptoms. *Substance Use and Misuse*, 2016; 51(2):165–78. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26846421>

Fischer B, Ialomiteanu AR, Aebly S, Rudzinski K, Kurdyak P, et al. Substance use, health, and functioning characteristics of medical marijuana program participants compared to the general adult population in ontario (canada). *J Psychoactive Drugs*, 2016:1-8. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28001910>

Dickerson F, Adamos MB, Katsafanas E, Khushalani S, Origoni A, et al. The association among smoking, hsv-1 exposure, and cognitive functioning in schizophrenia, bipolar disorder, and non-psychiatric controls. *Schizophr Res*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27262384>

Cooper J, Borland R, McKee SA, Yong HH, and Dugue PA. Depression motivates quit attempts but predicts relapse: Differential findings for gender from the international Tobacco Control study.

Addiction, 2016; 111(8):1438–47. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26888199>

Chen LS, Baker T, Brownson RC, Carney RM, Jorenby D, et al. Smoking cessation and electronic cigarettes in community mental health centers: Patient and provider perspectives. *Community Ment Health J*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27900650>

Burki TK. Smoking and mental health. *The Lancet Respiratory Medicine*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27180017>

Berlin I. Comorbidities and tobacco dependence treatment initiation. *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26911841>

Bazo-Alvarez JC, Peralta-Alvarez F, Bernabe-Ortiz A, Alvarado GF, and Miranda JJ. Tobacco consumption and positive mental health: An epidemiological study from a positive psychology perspective. *BMC Psychology*, 2016; 4:22. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27142187>

Anthenelli RM, Benowitz NL, West R, St Aubin L, McRae T, et al. Neuropsychiatric safety and efficacy of varenicline, bupropion, and nicotine patch in smokers with and without psychiatric disorders (eagles): A double-blind, randomised, placebo-controlled clinical trial. *The Lancet*, 2016; 387(10037):2507–20. Available from: [https://doi.org/10.1016/S0140-6736\(16\)30272-0](https://doi.org/10.1016/S0140-6736(16)30272-0)

Almadana Pacheco V, Gomez-Bastero Fernandez AP, Valido Morales A, Luque Crespo E, Monserrat S, et al. Anxiety, depression and tobacco abstinence. *Adicciones*, 2016:761. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27749973>

Welch AE, Jasek JP, Caramanica K, Chiles MC, and Johns M. Cigarette smoking and 9/11-related posttraumatic stress disorder among world trade center health registry enrollees, 2003-12. *Preventive Medicine*, 2015; 73C:94–9. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25655710>

Weidberg S, Garcia-Rodriguez O, Yoon JH, and Secades-Villa R. Interaction of depressive symptoms and smoking abstinence on delay discounting rates. *Psychol Addict Behav*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25844830>

Walsh R, Schweinfurth L, and Dickerson F. Smoking and smoking cessation treatment among hospitalized psychiatric patients. *Psychiatric Services*, 2015; 66(4):442–3. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25828985>

Trujillo MA, Khoddam R, Greenberg JB, Dyal SR, Ameringer KJ, et al. Distress tolerance as a correlate of tobacco dependence and motivation: Incremental relations over and above anxiety and depressive symptoms. *Behav Med*, 2015:0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26651507>

Torres OV and O'Dell LE. Stress is a principal factor that promotes tobacco use in females. *Prog Neuropsychopharmacol Biol Psychiatry*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25912856>

Thomson D, Berk M, Dodd S, Rapado-Castro M, Quirk SE, et al. Tobacco use in bipolar disorder. *Clinical Psychopharmacology and Neuroscience*, 2015; 13(1):1–11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25912533>

Taylor G, Taylor A, Munafo MR, McNeill A, and Aveyard P. Does smoking reduction worsen mental health? A comparison of two observational approaches. *BMJ Open*, 2015; 5(5):e007812. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25979871>

Szatkowski L and McNeill A. Diverging trends in smoking behaviors according to mental health status. *Nicotine & Tobacco Research*, 2015; 17(3):356-60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25180078>

Short NA, Oglesby ME, Raines AM, Zvolensky MJ, and Schmidt NB. Posttraumatic stress and emotion dysregulation: Relationships with smoking to reduce negative affect and barriers to smoking cessation. *Compr Psychiatry*, 2015; 61:15–22. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26070492>

Schauer GL, Berg CJ, Kegler MC, Donovan DM, and Windle M. Assessing the overlap between tobacco and marijuana: Trends in patterns of co-use of tobacco and marijuana in adults from 2003-2012. *Addictive Behaviors*, 2015; 49:26-32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26036666>

Schauer GL, Berg CJ, Kegler MC, Donovan DM, and Windle M. Differences in tobacco product use among past month adult marijuana users and nonusers: Findings from the 2003-2012 national survey on drug use and health. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26009578>

Oliveira RM, Siqueira Junior AC, and Furegato AR. The meaning of smoking for patients with mental disorder. *Issues Ment Health Nurs*, 2015; 36(2):127-34. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25625713>

Mitchell AJ, Vancampfort D, De Hert M, and Stubbs B. Do people with mental illness receive adequate smoking cessation advice? A systematic review and meta-analysis. *General Hospital Psychiatry*, 2015; 37(1):14–23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25510845>

Manzella F, Maloney SE, and Taylor GT. Smoking in schizophrenic patients: A critique of the self-medication hypothesis. *World J Psychiatry*, 2015; 5(1):35–46. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25815253>

Luo SX, Covey LS, Hu MC, Levin FR, Nunes EV, et al. Toward personalized smoking-cessation treatment: Using a predictive modeling approach to guide decisions regarding stimulant medication treatment of attention-deficit/hyperactivity disorder (adhd) in smokers. *American Journal on Addictions*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25659348>

Khati I, Menvielle G, Chollet A, Younes N, Metadieu B, et al. What distinguishes successful from unsuccessful tobacco smoking cessation? Data from a study of young adults (tempo). *Preventive Medicine Reports*, 2015; 2:679–85. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26844137>

Jardin C, Bakshaie J, Schmidt NB, Sharp C, and Zvolensky M. Examination of smoking inflexibility as a mechanism linking anxiety sensitivity and severity of smoking behavior. *American Journal on Addictions*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25808783>

Janes AC, Pedrelli P, Whitton AE, Pechtel P, Douglas S, et al. Reward responsiveness varies by smoking status in women with a history of major depressive disorder. *Neuropsychopharmacology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25662839>

Gurillo P, Jauhar S, Murray RM, and MacCabe JH. Does tobacco use cause psychosis? Systematic review and meta-analysis. *The Lancet Psychiatry*, 2015; 2(8):718–25. Available from: [http://dx.doi.org/10.1016/S2215-0366\(15\)00152-2](http://dx.doi.org/10.1016/S2215-0366(15)00152-2)

Foster DW, Langdon KJ, Schmidt NB, and Zvolensky M. Smoking processes, panic, and depressive symptoms among treatment-seeking smokers. *Substance Use and Misuse*, 2015; 50(3):394–402. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25496170>

Chapman S. A complete smoking ban in psychiatric hospitals is ethically wrong. *British Medical Journal*, 2015; 351:h6288. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26607749>

Buckner JD, Farris SG, Zvolensky MJ, Shah SM, Leventhal AM, et al. Dysphoria and smoking among treatment seeking smokers: The role of smoking-related inflexibility/avoidance. *American Journal of Drug and Alcohol Abuse*, 2015; 41(1):45–51. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25396705>

Brandt CP, Bakhshaie J, Garey L, Schmidt NB, Leventhal AM, et al. The moderating role of smoking amount per day on the relations between anxiety sensitivity, smoking dependence, and cognitive-affective aspects of smoking among treatment seeking smokers. *Addictive Behaviors Reports*, 2015; 1:26–33. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26114159>

Asnaani A, Alpert E, McLean CP, and Foa EB. Resilient but addicted: The impact of resilience on the relationship between smoking withdrawal and ptsd. *Journal of Psychiatric Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25881517>

Arnott D, Wessely S, and Fitzpatrick M. Should psychiatric hospitals completely ban smoking? *British Medical Journal*, 2015; 351:h5654. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26536887>

Zvolensky MJ, Farris SG, Leventhal AM, and Schmidt NB. Anxiety sensitivity mediates relations between emotional disorders and smoking. *Psychol Addict Behav*, 2014; 28(3):912–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25222178>

Zvolensky MJ, Farris SG, Leventhal AM, Ditre JW, and Schmidt NB. Emotional disorders and smoking: Relations to quit attempts and cessation strategies among treatment-seeking smokers. *Addictive Behaviors*, 2014; 40C:126–31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25260199>

Zvolensky MJ, Bakhshaie J, Sheffer C, Perez A, and Goodwin RD. Major depressive disorder and smoking relapse among adults in the United States: A 10-year, prospective investigation. *Psychiatry Res*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25650047>

Zorick T, Mandelkern MA, and Brody AL. A naturalistic study of the association between antidepressant treatment and outcome of smoking cessation treatment. *Journal of Clinical Psychiatry*, 2014; 75(12):e1433–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25551240>

Young-Wolff KC, Fromont SC, Delucchi K, Hall SE, Hall SM, et al. Ptsd symptomatology and readiness to quit smoking among women with serious mental illness. *Addictive Behaviors*, 2014; 39(8):1231–4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24813548>

Wilson SM, Dedert EA, Dennis PA, Dennis MF, Calhoun PS, et al. Do ethnicity and gender moderate the influence of posttraumatic stress disorder on time to smoking lapse? *Addictive Behaviors*, 2014; 39(7):1163–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24727113>

Williams JM, Stroup TS, Brunette MF, and Raney LE. Tobacco use and mental illness: A wake-up call for psychiatrists. *Psychiatric Services*, 2014; 65(12):1406–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25270381>

Twyman L, Bonevski B, Paul C, and Bryant J. Perceived barriers to smoking cessation in selected vulnerable groups: A systematic review of the qualitative and quantitative literature. *BMJ Open*, 2014; 4(12):e006414. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25534212>

The Royal Australian College of General Practitioners, Supporting smoking cessation: A guide for health professionals. Melbourne: RACGP; 2014. Available from: <http://www.racgp.org.au/your-practice/guidelines/smoking-cessation/>.

Taylor AE, Fluharty ME, Bjorngaard JH, Gabrielsen ME, Skorpen F, et al. Investigating the possible causal association of smoking with depression and anxiety using mendelian randomisation meta-analysis: The carta consortium. *BMJ Open*, 2014; 4(10):e006141. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25293386>

Stewart DW, Reitzel LR, Correa-Fernandez V, Cano MA, Adams CE, et al. Social support mediates the association of health literacy and depression among racially/ethnically diverse smokers with low socioeconomic status. *Journal of Behavioral Medicine*, 2014; 37(6):1169–79. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25391450>

Smith PH, Mazure CM, and McKee SA. Smoking and mental illness in the US population. *Tobacco Control*, 2014; 23(e2):e147–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24727731>

Smith PH, Homish GG, Giovino GA, and Kozlowski LT. Cigarette smoking and mental illness: A study of nicotine withdrawal. *American Journal of Public Health*, 2014; 104(2):e127-33. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24328637>

Shi Y. Smoking cessation among people seeking mental health treatment. *Psychiatric Services*, 2014; 65(7):957–60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26037007>

Schluger NW. The electronic cigarette: A knight in shining armour or a trojan horse? *Psychiatr Bull* (2014), 2014; 38(5):201–3. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25285216>

Sanderson SC, Taylor A, and Munafo M. Article does not prove that smoking cessation has an "effect" on mental health. *British Medical Journal*, 2014; 348:g2018. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24615004>

Ratschen E. Electronic cigarettes in mental health settings - solving a conundrum? *Psychiatr Bull* (2014), 2014; 38(5):226–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25285221>

Rae J, Pettey D, Aubry T, and Stol J. Factors affecting smoking cessation efforts of people with severe mental illness: A qualitative study. *Journal of Dual Diagnosis*, 2014:0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25491704>

Pang RD, Khoddam R, Guillot CR, and Leventhal AM. Depression and anxiety symptoms moderate the relation between negative reinforcement smoking outcome expectancies and nicotine dependence. *J Stud Alcohol Drugs*, 2014; 75(5):775–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25208195>

Okoli CT and Khara M. Smoking cessation outcomes and predictors among individuals with co-occurring substance use and/or psychiatric disorders. *Journal of Dual Diagnosis*, 2014; 10(1):9–18. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25392058>

Morris CD, Burns EK, Waxmonsky JA, and Levinson AH. Smoking cessation behaviors among persons with psychiatric diagnoses: Results from a population-level state survey. *Drug and Alcohol Dependence*, 2014; 136:63–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24417963>

Minami H, Kahler CW, Bloom EL, Prince MA, Abrantes AM, et al. Effects of sequential fluoxetine and gender on prequit depressive symptoms, affect, craving, and quit day abstinence in smokers with elevated depressive symptoms: A growth curve modeling approach. *Experimental and Clinical Psychopharmacology*, 2014; 22(5):392–406. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25089930>

McClure EA, Campbell AN, Pavlicova M, Hu M, Winhusen T, et al. Cigarette smoking during substance use disorder treatment: Secondary outcomes from a national drug abuse treatment clinical trials network study. *Journal of Substance Abuse Treatment*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25595301>

Malte CA, Dennis PA, Saxon AJ, McFall M, Carmody TP, et al. Tobacco use trajectories among a large cohort of treated smokers with posttraumatic stress disorder. *Addictive Behaviors*, 2014; 41C:238–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25452071>

Leventhal AM and Zvolensky MJ. Anxiety, depression, and cigarette smoking: A transdiagnostic vulnerability framework to understanding emotion-smoking comorbidity. *Psychol Bull*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25365764>

Leventhal AM, Trujillo M, Ameringer KJ, Tidey JW, Sussman S, et al. Anhedonia and the relative reward value of drug and nondrug reinforcers in cigarette smokers. *J Abnorm Psychol*, 2014; 123(2):375–86. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24886011>

Leventhal AM, Piper ME, Japuntich SJ, Baker TB, and Cook JW. Anhedonia, depressed mood, and smoking cessation outcome. *Journal of Consulting and Clinical Psychology*, 2014; 82(1):122–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24219183>

Langdon KJ and Leventhal AM. Posttraumatic stress symptoms and tobacco abstinence effects in a non-clinical sample: Evaluating the mediating role of negative affect reduction smoking expectancies. *J Psychopharmacol*, 2014; 28(11):1009–17. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25142407>

Kollins SH and Adcock RA. Adhd, altered dopamine neurotransmission, and disrupted reinforcement processes: Implications for smoking and nicotine dependence. *Prog Neuropsychopharmacol Biol Psychiatry*, 2014; 52:70–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24560930>

Kimbrel NA, Morissette SB, Gulliver SB, Langdon KJ, and Zvolensky MJ. The effect of social anxiety on urge and craving among smokers with and without anxiety disorders. *Drug and Alcohol Dependence*, 2014; 135:59–64. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24331637>

Kelly MM, Sido H, Forsyth JP, Ziedonis DM, Kalman D, et al. Acceptance and commitment therapy smoking cessation treatment for veterans with posttraumatic stress disorder: A pilot study. *Journal of Dual Diagnosis*, 2014;0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25491589>

Hruska B, Bernier J, Kenner F, Kenne DR, Boros AP, et al. Examining the relationships between posttraumatic stress disorder symptoms, positive smoking outcome expectancies, and cigarette smoking in people with substance use disorders: A multiple mediator model. *Addictive Behaviors*, 2014; 39(1):273–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24144587>

Hooker SA, MacGregor KL, Funderburk JS, and Maisto SA. Body mass index and depressive symptoms in primary care settings: Examining the moderating roles of smoking status, alcohol consumption and vigorous exercise. *Clin Obes*, 2014; 4(1):21–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25425129>

Haller CS, Etter JF, and Courvoisier DS. Trajectories in cigarette dependence as a function of anxiety: A multilevel analysis. *Drug and Alcohol Dependence*, 2014; 139:115–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24703608>

Guillot CR, Pang RD, and Leventhal AM. Anxiety sensitivity and negative urgency: A pathway to negative reinforcement-related smoking expectancies. *Journal of Addiction Medicine*, 2014; 8(3):189–94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24662369>

Glasheen C, Hedden SL, Forman-Hoffman VL, and Colpe LJ. Cigarette smoking behaviors among adults with serious mental illness in a nationally representative sample. *Annals of Epidemiology*, 2014; 24(10):776–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25169683>

Garcia-Portilla MP, Garcia-Alvarez L, Saiz PA, Diaz-Mesa E, Galvan G, et al. Effectiveness of a multi-component smoking cessation support programme (mcscsp) for patients with severe mental disorders: Study design. *International Journal of Environmental Research and Public Health*, 2014; 11(1):373–89. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24368428>

Furukawa TA. Helping people with schizophrenia to quit smoking. *Evid Based Ment Health*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25527712>

Fucito LM, Redeker NS, Ball SA, Toll BA, Ikomi JT, et al. Integrating a behavioural sleep intervention into smoking cessation treatment for smokers with insomnia: A randomised pilot study. *J Smok Cessat*, 2014; 9(1):31–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24995044>

Fraser D, Gartner C, and Hall W. Patterns of tobacco smoking among illicit drug users in Australia 2001–2010. *Drug and Alcohol Review*, 2014; 33(5):534–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25069871>

Filia SL, Baker AL, Gurvich CT, Richmond R, Lewin TJ, et al. Gender differences in characteristics and outcomes of smokers diagnosed with psychosis participating in a smoking cessation intervention. *Psychiatry Res*, 2014; 215(3):586–93. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24485064>

Filia SL, Baker AL, Gurvich CT, Richmond R, and Kulkarni J. The perceived risks and benefits of quitting in smokers diagnosed with severe mental illness participating in a smoking cessation intervention: Gender differences and comparison to smokers without mental illness. *Drug and Alcohol Review*, 2014; 33(1):78–85. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24256336>

Fergusson DM and Boden JM. Commentary on prince van leeuwen et al. (2014): Tobacco and cannabis use. *Addiction*, 2014; 109(2):312–3. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24422619>

Farris SG, Zvolensky MJ, Norton PJ, Hogan J, Smith AH, et al. Smoking-specific experiential avoidance is indirectly associated with trait worry and smoking processes among treatment-seeking smokers. *Behav Med*, 2014:0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25398072>

Ezquiaga E, Garcia-Lopez A, de Dios C, Agud JL, Albillo D, et al. Seasonality, smoking and history of poor treatment compliance are strong predictors of dropout in a naturalistic 6 year follow-up of bipolar patients. *Psychiatr Q*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24986371>

Evins AE, Cather C, Pratt SA, Pachas GN, Hoepfner SS, et al. Maintenance treatment with varenicline for smoking cessation in patients with schizophrenia and bipolar disorder: A randomized clinical trial. *Journal of the American Medical Association*, 2014; 311(2):145–54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24399553>

Doyle F, Rohde D, Rutkowska A, Morgan K, Cousins G, et al. Systematic review and meta-analysis of the impact of depression on subsequent smoking cessation in patients with coronary heart disease: 1990 to 2013. *Psychosom Med*, 2014; 76(1):44–57. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24367125>

Dedert EA, Dennis PA, Swinkels CM, Calhoun PS, Dennis MF, et al. Ecological momentary assessment of posttraumatic stress disorder symptoms during a smoking quit attempt. *Nicotine & Tobacco Research*, 2014; 16(4):430–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24191981>

Cosgrove KP, McKay R, Esterlis I, Kloczynski T, Perkins E, et al. Tobacco smoking interferes with gabaa receptor neuroadaptations during prolonged alcohol withdrawal. *Proc Natl Acad Sci U S A*, 2014; 111(50):18031–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25453062>

Cook BL, Wayne GF, Kafali EN, Liu Z, Shu C, et al. Trends in smoking among adults with mental illness and association between mental health treatment and smoking cessation. *Journal of the American Medical Association*, 2014; 311(2):172–82. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24399556>

Cerimele JM, Halperin AC, and Saxon AJ. Tobacco use treatment in primary care patients with psychiatric illness. *Journal of the American Board of Family Medicine*, 2014; 27(3):399–410. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24808119>

Cavazos-Rehg PA, Breslau N, Hatsukami D, Krauss MJ, Spitznagel EL, et al. Smoking cessation is associated with lower rates of mood/anxiety and alcohol use disorders. *Psychological Medicine*, 2014; 44(12):2523–35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25055171>

Capron DW, Allan NP, Norr AM, Zvolensky MJ, and Schmidt NB. The effect of successful and unsuccessful smoking cessation on short-term anxiety, depression, and suicidality. *Addictive Behaviors*, 2014; 39(4):782–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24457901>

Callaghan RC, Veldhuizen S, Jeysingh T, Orlan C, Graham C, et al. Patterns of tobacco-related mortality among individuals diagnosed with schizophrenia, bipolar disorder, or depression. *Journal of Psychiatric Research*, 2014; 48(1):102–10. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24139811>

Burki TK. Smoking and mental illness-time to break the link. *The Lancet Respiratory Medicine*, 2014; 2(3):177. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24745048>

Brodbeck J, Bachmann MS, Brown A, and Znoj HJ. Effects of depressive symptoms on antecedents of lapses during a smoking cessation attempt: An ecological momentary assessment study. *Addiction*, 2014; 109(8):1363–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24690068>

Bonevski B, Regan T, Paul C, Baker AL, and Bisquera A. Associations between alcohol, smoking, socioeconomic status and comorbidities: Evidence from the 45 and up study. *Drug and Alcohol Review*, 2014; 33(2):169–76. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24372899>

Boggs DL, Carlson J, Cortes-Briones J, Krystal JH, and D'Souza DC. Going up in smoke? A review of nAchRs-based treatment strategies for improving cognition in schizophrenia. *Curr Pharm Des*, 2014; 20(31):5077-92. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24345265>

Bloom EL, Oliver JA, Sutton SK, Brandon TH, Jacobsen PB, et al. Post-operative smoking status in lung and head and neck cancer patients: Association with depressive symptomatology, pain, and fatigue. *Psychooncology*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25257853>

Bland P. Smoking cessation improves anxiety depression. *Practitioner*, 2014; 258(1769):5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24791405>

Barros FC, Melo AP, Cournos F, Cherchiglia ML, Peixoto ER, et al. Cigarette smoking among psychiatric patients in Brazil. *Cad Saude Publica*, 2014; 30(6):1195-206. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25099043>

Balbuena L and Tempier R. Independent association of chronic smoking and abstinence with suicide. *Psychiatric Services*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25269443>

Ashare R, Strasser AA, Wileyto EP, Cuevas J, and Audrain-McGovern J. Cognitive deficits specific to depression-prone smokers during abstinence. *Experimental and Clinical Psychopharmacology*, 2014; 22(4):323–31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24932895>

Aschbrenner KA, Ferron JC, Mueser KT, Bartels SJ, and Brunette MF. Social predictors of cessation treatment use among smokers with serious mental illness. *Addictive Behaviors*, 2014; 41C:169–74. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25452062>

Abramovitch A, Pizzagalli DA, Geller DA, Reuman L, and Wilhelm S. Cigarette smoking in obsessive-compulsive disorder and unaffected parents of ocd patients. *European Psychiatry*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24637253>

Bulletin comment: Smoking, mental ill health and electronic cigarettes. *Psychiatr Bull* (2014), 2014; 38(5):249. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25285225>

Weinberger A, Pilver C, Hoff R, Mazure C, and McKee S. Changes in smoking for adults with and without alcohol and drug use disorders: Longitudinal evaluation in the US population. *The American Journal of Drug and Alcohol Abuse*, 2013; 39(3):186–93. Available from: <http://informahealthcare.com/doi/full/10.3109/00952990.2013.785557>
<http://www.ncbi.nlm.nih.gov/pubmed/23721534>

van der Meer RM, Willemsen MC, Smit F, and Cuijpers P. Smoking cessation interventions for smokers with current or past depression. *Cochrane Database of Systematic Reviews*, 2013; 8(8):CD006102. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23963776>

Tsoi DT, Porwal M, and Webster AC. Interventions for smoking cessation and reduction in individuals with schizophrenia. *Cochrane Database of Systematic Reviews*, 2013; 2(2):CD007253. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23450574>

Ragg M, Gordon R, Ahmed T, and Allan J. The impact of smoking cessation on schizophrenia and major depression. *Australas Psychiatry*, 2013; 21(3):238–45. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23616382>

Minichino A, Bersani FS, Calo WK, Spagnoli F, Francesconi M, et al. Smoking behaviour and mental health disorders--mutual influences and implications for therapy. *International Journal of Environmental Research and Public Health*, 2013; 10(10):4790–811. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24157506>

Leyro TM, Hall SM, Hickman N, Kim R, Hall SE, et al. Clinical management of tobacco dependence in inpatient psychiatry: Provider practices and patient utilization. *Psychiatric Services*, 2013; 64(11):1161–5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24185538>

Hermens DF, Scott EM, White D, Lynch M, Lagopoulos J, et al. Frequent alcohol, nicotine or cannabis use is common in young persons presenting for mental healthcare: A cross-sectional study. *BMJ Open*, 2013; 3(2):e002229. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23381649>

Hahn L, Rigby A, and Galletly C. Determinants of high rates of smoking among people with psychosis living in a socially disadvantaged region in south Australia. *The Australian and New Zealand Journal of Psychiatry*, 2013; [Epub ahead of print]. Available from: <http://anp.sagepub.com/content/early/2013/06/04/0004867413491158.long>

<http://www.ncbi.nlm.nih.gov/pubmed/23739313>

Gehricke JG, Polzonetti C, Caburian C, and Gratton E. Prefrontal hemodynamic changes during cigarette smoking in young adult smokers with and without adhd. *Pharmacol Biochem Behav*, 2013; 112:78–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24125785>

Deruiter W, Cheng C, Gehrs M, Langley J, and Dewa C. Substance abuse and smoking among a canadian cohort of first episode psychosis patients. *Community Mental Health Journal*, 2013; [Epub ahead of print]. Available from: <http://link.springer.com/article/10.1007%2Fs10597-013-9634-2>

<http://www.ncbi.nlm.nih.gov/pubmed/23775243>

Department of Health, National mental health report. Australian Government; 2013.

Chen WJ, Ting TT, Chang CM, Liu YC, and Chen CY. Ketamine use among regular tobacco and alcohol users as revealed by respondent driven sampling in taipei: Prevalence, expectancy, and users' risky decision making. *Journal of Food and Drug Analysis*, 2013; 21(4):S102-S5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25264412>

Castaldelli-Maia JM, Carvalho CF, Armentano F, Frallonardo FP, Alves TC, et al. Outcome predictors of smoking cessation treatment provided by an addiction care unit between 2007 and 2010. *Rev Bras Psiquiatr*, 2013; 35(4):338–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24402207>

Berg CJ, Wen H, Cummings JR, Ahluwalia JS, and Druss BG. Depression and substance abuse and dependency in relation to current smoking status and frequency of smoking among nondaily and daily smokers. *American Journal on Addictions*, 2013; 22(6):581-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24131166>

Anthenelli RM, Morris C, Ramey TS, Dubrava SJ, Tsilkos K, et al. Effects of varenicline on smoking cessation in adults with stably treated current or past major depression: A randomized trial. *Annals*

of Internal Medicine, 2013; 159(6):390–400. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24042367>

Amundson EP. Extenuating circumstances--tobacco use and mental illness: Treating addiction. South Dakota Medicine, 2013; 66(8):331. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24175501>

Wu BS, Weinberger AH, Mancuso E, Wing VC, Haji-Khamneh B, et al. A preliminary feasibility study of varenicline for smoking cessation in bipolar disorder. Journal of Dual Diagnosis, 2012; 8(2):131–2. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22962546>

Wilson N, Weerasekera D, Kahler C, Borland R, and Edwards R. Hazardous patterns of alcohol use are relatively common in smokers: Itc project (New Zealand). The New Zealand Medical Journal, 2012; 125(1348):34–41. Available from: <http://journal.nzma.org.nz/journal/abstract.php?id=5022>

Willcutt EG. The prevalence of dsm-iv attention-deficit/hyperactivity disorder: A meta-analytic review. Neurotherapeutics, 2012; 9(3):490–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22976615>

Weinberger AH, Pilver CE, Desai RA, Mazure CM, and McKee SA. The relationship of major depressive disorder and gender to changes in smoking for current and former smokers: Longitudinal evaluation in the US population. Addiction, 2012; 107(10):1847–56. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22429388>

Punnoose S and Belgamwar MR. Nicotine for schizophrenia. The Cochrane Library, 2012; (Issue 1, Art. No.: CD004838). Available from: http://www.cochrane.org/CD004838/SCHIZ_nicotine-for-schizophrenia

Kelly DL, Raley HG, Lo S, Wright K, Liu F, et al. Perception of smoking risks and motivation to quit among nontreatment-seeking smokers with and without schizophrenia. Schizophrenia Bulletin, 2012; 38(3):543–51. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21041835>

Harrell P, Trenz R, Scherer M, Pacek L, and Latimer W. Cigarette smoking, illicit drug use, and routes of administration among heroin and cocaine users. Addictive Behaviors, 2012; [Epub ahead of print]. Available from: <http://www.sciencedirect.com/science/article/pii/S0306460312000123>

George TP, Wu BS, and Weinberger AH. A review of smoking cessation in bipolar disorder: Implications for future research. Journal of Dual Diagnosis, 2012; 8(2):126–30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22737046>

Cooper J, Mancuso SG, Borland R, Slade T, Galletly C, et al. Tobacco smoking among people living with a psychotic illness: The second Australian survey of psychosis. Australian and New Zealand Journal of Psychiatry, 2012; 46(9):851–63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22645396>

Australian Bureau of Statistics. 4364.0.55.001 - Australian health survey: First results, 2011–12. 2012. Available from: <http://www.abs.gov.au/ausstats/abs@.nsf/lookup/4364.0.55.001main+features12011-12>

Aubin HJ, Rollema H, Svensson TH, and Winterer G. Smoking, quitting, and psychiatric disease: A review. *Neurosci Biobehav Rev*, 2012; 36(1):271–84. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21723317>

Yaworski D, Robinson J, Sareen J, and Bolton JM. The relation between nicotine dependence and suicide attempts in the general population. *Canadian Journal of Psychiatry*, 2011; 56(3):161–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21443823>

Williams JM, Gandhi KK, Lu SE, Kumar S, Steinberg ML, et al. Shorter interpuff interval is associated with higher nicotine intake in smokers with schizophrenia. *Drug and Alcohol Dependence*, 2011; 118(2-3):313–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21596491>

Weiner E, Buchholz A, Coffay A, Liu F, McMahon R, et al. Varenicline for smoking cessation in people with schizophrenia: A double blind randomized pilot study. *Schizophrenia Research*, 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21376537>

Vinci C, McVay M, Copeland A, and Carrigan M. The relationship between depression level and smoking motives in college smokers. *Psychology of Addictive Behaviors*, 2011; [Epub ahead of print]. Available from: <http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2011-09194-001>

Vansickel A, Stoops W, Glaser P, Poole M, and Rush C. Methylphenidate increases cigarette smoking in participants with adhd. *Psychopharmacology* 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21590284>

Tregellas J, Tanabe J, Rojas D, Shatti S, Olincy A, et al. Effects of an alpha 7-nicotinic agonist on default network activity in schizophrenia. *Biological Psychiatry*, 2011; 69(1):7–11. Available from: <http://www.biologicalpsychiatryjournal.com/article/PIIS0006322310007079/fulltext>

Torchalla I, Okoli C, Malchy L, and Johnson J. Nicotine dependence and gender differences in smokers accessing community mental health services. *Journal of Psychiatric and Mental Health Nursing*, 2011; 18(4):349–58. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2850.2010.01674.x/full>

Tidey J, Rohsenow D, Kaplan G, Swift R, and Reid N. Effects of contingency management and bupropion on cigarette smoking in smokers with schizophrenia. *Psychopharmacology*, 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21475970>

Sung H, Prochaska J, Ong M, Shi Y, and Max W. Cigarette smoking and serious psychological distress: A population-based study of california adults. *Nicotine & Tobacco Research*, 2011; 13(12):1183–92. Available from: <http://ntr.oxfordjournals.org/content/early/2011/08/17/ntr.ntr148.full>

Schechter M. Nicotine replacement therapy in the psychiatric emergency department. *The American Journal of Psychiatry*, 2011; 168(4):347–9. Available from: <http://ajp.psychiatryonline.org/cgi/content/full/168/4/347>

Prochaska JJ. Smoking and mental illness--breaking the link. *New England Journal of Medicine*, 2011; 365(3):196–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21774707>

Piper ME, Cook JW, Schlam TR, Jorenby DE, and Baker TB. Anxiety diagnoses in smokers seeking cessation treatment: Relations with tobacco dependence, withdrawal, outcome and response to treatment. *Addiction*, 2011; 106(2):418–27. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/20973856>

Okoli C, Khara M, Torchalla I, Ensom M, Oliffe J, et al. Sex differences in smoking cessation outcomes of a tailored program for individuals with substance use disorders and mental illness. *Addictive Behaviors*, 2011; [Epub ahead of print]. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/21277095>

Ohta C, Yasui-Furukori N, Furukori H, Tsuchimine S, Saito M, et al. The effect of smoking status on the plasma concentration of prolactin already elevated by risperidone treatment in schizophrenia patients. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21216266>

Morris C, Waxmonsky J, May M, Tinkelman D, Dickinson M, et al. Smoking reduction for persons with mental illnesses: 6-month results from community-based interventions. *Community Mental Health Journal*, 2011; [Epub ahead of print]. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/21556784>

Morgan V WA, Jablensky A, Mackinnon A, McGrath J, Carr V, Bush R, Castle D, Cohen M, Harvey C, Galletly C, Stain H, Neil A, McGorry P, Hocking B, Shah S, Saw S. People living with psychotic illness 2010: Report on the second Australian national survey. 2011. Available from:

<http://www.health.gov.au/internet/main/publishing.nsf/Content/mental-pubs-p-psych10>

Molina-Linde J. Effectiveness of smoking cessation programs for seriously mentally ill. *Actas Espanolas de Psiquiatria*, 2011; 39(2):106–14. Available from:

<http://actaspsiquiatria.es/repositorio/13/70/ESP/13-70-ESP-106-114-494933.pdf>

McNally L, Todd C, and Ratschen E. The prevalence of mental health problems among users of nhs stop smoking services: Effects of implementing a routine screening procedure. *BMC Health Services Research*, 2011; 11(1):190. Available from: <http://www.biomedcentral.com/1472-6963/11/190>

McClernon F, Van Voorhees E, English J, Hallyburton M, Holdaway A, et al. Smoking withdrawal symptoms are more severe among smokers with adhd and independent of adhd symptom change: Results from a 12-day contingency-managed abstinence trial. *Nicotine & Tobacco Research*, 2011; [Epub ahead of print]. Available from:

<http://ntr.oxfordjournals.org/content/early/2011/05/12/ntr.ntr073.full>

Matthews A, Wilson V, and Mitchell S. The role of antipsychotics in smoking and smoking cessation. *CNS Drugs*, 2011; 25(4):299–315. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=21425883

Mathews R, Hall W, and Gartner C. Depression and psychological distress in tobacco smokers and people with cannabis dependence in the national survey of mental health and wellbeing. *The Medical Journal of Australia*, 2011; 195(3):S12–5. Available from:

http://www.mja.com.au/public/issues/195_03_010811/mat10915_fm.html

Maremmanni I, Perugi G, Rovai L, Maremmanni A, Pacini M, et al. Are "social drugs" (tobacco, coffee and chocolate) related to the bipolar spectrum? *Journal of Affective Disorders*, 2011; 133(1-2):227–33. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21605911>

Liu M, Tsai S, Jeang S, Peng S, Wu S, et al. Varenicline prevents affective and cognitive exacerbation during smoking abstinence in male patients with schizophrenia. *Psychiatry Research*, 2011; [Epub ahead of print]. Available from: <http://www.sciencedirect.com/science/article/pii/S0165178111003490>

Lin M, Burgess J, Jr, and Carey K. The association between serious psychological distress and emergency department utilization among young adults in the USA. *Social Psychiatry and Psychiatric Epidemiology*, 2011; [Epub ahead of print]. Available from: <https://springerlink3.metapress.com/content/el55718687184140/resource-secured/?target=fulltext.pdf&sid=nhvqdh551bjhvt455djz53mu&sh=www.springerlink.com>

Leventhal A, Zvolensky M, and Schmidt N. Smoking-related correlates of depressive symptom dimensions in treatment-seeking smokers. *Nicotine & Tobacco Research*, 2011; 13(8):668–76. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21471305>

Lee SS, Humphreys KL, Flory K, Liu R, and Glass K. Prospective association of childhood attention-deficit/hyperactivity disorder (adhd) and substance use and abuse/dependence: A meta-analytic review. *Clin Psychol Rev*, 2011; 31(3):328–41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21382538>

Lawn S, Hersh D, Ward PR, Tsourtos G, Muller R, et al. 'I just saw it as something that would pull you down, rather than lift you up': Resilience in never-smokers with mental illness. *Health Education Research*, 2011; 26(1):26–38. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21062967>

Hong L, Schroeder M, Ross T, Buchholz B, Salmeron B, et al. Nicotine enhances but does not normalize visual sustained attention and the associated brain network in schizophrenia. *Schizophrenia Bulletin*, 2011; 37(2):416–25. Available from: <http://schizophreniabulletin.oxfordjournals.org/content/37/2/416.long>

Hitsman B, Buka S, Veluz-Wilkins A, Mohr D, Niaura R, et al. Accuracy of a brief screening scale for lifetime major depression in cigarette smokers. *Psychology of Addictive Behaviors*, 2011; 25(3):559–64. Available from: https://my.apa.org/apa/idm/login.seam?ERIGHTS_TARGET=http%3A%2F%2Fpsycnet.apa.org%2Fpsycarticles%2F2011-06251-001.pdf&AUTHENTICATION_REQUIRED=true

Herrera M and McChargue D. Mood and cue reactivity among smokers with a history of major depression: The role of rumination and impulsivity. *American Journal on Addictions*, 2011; 20(2):161–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21314759>

Hebert K, Cummins S, Hernandez S, Tedeschi G, and Zhu S. Current major depression among smokers using a state quitline. *American Journal of Preventive Medicine*, 2011; 40(1):47–53. Available from: <http://www.ajpm-online.net/article/S0749-3797%2810%2900555-6/fulltext>

Gehricke J, Hong N, Wigal T, Chan V, and Doan A. Adhd medication reduces cotinine levels and withdrawal in smokers with adhd. *Pharmacology, Biochemistry, and Behavior*, 2011; 98(3):485–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21356232>

Gahr M, Freudenmann R, and Schonfeldt-Lecuona C. Non-suicidal self-injury by intravenous application of nicotine in a patient with borderline personality disorder resulting in substance dependence. *Addiction*, 2011; 106(6). Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1360-0443.2011.03393.x/pdf>

Dickerson F, Bennett M, Dixon L, Burke E, Vaughan C, et al. Smoking cessation in persons with serious mental illnesses: The experience of successful quitters. *Psychiatric Rehabilitation Journal*, 2011; 34(4):311–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21459747>

D'Souza M and Markou A. Schizophrenia and tobacco smoking comorbidity: Nachr agonists in the treatment of schizophrenia-associated cognitive deficits. *Neuropharmacology*, 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21288470>

Chen X, Li C, Smith R, Xiao Z, and Wang J. Differential sensory gating functions between smokers and non-smokers among drug-naïve first episode schizophrenic patients. *Psychiatry Research*, 2011; 188(3):327–33. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21216472>

Brunette M, Ferron J, McHugo G, Davis K, Devitt T, et al. An electronic decision support system to motivate people with severe mental illnesses to quit smoking. *Psychiatric Services*, 2011; 62(4):360–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21459986>

Bolam B, West R, and Gunnell D. Does smoking cessation cause depression and anxiety? Findings from the attempt cohort. *Nicotine & Tobacco Research*, 2011; 13(3):209–14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21330275>

Berlin I, Covey L, Donohue M, and Agostiv V. Duration of smoking abstinence and suicide related outcomes. *Nicotine & Tobacco Research*, 2011; [Epub ahead of print]. Available from: <http://ntr.oxfordjournals.org/content/early/2011/05/25/ntr.ntr089.full>

Wye P, Bowman J, Wiggers J, Baker A, Carr V, et al. An audit of the prevalence of recorded nicotine dependence treatment in an Australian psychiatric hospital. *Australian and New Zealand Journal of Public Health*, 2010; 34(3):298–303. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20618273>

Wye P, Bowman J, Wiggers J, Baker A, Carr V, et al. Providing nicotine dependence treatment to psychiatric inpatients: The views of Australian nurse managers. *J Psychiatr Ment Health Nurs*, 2010; 17(4):319–27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20529182>

Winhusen TM, Somoza EC, Brigham GS, Liu DS, Green CA, et al. Impact of attention-deficit/hyperactivity disorder (adhd) treatment on smoking cessation intervention in adhd smokers: A randomized, double-blind, placebo-controlled trial. *Journal of Clinical Psychiatry*, 2010; 71(12):1680–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20492837>

Welfare AloHa. Australian institute of health and welfare 2010. Australia's health 2010. Australia's health series. 2010. Available from: <http://www.aihw.gov.au/publication-detail/?id=6442468376>

Uehara K, Sone R, and Yamazaki F. Cigarette smoking following a prolonged mental task exaggerates vasoconstriction in glabrous skin in habitual smokers. *Journal of UOEH*, 2010; 32(4):303–16. Available from: http://www.uoeh-u.ac.jp/kouza/journal/intro_e.html

Tsoi DT, Porwal M, and Webster AC. Efficacy and safety of bupropion for smoking cessation and reduction in schizophrenia: Systematic review and meta-analysis. *British Journal of Psychiatry*, 2010; 196(5):346–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20435957>

Tordeurs D, Zdanowicz N, and Reynaert C. Various facets of anxiety in nicotinic consumption and cessation. *Psychiatria Danubina*, 2010; 22 Suppl 1:S167. Available from: http://www.unboundmedicine.com/medline/ebm/record/21057436/full_citation/Various_facets_of_anxiety_in_nicotinic_consumption_and_cessation

Takeuchi T, Nakao M, Shinozaki Y, and Yano E. Validity of self-reported smoking in schizophrenia patients. *Psychiatry and Clinical Neurosciences*, 2010; 64(3):274–8. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1440-1819.2010.02082.x/full>

Strong D, Cameron A, Feuer S, Cohn A, Abrantes A, et al. Single versus recurrent depression history: Differentiating risk factors among current US smokers. *Drug and Alcohol Dependence*, 2010; [Epub ahead of print]. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=20074868

Steinberg M, Williams J, Gandhi K, Foulds J, and Brandon T. Lower task persistence in smokers with schizophrenia as compared to non-psychiatric control smokers. *Psychology of Addictive Behaviors*, 2010; 24(4):724–9. Available from: <http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2010-26520-014>

Sonne S, Nunes E, Jiang H, Tyson C, Rotrosen J, et al. The relationship between depression and smoking cessation outcomes in treatment-seeking substance abusers. *American Journal on Addictions*, 2010; 19(2):111–8. Available from: <http://www3.interscience.wiley.com/user/accessdenied?ID=123249311&Act=2138&Code=4719&Page=/cgi-bin/fulltext/123249311/HTMLSTART>

Siru R, Hulse GK, Khan RJ, and Tait RJ. Motivation to quit smoking among hospitalised individuals with and without mental health disorders. *Australian and New Zealand Journal of Psychiatry*, 2010; 44(7):640–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20560851>

Shah A and Bhatia V. The relationship between elderly suicide rates and smoking in England and Wales. *Int Psychogeriatr*, 2010; 22(3):504–7. Available from: <http://journals.cambridge.org/action/displayAbstract?fromPage=online&aid=7453624>

Roberts SJ, Glod CA, Kim R, and Houchell J. Relationships between aggression, depression, and alcohol, tobacco: Implications for healthcare providers in student health. *Journal of the American Academy of Nurse Practitioners*, 2010; 22(7):369–75. Available from:

<http://www3.interscience.wiley.com/user/accessdenied?ID=123563481&Act=2138&Code=4719&Page=/cgi-bin/fulltext/123563481/HTMLSTART>

Prochaska JJ. Failure to treat tobacco use in mental health and addiction treatment settings: A form of harm reduction? *Drug and Alcohol Dependence*, 2010; 110(3):177–82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20378281>

Prochaska J. Integrating tobacco treatment into mental health settings. *Journal of the American Medical Association*, 2010; 304(22):2534–5. Available from: <http://jama.ama-assn.org/content/304/22/2534.long>

Pratt L and Brody D. Depression and smoking in the u.S. Household population aged 20 and over, 2005–2008. *NCHS Data Brief*, 2010; (34):1–8. Available from: <http://www.cdc.gov/nchs/data/databriefs/db34.pdf>

Munafo M and Araya R. Cigarette smoking and depression: A question of causation. *The British Journal of Psychiatry*, 2010; 196(6):425–6. Available from: <http://bjp.rcpsych.org/cgi/content/full/196/6/425>

Moss T, Weinberger A, Vessicchio J, Mancuso V, Cushing S, et al. A tobacco reconceptualization in psychiatry: Toward the development of tobacco-free psychiatric facilities. *American Journal on Addictions*, 2010; 19(4):293–311. Available from: <http://www3.interscience.wiley.com/user/accessdenied?ID=123469557&Act=2138&Code=4719&Page=/cgi-bin/fulltext/123469557/HTMLSTART>

Miller G and Dixon L. Half full or half empty? *Psychiatric Services*, 2010; 61(9):859. Available from: <http://www.nature.com/bjc/journal/vaop/ncurrent/full/6605842a.html>

McNally L and Ratschen E. The delivery of stop smoking support to people with mental health conditions: A survey of nhs stop smoking services. *BMC Health Services Research*, 2010; 10(1):179. Available from: <http://www.biomedcentral.com/content/pdf/1472-6963-10-179.pdf>

McFall M, Saxon AJ, Malte CA, Chow B, Bailey S, et al. Integrating tobacco cessation into mental health care for posttraumatic stress disorder: A randomized controlled trial. *Journal of the American Medical Association*, 2010; 304(22):2485–93. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21139110>

Le Strat Y, Ramoz N, and Gorwood P. In alcohol-dependent drinkers, what does the presence of nicotine dependence tell us about psychiatric and addictive disorders comorbidity? *Alcohol and Alcoholism*, 2010; 45(2):167–72. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/45/2/167>

Lawrence D, Considine J, Mitrou F, and Zubrick S. Anxiety disorders and cigarette smoking: Results from the Australian survey of mental health and wellbeing. *The Australian and New Zealand Journal of Psychiatry*, 2010; 44(6):520–7. Available from: <http://informahealthcare.com/doi/abs/10.3109/00048670903571580>

Korhonen T, Prince van Leeuwen A, Reijneveld SA, Ormel J, Verhulst FC, et al. Externalizing behavior problems and cigarette smoking as predictors of cannabis use: The trails study. *J Am Acad Child*

Adolesc Psychiatry, 2010; 49(1):61–9. Available from:

<http://www.jaacap.com/article/PIIS0890856709000033/fulltext>

Kollins SH, McClernon FJ, and Van Voorhees EE. Monetary incentives promote smoking abstinence in adults with attention deficit hyperactivity disorder (adhd). Experimental and Clinical Psychopharmacology, 2010; 18(3):221–8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/20545386>

Knuts I, Cosci F, Esquivel G, Goossens L, van Duinen M, et al. Cigarette smoking and 35% co(2) induced panic in panic disorder patients. Journal of Affective Disorders, 2010; 124(1–2):215–8.

Available from: www.ncbi.nlm.nih.gov/pubmed/19896718

Kapson H and Haaga D. Depression vulnerability moderates the effects of cognitive behavior therapy in a randomized controlled trial for smoking cessation. Behavior Therapy, 2010; 41(4):447–60.

Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=21035610

Kang E and Lee J. A longitudinal study on the causal association between smoking and depression.

Journal of Preventive Medicine and Public Health, 2010; 43(3):193–204. Available from:

http://www.healthis.org/search_jour/pdf_view.html?code=pv&year=2010&page=193

Johnson J, Moffat B, and Malchy L. In the shadow of a new smoke free policy: A discourse analysis of health care providers' engagement in tobacco control in community mental health. International Journal of Mental Health Systems, 2010; 4(1):23. Available from:

<http://www.ijmhs.com/content/4/1/23>

Haustein K-O and Groneberg D. Other organ systems. Tobacco or Health? 2010. p 149–220 Available from:

<https://springerlink3.metapress.com/content/p18h677167t70212/resource-secured/?target=fulltext.html&sid=p5vwkv45auw2nlzc4dirmfbz&sh=www.springerlink.com>.

Griffiths M, Kidd S, Pike S, and Chan J. The tobacco addiction recovery program: Initial outcome findings. Archives of Psychiatric Nursing, 2010; 24(4):239–46. Available from:

<http://www.psychiatricnursing.org/article/PIIS0883941709000995/fulltext>

Goldberg JO. Successful change in tobacco use in schizophrenia. Journal of the American Psychiatric Nurses 2010; 16(1):30–5. Available from:

<http://jap.sagepub.com/cgi/reprint/16/1/30>

Gierisch J, Bastian L, Calhoun P, McDuffie J, Williams J, et al. Comparative effectiveness of smoking cessation treatments for patients with depression: A systematic review and meta-analysis of the evidence. Department of Veterans Affairs, Veterans Health Administration, and Health Services Research & Development Service, Editors. 2010. Available from:

<http://www.ncbi.nlm.nih.gov/books/NBK51226/pdf/TOC.pdf>.

Fountoulakis K, Siamouli M, Panagiotidis P, Magiria S, Kantartzis S, et al. Obesity and smoking in patients with schizophrenia and normal controls: A case-control study. Psychiatry Research, 2010; 176(1):13–6. Available from:

www.ncbi.nlm.nih.gov/pubmed/20079934

Fernandez Del Rio E, Lopez A, and Becona E. Smoking cessation: Case study of a client with probable borderline personality disorder. *Psychological Reports*, 2010; 106(3):918–26. Available from: <http://pubget.com/paper/20712180>

Fernandez Del Rio E, Lopez A, and Becona E. Personality disorders and premature dropout from psychological treatment for smoking cessation. *Psychological Reports*, 2010; 106(3):679–84. Available from: http://www.unboundmedicine.com/medline/ebm/record/20712155/full_citation/Personality_disorders_and_premature_dropout_from_psychological_treatment_for_smoking_cessation

Dos Santos V, Migott A, Bau C, and Chatkin J. Tobacco smoking and depression: Results of a cross-sectional study. *The British Journal of Psychiatry*, 2010; 197:413–4. Available from: <http://bjp.rcpsych.org/cgi/content/full/197/5/413>

de Miguel Diez J, Esteban y Pena M, Puente Maestu L, Hernandez Barrera V, Carrasco Garrido P, et al. Relationship between tobacco consumption and health-related quality of life in adults living in a large metropolitan area. *Lung*, 2010; 188(5):393–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20721573>

Dalby R, Chakravarty M, Ahdidan J, Sorensen L, Frandsen J, et al. Localization of white-matter lesions and effect of vascular risk factors in late-onset major depression. *Psychological Medicine*, 2010; 40(8):1389–99. Available from: <http://journals.cambridge.org/action/displayAbstract?fromPage=online&aid=7826354>

Covey LS, Hu MC, Winhusen T, Weissman J, Berlin I, et al. Oros-methylphenidate or placebo for adult smokers with attention deficit hyperactivity disorder: Racial/ethnic differences. *Drug and Alcohol Dependence*, 2010; 110(1-2):156–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20219292>

Cohn A, Strong D, Abrantes A, and Brown R. History of single episode and recurrent major depressive disorder among smokers in cessation treatment: Associations with depressive symptomatology and early cessation failure. *Addictive Disorders & Their Treatment*, 2010; 9(1):41–52. Available from: http://journals.lww.com/addictiondisorders/Abstract/2010/03000/History_of_Single_Episode_and_Recurrent_Major.5.aspx

Boden J, Fergusson D, and Horwood L. Cigarette smoking and depression: Tests of causal linkages using a longitudinal birth cohort. *The British Journal of Psychiatry*, 2010; 196(6):440–6. Available from: <http://bjp.rcpsych.org/cgi/content/full/196/6/440>

Banham L and Gilbody S. Smoking cessation in severe mental illness: What works? *Addiction*, 2010; 105(7):1176–89. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20491721>

Baker A, Richmond R, Lewin TJ, and Kay-Lambkin F. Cigarette smoking and psychosis: Naturalistic follow up 4 years after an intervention trial. *Australian and New Zealand Journal of Psychiatry*, 2010; 44(4):342–50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20307166>

[No authors listed]. For the patient. Do african american teens have equal access to nicotine patch and gum? *Ethnicity & Disease*, 2010; 20(2):195. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20503903>

Weinberger A, Maciejewski P, McKee S, Reutenauer E, and Mazure C. Gender differences in associations between lifetime alcohol, depression, panic disorder, and posttraumatic stress disorder and tobacco withdrawal. *American Journal on Addictions*, 2009; 18(2):140–7. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/10550490802544888>

Webb M and Carey M. The early health consequences of smoking: Relationship with psychosocial factors among treatment-seeking black smokers. *Nicotine & Tobacco Research*, 2009; 11(5):564–71. Available from: <http://ntr.oxfordjournals.org/cgi/content/full/11/5/564>

Vorspan F, Bloch V, Guillem E, Dupuy G, Pirnay S, et al. Smoking ban in a psychiatry department: Are nonsmoking employees less exposed to environmental tobacco smoke? *European Psychiatry*, 2009; 24(8):529–32. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19540729>

Stapleton J, Keaney F, and Sutherland G. Illicit drug use as a predictor of smoking cessation treatment outcome. *Nicotine and Tobacco Research* 2009; 11(6):685–9. Available from: <http://ntr.oxfordjournals.org/content/11/6/685.full>

St Gray K and Upadhyaya H. Tobacco smoking in individuals with attention-deficit hyperactivity disorder: Epidemiology and pharmacological approaches to cessation. *CNS Drugs*, 2009; 23(8):661–8. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19594195

Solway E. Windows of opportunity for culture change round tobacco use in mental health settings *Journal of the American Psychiatric Nurses Association*, 2009; 15(1):41–9. Available from: <http://jap.sagepub.com/cgi/reprint/15/1/41>

Solty H, Crockford D, White W, and Currie S. Cigarette smoking, nicotine dependence, and motivation for smoking cessation in psychiatric inpatients. *Canadian Journal of Psychiatry*, 2009; 54(1):36–45. Available from: publications.cpa-apc.org/media.php?mid=729&xwm=true

Smith W. Housing: A human right. *New Paradigm*, 2009; Winter 2009:15–9. Available from: www.vicserv.org.au/uploads/documents/articles%20index.pdf

Smith G, Wong H, Macewan G, Kopala L, Ehmann T, et al. Predictors of starting to smoke cigarettes in patients with first episode psychosis. *Schizophrenia Research*, 2009; 108(1):258–64. Available from: <http://www.schres-journal.com/article/PIIS0920996408005598/fulltext>

Siru R, Hulse GK, and Tait RJ. Assessing motivation to quit smoking in people with mental illness: A review. *Addiction*, 2009; 104(5):719–33. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19413788>

Sharp DL, Blaakman SW, Cole RE, and Evinger JS. Report from a national tobacco dependence survey of psychiatric nurses. *Journal of the American Psychiatric Nurses Association*, 2009; 15(3):172–81. Available from: <http://jap.sagepub.com/cgi/reprint/15/3/172>

Sharp DL and Blaakman SW. Report from the apna tobacco dependence council Journal of the American Psychiatric Nurses Association, 2009; 15(6):412–4. Available from: <http://jap.sagepub.com/cgi/reprint/15/6/412>

Richmond R, Baker A, Lewin T, and Kay-Lambkin F. Smoking cessation intervention among people with a psychotic disorder: 4 year follow up. Abstract pa11-5. in 2009 Joint Conference of the Society for Research on Nicotine and Tobacco (SRNT) and SRNT-Europe. Dublin, Ireland. 2009. Available from: http://www.srnt.org/meeting/2009/pdf/2009_Paper_Sessions.pdf.

Ratschen E, Britton J, Doody G, and McNeill A. Smoke-free policy in acute mental health wards: Avoiding the pitfalls. General Hospital Psychiatry, 2009; 31(2):131–6. Available from: <http://www.ghpjournal.com/article/PIIS0163834308002132/fulltext>

Rao U, Hammen CL, and Poland RE. Drs. Rao, hammen, and Poland reply. The American Journal of Psychiatry, 2009; 166(7):831–a–2. Available from: <http://ajp.psychiatryonline.org/cgi/reprint/166/7/831-a>

Prochaska JJ. Ten critical reasons for treating tobacco dependence in inpatient psychiatry Journal of the American Psychiatric Nurses Association, 2009; 15(6):404–9. Available from: <http://jap.sagepub.com/cgi/reprint/15/6/404>

Pickett K, Wilkinson R, and Wakschlag L. The psychosocial context of pregnancy smoking & quitting in the millennium cohort study. Journal of Epidemiology and Community Health 2009; 63(6):474–80. Available from: <http://jech.bmj.com/content/63/6/474.long>

Naegle M, Baird C, and Farchaus-Stein K. Psychiatric nurses as champions for smoking cessation Journal of the American Psychiatric Nurses Association, 2009; 15(1):21–3. Available from: <http://jap.sagepub.com/cgi/reprint/15/1/21>

Morris CD, Tedeschi GJ, Waxmonsky JA, May M, and Giese AA. Tobacco quitlines and persons with mental illnesses: Perspective, practice, and direction Journal of the American Psychiatric Nurses Association, 2009; 15(1):32–40. Available from: <http://jap.sagepub.com/cgi/reprint/15/1/32>

Morisano D, Bacher I, Audrain-McGovern J, and George TP. Mechanisms underlying the comorbidity of tobacco use in mental health and addictive disorders. Can J Psychiatry, 2009; 54(6):356–67. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19527556>

McGrath JJ and Susser ES. New directions in the epidemiology of schizophrenia. Medical Journal of Australia, 2009; 190(S4):S7–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19220176>

McGrath D and Barrett S. The comorbidity of tobacco smoking and gambling: A review of the literature. Drug and Alcohol Review, 2009; 28(6):676–81. Available from: <http://www3.interscience.wiley.com/user/accessdenied?ID=122539709&Act=2138&Code=4719&Page=/cgi-bin/fulltext/122539709/HTMLSTART>

Marshall EC, Vujanovic AA, Kutz A, Gibson L, Leyro T, et al. Reasons for quitting smoking prior to a self-quit attempt among smokers with and without posttraumatic stress disorder or other anxiety/mood psychopathology. American Journal of Addiction, 2009; 18(4):309–15. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/10550490902925763>

Lising-Enriquez K and George T. Treatment of comorbid tobacco use in people with serious mental illness. *Journal of Psychiatry & Neuroscience*, 2009; 34(3):E1–2. Available from:

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pubmed&pubmedid=19448844>

Lima JE, Reid MS, Smith JL, Zhang Y, Jiang H, et al. Medical and mental health status among drug dependent patients participating in a smoking cessation treatment study. *Journal of Drug Issues*, 2009; 39(2). Available from: <http://www.encyclopedia.com/doc/1P3-1826120731.html>

Kim S, Han D, Joo S, and Min K. The effect of dopamine partial agonists on the nicotine dependency in patients with schizophrenia. *Human Psychopharmacology*, 2009; 25(2):187–90. Available from:

<http://onlinelibrary.wiley.com/doi/10.1002/hup.1089/pdf>

Khaled SM, Bulloch A, Exner DV, and Patten SB. Cigarette smoking, stages of change, and major depression in the canadian population. *Canadian Journal of Pyschiatry*, 2009; 54(3):204–8. Available from: publications.cpa-apc.org/media.php?mid=759&xwm=true

Keizer I, Descloux V, and Eytan A. Variations in smoking after admission to psychiatric inpatient units and impact of a partial smoking ban on smoking and on smoking-related perceptions. *Int J Soc Psychiatry*, 2009; 55(2):109–23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19240201>

Johnson JL, Malchy LA, Ratner PA, Hossain S, Procyshyn RM, et al. Community mental healthcare providers' attitudes and practices related to smoking cessation interventions for people living with severe mental illness. *Patient Education and Counseling*, 2009; 77(2):289–95. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/19398293>

Hitsman B, Moss TG, Montoya ID, and George TP. Treatment of tobacco dependence in mental health and addictive disorders. *Can J Psychiatry*, 2009; 54(6):368–78. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/19527557>

Higgins S and Chilcoat H. Women and smoking: An interdisciplinary examination of socioeconomic influences. *Drug and Alcohol Dependence*, 2009; 104(suppl 1.):S1–S5. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/19586726>

Gupta A and Craig T. Diet, smoking and cardiovascular risk in schizophrenia in high and low care supported housing. *Epidemiologia E Psichiatria Sociale*, 2009; 18(3):200–7. Available from:

http://www.eps-journal.com/articoli.php?archivio=yes&vol_id=446&id=5281

George T and Ziedonis D. Addressing tobacco dependence in psychiatric practice: Promises and pitfalls. *Canadian Journal of Psychiatry*, 2009; 54(6):353–5. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19527555

Gehricke J, Hong N, Whalen C, Steinhoff K, and Wigal T. Effects of transdermal nicotine on symptoms, moods, and cardiovascular activity in the everyday lives of smokers and nonsmokers with attention-deficit/hyperactivity disorder. *Psychology of Addictive Behaviors*, 2009; 23(4):644–55. Available from:

<http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2009-24023-010&CFID=27436988&CFTOKEN=17479918>

Fusar-Poli P, De Marco L, Cavallin F, Bertorello A, Nicolasi M, et al. Lifestyles and cardiovascular risk in individuals with functional psychoses. *Perspectives in Psychiatric Care*, 2009; 45(2):87–99.

Available from:

<http://www3.interscience.wiley.com/user/accessdenied?ID=122298580&Act=2138&Code=4719&Page=/cgi-bin/fulltext/122298580/HTMLSTART>

Fagerstrom K and Aubin HJ. Management of smoking cessation in patients with psychiatric disorders. *Current Medical Research and Opinion*, 2009; 25(2):511–8. Available from:

<http://www.informapharmascience.com/doi/abs/10.1185/03007990802707568>

Essenmacher C, Karvonen-Gutierrez C, Lynch-Sauer J, and Duffy SA. Staff's attitudes toward the delivery of tobacco cessation services in a primarily psychiatric veterans affairs hospital. *Archives of Psychiatric Nursing*, 2009; 23(3):231–42. Available from:

[http://www.psychiatricnursing.org/article/S0883-9417\(08\)00110-6/abstract](http://www.psychiatricnursing.org/article/S0883-9417(08)00110-6/abstract)

Dwyer T, Bradshaw J, and Happell B. Comparison of mental health nurses' attitudes towards smoking and smoking behaviour. *International Journal of Mental Health Nursing*, 2009; 18(6):424–33.

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19883414>

Dunn KE, Sigmon SC, Reimann E, Heil SH, and Higgins ST. Effects of smoking cessation on illicit drug use among opioid maintenance patients: A pilot study. *Journal of Drug Issues* 2009; 39(2). Available from:

<http://www.encyclopedia.com/doc/1P3-1826120741.html>

Diaz F, James D, Botts S, Maw L, Susce M, et al. Tobacco smoking behaviors in bipolar disorder: A comparison of the general population, schizophrenia, and major depression. *Bipolar Disorders*, 2009; 11(2):154–65. Available from:

<http://www3.interscience.wiley.com/user/accessdenied?ID=122218275&Act=2138&Code=4719&Page=/cgi-bin/fulltext/122218275/HTMLSTART>

de Ruiter M, Veltman D, Goudriaan A, Oosterlaan J, Sjoerds Z, et al. Response perseveration and ventral prefrontal sensitivity to reward and punishment in male problem gamblers and smokers. *Neuropsychopharmacology*, 2009; 34:1027–38. Available from:

<http://www.nature.com/npp/journal/vaop/ncurrent/full/npp2008175a.html>

Currie S and Crockford D. Authors' response to commentary *Journal of the American Psychiatric Nurses Association*, 2009; 15(6):410–11. Available from:

<http://jap.sagepub.com/cgi/reprint/15/6/410>

Conway J. Exogenous nicotine normalises sensory gating in schizophrenia; therapeutic implications. *Med Hypotheses*, 2009; 73(2):259–62. Available from:

www.ncbi.nlm.nih.gov/pubmed/19328631

Chapman S, Ragg M, and McGeechan K. Citation bias in reported smoking prevalence in people with schizophrenia. *Australian and New Zealand Journal of Psychiatry*, 2009; 43(3):277–82. Available from:

<http://www.informaworld.com/smpp/content~db=all?content=10.1080/00048670802653372>

Baker A, Richmond R, Castle D, Kulkarni J, Kay-Lambkin F, et al. Coronary heart disease risk reduction intervention among overweight smokers with a psychotic disorder: Pilot trial. *Australian and New*

Zealand Journal of Psychiatry, 2009; 43(2):129–35. Available from:

<http://www.informaworld.com/smpp/content~db=all?content=10.1080/00048670802607147>

380. Zvolensky MJ, Gibson LE, Vujanovic AA, Gregor K, Bernstein A, et al. Impact of posttraumatic stress disorder on early smoking lapse and relapse during a self-guided quit attempt among community-recruited daily smokers. *Nicotine & Tobacco Research*, 2008; 10(8):1415–27.

Available from:

<http://www.informaworld.com/smpp/content~content=a901415630~db=all~order=page>

Wilens TE, Adamson J, Monuteaux MC, Faraone SV, Schillinger M, et al. Effect of prior stimulant treatment for attention-deficit/hyperactivity disorder on subsequent risk for cigarette smoking and alcohol and drug use disorders in adolescents. *Archives of Pediatrics & Adolescent Medicine*, 2008; 162(10):916–21. Available from: <http://archpedi.ama-assn.org/cgi/content/full/162/10/916>

Wilens T, Biederman J, Adamson J, Henin A, Sgambati S, et al. Further evidence of an association between adolescent bipolar disorder with smoking and substance use disorders: A controlled study *Drug and Alcohol Dependence*, 2008; 95(3):188 – 98. Available from:

[http://linkinghub.elsevier.com/retrieve/pii/S0376-8716\(08\)00050-1](http://linkinghub.elsevier.com/retrieve/pii/S0376-8716(08)00050-1)

Weinberger AH, Vessicchio JC, Sacco KA, Creedon CL, Chengappa KN, et al. A preliminary study of sustained-release bupropion for smoking cessation in bipolar disorder. *J Clin Psychopharmacol*, 2008; 28(5):584–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18794666>

Weinberger A, Reutenauer E, Vessicchio J, and George T. Survey of clinician attitudes toward smoking cessation for psychiatric and substance abusing clients. *Journal of Addictive Diseases*, 2008; 27(1):55–63. Available from:

https://www.haworthpress.com/store/Toc_views.asp?sid=THPMHA1RBXQX9MVXDCB28EQFGV4DCV12&TOCName=J069v27n01%5FTOC&desc=Volume%3A%2027%20Issue%3A%201

Weinberger A, George T, Perkins K, and Chengappa K. Effects of topiramate on smoking in patients with schizoaffective disorder, bipolar type. *Journal of Clinical Psychopharmacology*, 2008; 28(2):247–8. Available from:

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=18344743

Tobias M, Templeton R, and Collings S. How much do mental disorders contribute to new zealand's tobacco epidemic? *Tobacco Control*, 2008; 17(5):347–50. Available from:

<http://tobaccocontrol.bmj.com/cgi/content/abstract/17/5/347>

Tidey JW, Rohsenow DJ, Kaplan GB, Swift RM, and Adolfo AB. Effects of smoking abstinence, smoking cues and nicotine replacement in smokers with schizophrenia and controls. *Nicotine & Tobacco Research*, 2008; 10(6):1047–56. Available from:

<http://www.informaworld.com/smpp/content~content=a794502342~db=all~order=page>

Snyder M, McDevitt J, and Painter S. Smoking cessation and serious mental illness. *Archives of Psychiatric Nursing*, 2008; 22(5):297–304. Available from:

<http://www.sciencedirect.com/science/journal/08839417>

Shmueli D, Fletcher L, Hall S, Hall S, and Prochaska J. Changes in psychiatric patients' thoughts about quitting smoking during a smoke-free hospitalization. *Nicotine & Tobacco Research*, 2008; 10(5):875–81. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18569762>

Ribeiro S, Jennen-Steinmetz C, Schmidt M, and Becker K. Nicotine and alcohol use in adolescent psychiatric inpatients: Associations with diagnoses, psychosocial factors, gender and age. *Nordic Journal of Psychiatry*, 2008; 10:1–8. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/08039480801984073>

Ragg M and Ahmed T, Smoke and mirrors: A review of the literature on smoking and mental illness. Tackling tobacco program research series no 1 Sydney: Cancer Council NSW; 2008. Available from: <http://www.cancerdirectory.com.au/resource/view?slug=Smoke-and-Mirrors-A-review-of-the-literature-on-smoking-and-mental-illness&page=1>.

Moeller-Saxone K. Cigarette smoking and interest in quitting among consumers at a psychiatric disability rehabilitation and support service in victoria. *Australian and New Zealand Journal of Public Health*, 2008; 32(5):479–81. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1753-6405.2008.00283.x/abstract>

Mobascher A and Winterer G. The molecular and cellular neurobiology of nicotine abuse in schizophrenia. *Pharmacopsychiatry*, 2008; 41(Suppl 1):S51–9. Available from: <http://www.thieme-connect.com/ejournals/html/pharmacology/doi/10.1055/s-2008-1081463>

McClernon FJ and Kollins SH. Adhd and smoking: From genes to brain to behavior. *Annals of the New York Academy of Sciences*, 2008; 1141:131–47. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18991955>

Martin C, Allan R, Fleming M, and Atkinson J. Mood and smoking in schizophrenia. *Journal of Psychiatric and Mental Health Nursing*, 2008; 15(9):722–7. Available from: <http://www3.interscience.wiley.com/journal/121430260/abstract>

Lyons M, Hitsman B, Xian H, Panizzon M, Jerskey B, et al. A twin study of smoking, nicotine dependence, and major depression in men. *Nicotine & Tobacco Research*, 2008; 10(1):97–108. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200701705332>

Lawn S. Tobacco control policies, social inequality and mental health populations: Time for a comprehensive treatment response. *Australian and New Zealand Journal of Psychiatry*, 2008; 42(5):353–6. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/00048670801961180>

Lasser KE and Boyd JW. Varenicline and smokers with mental illnesses. *Lancet*, 2008; 372(9645):1218–9. Available from: http://www.sciencedirect.com/science?_ob=MIimg&_imagekey=B6T1B-4TKB3T6-P-1&_cdi=4886&_user=10&_orig=browse&_coverDate=10%2F10%2F2008&_sk=996270354&view=c&wchp=dGLbVlz-zSkzV&md5=83e8701b4c0e304826b42b3b729c2d6b&ie=/sdarticle.pdf

Lal A and Siahpush M. The effect of smoke-free policies on electronic gaming machine expenditure in victoria, Australia. *Journal of Epidemiology and Community Health*, 2008; 62(1):11–5. Available from: <http://jech.bmj.com/cgi/content/abstract/62/1/11>

Kurdyak P, Cairney J, Sarnocinska-Hart A, Callahan R, and Strike C. The impact of a smoking cessation policy on visits to a psychiatric emergency department. *Canadian Journal of Psychiatry*, 2008; 53(11):779–82. Available from: <http://publications.cpa-apc.org/media.php?mid=697&xwm=true>

Kristi A, Creedena SC, Reutenauera EL, and George TP. Effects of galantamine on cognitive deficits in smokers and non-smokers with schizophrenia. *Schizophrenia Research*, 2008; 103(1):326–7. Available from: [http://www.schres-journal.com/article/S0920-9964\(08\)00234-X/abstract](http://www.schres-journal.com/article/S0920-9964(08)00234-X/abstract)

Kotov R, Guey L, Bromet E, and Schwartz J. Smoking in schizophrenia: Diagnostic specificity, symptom correlates, and illness severity. *Schizophrenia Bulletin*, 2008; 36(1):173– 81. Available from: <http://schizophreniabulletin.oxfordjournals.org/content/36/1/173.full>

Kodl MM, Fu SS, Willenbring ML, Gravely A, Nelson DB, et al. The impact of depressive symptoms on alcohol and cigarette consumption following treatment for alcohol and nicotine dependence. *Alcoholism: Clinical and Experimental Research*, 2008; OnlineEarly Articles. Available from: <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1530-0277.2007.00556.x>

Kisely S and Campbell L. Use of smoking cessation therapies in individuals with psychiatric illness: An update for prescribers. *CNS Drugs*, 2008; 22(4):263–73. Available from: <http://www.ingentaconnect.com/content/adis/cns/2008/00000022/00000004/art00001?token=003f1af1055929990441333c4a2f24416a433b316721482867444f6d4e2224b>

Kinnunen T, Korhonen T, and Garvey A. Role of nicotine gum and pretreatment depressive symptoms in smoking cessation: Twelve-month results of a randomized placebo controlled trial. *The International Journal of Psychiatry in Medicine* 2008; 38(3):373–89. Available from: <http://baywood.metapress.com/app/home/contribution.asp?referrer=parent&backto=searcharticle&results,1,1;>

Ker S and Owens D. Admission to a psychiatric unit and changes in tobacco smoking. *Clinical Practice and Epidemiology in Mental Health* 2008; 4(12). Available from: <http://www.cpementalhealth.com/content/4/1/12>

Jenkins R, Bhugra D, Bebbington P, Brugha T, Farrell M, et al. Debt, income and mental disorder in the general population. *Psychological Medicine*, 2008; 38(10):1485–93. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18184442>

Haas A, Sorensen J, Hall S, Lin C, Delucchi K, et al. Cigarette smoking in opioid-using patients presenting for hospital-based medical services. *American Journal on Addictions*, 2008; 17(1):65–9. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/10550490701756112>

Government Media Office Ministerial media statements: Kalgoorlie mental health care in good hands. 2008. Available from:

<http://www.mediastatements.wa.gov.au/Pages/WACabinetMinistersSearch.aspx?ItemId=130903&minister=Jacobs&admin=Barnett>

Goldstein B, Birmaher B, Axelson D, Goldstein T, Esposito-Smythers C, et al. Significance of cigarette smoking among youths with bipolar disorder. *American Journal on Addictions*, 2008; 17(5):364–71. Available from:

<http://www.informaworld.com/smpp/content~db=all?content=10.1080/10550490802266151>

Garcia-Portilla M, Saiz P, Bascaran M, Martinez A, Benabarre A, et al. Cardiovascular risk in patients with bipolar disorder. *Journal of Affective Disorders*, 2008; 115(3):302–8. Available from:

<http://www.jad-journal.com/article/PIIS0165032708003650/fulltext>

Friedman-Wheeler D, Haaga D, Gunthert K, Ahrens A, and McIntosh E. Depression, neuroticism, and mood-regulation expectancies for engagement and disengagement coping among cigarette smokers. *Cognitive Therapy and Research*, 2008; 32(1):105–15. Available from:

<http://www.springerlink.com/content/j12l4g03106q8180/fulltext.pdf>

Evins A, Culhane M, Alpert J, Pava J, Liese B, et al. A controlled trial of bupropion added to nicotine patch and behavioral therapy for smoking cessation in adults with unipolar depressive disorders. *Journal of Clinical Psychopharmacology*, 2008; 28(6):660–6. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/19011435?ordinalpos=5&itool=EntrezSystem>

Diaz F, Velasquez D, Susce M, and de Leon J. The association between schizophrenia and smoking: Unexplained by either the illness or the prodromal period. *Schizophrenia Research*, 2008; 104(1–3):214–9. Available from: <http://www.schres-journal.com/article/PIIS0920996408002831/fulltext>

Culhane M, Schoenfeld D, Barr R, Cather C, Deckersbach T, et al. Predictors of early abstinence in smokers with schizophrenia. *Journal of Clinical Psychiatry*, 2008; 69(11):1743–50. Available from:

http://article.psychiatrist.com/dao_1-login.asp?ID=10003783&RSID=98999193354782

Covey L, Manubay J, Jiang H, Nortick M, and Palumbo D. Smoking cessation and inattention or hyperactivity/impulsivity: A post hoc analysis. *Nicotine & Tobacco Research*, 2008; 10(12):1717–25. Available from:

<http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802443536>

Cooper TV, Hanson BS, Burke RS, and Hunt YM. Sequential treatment for nicotine dependence and postcessation weight concern in a female diagnosed with paranoid schizophrenia. *Clinical Case Studies*, 2008; 7(4):340–55. Available from: <http://ccs.sagepub.com/cgi/reprint/7/4/340>

Cocores J. Transdermal nicotine in adult adhd with depression and anxiety. *Primary Care Companion to the Journal of Clinical Psychiatry*, 2008; 10(3):253 – 4. Available from:

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pubmed&pubmedid=18615164>

Chaves L and Shirakawa I. Nicotine use in patients with schizophrenia evaluated by the fagerström tolerance questionnaire: A descriptive analysis from a brazilian sample. *Revista Brasileira de Psiquiatria*, 2008:S1516–44462008005000014. Available from:

http://www.scielo.br/scielo.php?script=sci_pdf&pid=S1516-44462008005000014&lng=en&nrm=iso&tling=en

Chapman S. Questionable inflated estimate of smoking prevalence among mentally ill persons in Australia. *Australian and New Zealand Journal of Psychiatry*, 2008; 42(7):646 – 8. Available from: http://www.unboundmedicine.com/medline/ebm/record/18623628/full_citation/Questionable_inflated_estimate_of_smoking_prevalence_among_mentally_ill_persons_in_Australia

Campion J, Checinski K, Nurse J, and McNeill A. Smoking by people with mental illness and benefits of smoke-free mental health services. *Advances in Psychiatric Treatment*, 2008; 14:217–28. Available from: <http://apt.rcpsych.org/cgi/content/abstract/14/3/217>

Campion J, Checinski K, and Nurse J. Review of smoking cessation treatments for people with mental illness. *Advances in Psychiatric Treatment*, 2008; 14:208–16. Available from: <http://apt.rcpsych.org/cgi/reprint/14/3/208>

Campayo JG, Sobradie N, Alda M, Mas A, Andrés E, et al. Effectiveness of topiramate for tobacco dependence in patients with depression; a randomised, controlled trial. *BMC Family Practice*, 2008; 9(28). Available from: <http://www.biomedcentral.com/1471-2296/9/28>

Buckner JD and Schmidt NB. Marijuana effect expectancies: Relations to social anxiety and marijuana use problems *Addictive Behaviors*, 2008; 33(11):1477–83. Available from: http://www.sciencedirect.com/science?_ob=MIimg&_imagekey=B6VC9-4SWN0VF-2-1&_cdi=5949&_user=10&_orig=browse&_coverDate=11%2F30%2F2008&_sk=999669988&view=c&_wchp=dGLbVlb-zSkWb&md5=cd09850e5d6924db6f1101a6f937c428&ie=/sdarticle.pdf

Borras L, Mohr S, Brandt P, Gillieron C, Eytan A, et al. Influence of spirituality and religiousness on smoking among patients with schizophrenia or schizo-affective disorder in Switzerland. *International Journal of Social Psychiatry*, 2008; 54(6):539–49. Available from: <http://isp.sagepub.com/cgi/reprint/54/6/539>

Beckham J, Wiley M, Miller S, Dennis M, Wilson S, et al. Ad lib smoking in post-traumatic stress disorder: An electronic diary study. *Nicotine and Tobacco Research* 2008; 10(7):1149–57. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802123302>

Barnett PG, Wong W, and Hall S. The cost-effectiveness of a smoking cessation program for out-patients in treatment for depression. *Addiction*, 2008; 103(5):834–40. Available from: <http://dx.doi.org/10.1111/j.1360-0443.2008.02167.x>

Williams JM and Foulds J. Successful tobacco dependence treatment in schizophrenia. *American Journal of Psychiatry*, 2007; 164(2):222–7; quiz 373. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17267783>

SANE Australia. Smoking and mental illness: Costs. Access Economics Melbourne, 2007. Available from: <http://www.sane.org/information/research>

Quit Group. Minimum standards for Australian quitline services, 2007.

Myers M, Doran N, and Brown S. Is cigarette smoking related to alcohol use during the 8 years following treatment for adolescent alcohol and other drug abuse? *Alcohol and Alcoholism* 2007; 42(3):226–33. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/42/3/226>

Little H. Introduction to special issue of alcohol and alcoholism on alcohol and nicotine. *Alcohol and Alcoholism*, 2007; 42(3):165–6. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/42/3/165>

Leeman R, Huffman C, and O'Malley S. Alcohol history and smoking cessation in nicotine replacement therapy, bupropion sustained release and varenicline trials: A review. *Alcohol and Alcoholism*, 2007; 42(3):196 – 206. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/42/3/196>

Hughes J. Depression during tobacco abstinence. *Nicotine & Tobacco Research*, 2007; 9(4):443–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17454698>

Heffner J, Barrett S, and Anthenelli R. Predicting alcohol misusers' readiness and ability to quit smoking: A critical review. *Alcohol and Alcoholism*, 2007; 42(3):186–95. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/42/3/186>

Goi N, Hirai Y, Harada H, Ikari A, Ono T, et al. Comparison of peroxidase response to mental arithmetic stress in saliva of smokers and non-smokers. *Journal of Toxicological Sciences*, 2007; 332(2):121 – 7. Available from: http://www.jstage.jst.go.jp/article/jts/32/2/121/_pdf

Gallagher S, Penn P, Schindler E, and Layne W. A comparison of smoking cessation treatments for persons with schizophrenia and other serious mental illnesses *Journal of Psychoactive Drugs*, 2007; 39(4):487–97. Available from: <http://www.drDave.org/Articles/Journal-Of-PsychedelicDrugs.htm>

Fuller BE, Guydish J, Tsoh J, Reid MS, Resnick M, et al. Attitudes toward the integration of smoking cessation treatment into drug abuse clinics. *Journal of Substance Abuse Treatment*, 2007; 32(1):53–60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17175398>

Friend K and Pagano M. Timevarying predictors of smoking cessation among individuals in treatment for alcohol abuse and dependence: Findings from project match. *Alcohol and Alcoholism* 2007; 42(3):234 – 40. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/42/3/234>

Friedman-Wheeler D, Ahrens A, Haaga D, McIntosh E, and Thorndike F. Depressive symptoms, depression proneness, and outcome expectancies for cigarette smoking. *Cognitive Therapy and Research*, 2007; 31(4):547–57. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pubmed&pubmedid=18074003>
<http://www.pubmedcentral.nih.gov/picrender.fcgi?artid=2131705&blobtype=pdf>

Evins A, Cather C, Culhane M, Birnbaum A, Horowitz J, et al. A 12-week double-blind, placebo-controlled study of bupropion sr added to high-dose dual nicotine replacement therapy for smoking cessation or reduction in schizophrenia. *Journal of Clinical Psychopharmacology*, 2007; 27(4):380–6. Available from: <http://www.psychopharmacology.com/pt/re/jclnpsychopharm/abstract.00004714-200708000-00010.htm;jsessionid=LtJJ3LChv5rsTS77Tg2L6LhtgXjXsKGDJXyJnKYZth7f612L9hrM!-1912992316!181195628!8091!-1>

De Leon J, Rendon D, Baca-Garcia E, Aizpuru F, Gonzalez-Pinto A, et al. Association between smoking and alcohol use in the general population: Stable and unstable odds ratios across two years in two different countries. *Alcohol and Alcoholism*, 2007; 42(3):252–7. Available from:

<http://alcalc.oxfordjournals.org/cgi/content/full/42/3/252>

Bauld L, Ferguson J, Kerr A, McKenzie K, and McNeill A. Moving towards smoke-free mental health services. *Mental Health Review Journal* 2007; 12(2):18–24.

Baker A, Richmond R, Haile M, Lewin TJ, Carr VJ, et al. Characteristics of smokers with a psychotic disorder and implications for smoking interventions. *Psychiatry Res*, 2007; 150(2):141–52. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17289155>

Australian Bureau of Statistics, 4326.0 - national survey of mental health and wellbeing. Canberra 2007. Available from: <http://www.abs.gov.au/ausstats/abs@.nsf/mf/4326.0>.

Access Economics. Smoking and mental illness: Costs., Melbourne: SANE Australia, 2007. Available from: http://www.sane.org/images/stories/information/research/0712_info_smokecosts.pdf.

Williams JM and Ziedonis DM. Snuffing out tobacco dependence. Ten reasons behavioral health providers need to be involved. *Behav Healthc*, 2006; 26(5):27–31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16736916>

Wilhelm K, Wedgwood L, Niven H, and Kay-Lambkin F. Smoking cessation and depression: Current knowledge and future directions. *Drug and Alcohol Review*, 2006; 25(1):97–107. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16492582>

Thorndike FP, Wernicke R, Pearlman MY, and Haaga DA. Nicotine dependence, ptsd symptoms, and depression proneness among male and female smokers. *Addictive Behaviors*, 2006; 31(2):223–31. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15919158

Prochaska JJ, Fletcher L, Hall SE, and Hall SM. Return to smoking following a smoke-free psychiatric hospitalization. *American Journal on Addictions*, 2006; 15(1):15–22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16449089>

Lawn S and Condon J. Psychiatric nurses' ethical stance on cigarette smoking by patients: Determinants and dilemmas in their role in supporting cessation. *International Journal of Mental Health Nursing*, 2006; 15(2):111–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16643346>

Flick L, Cook C, Homan S, McSweeney M, Campbell C, et al. Persistent tobacco use during pregnancy and the likelihood of psychiatric disorders. *American Journal of Public Health* 2006; 96(10):1799–807. Available from: <http://www.ajph.org/cgi/content/abstract/96/10/1799>

Baker A, Richmond R, Haile M, TJ L, Carr V, et al. A randomised controlled trial of a smoking cessation intervention among people with a psychiatric disorder: Brief report of a pilot study. *American Journal of Psychiatry*, 2006; 163:1934–42.

Kollins SH, McClernon FJ, and Fuemmeler BF. Association between smoking and attention-deficit/hyperactivity disorder symptoms in a population-based sample of young adults. *Arch Gen Psychiatry*, 2005; 62(10):1142–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16203959>

Kalman D, Morissette S, and George T. Co-morbidity of smoking in patients with psychiatric and substance use disorders. *American Journal on Addictions*, 2005; 14:106–23. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pubmed&pubmedid=16019961>

Bertossa S. You can do it! A positive approach to supporting people to address tobacco. The Mental Health Services Conference, Sydney, 2005.

Ashton M. Policy within mental health services: Pro-actively addressing tobacco. The Mental Health Service Conference, 2005.

Wilhelm K, Richmond R, and Wodak A. Clinical aspects of nicotine dependence and depression. *Medicine Today*, 2004; 3:40–6. Available from: <http://medicinetoday.com.au/2004/january/feature-article/clinical-aspects-nicotine-dependence-and-depression>

Wilhelm K, Arnold K, Niven H, and Richmond R. Grey lungs and blue moods: Smoking cessation in the context of lifetime depression history. *Australian and New Zealand Journal of Psychiatry*, 2004; 38(11-12):896–905. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15555023>

Siahpush M, Borland R, and Scollo M. Is household smoking status associated with expenditure on food at restaurants, alcohol, gambling and insurance? Results from the 1998-99 household expenditure survey, Australia. *Tobacco Control*, 2004; 13(4):409–14. Available from: <http://tc.bmjournals.com/cgi/content/abstract/13/4/409>

Rohde P, Kahler CW, Lewinsohn PM, and Brown RA. Psychiatric disorders, familial factors, and cigarette smoking: II. Associations with progression to daily smoking. *Nicotine & Tobacco Research*, 2004; 6(1):119–32. Available from: <http://www.informaworld.com/smpp/content~db=all~content=a713938131~tab=content>

Quit South Australia, The Tobacco and Mental Illness Project, and The Queen Elizabeth Hospital and Health Service, 'I'm giving it a go'. A step-by-step guide for people with mental illness who smoke and want to cut down or quit. Adelaide, SA: Quit South Australia; 2004.

Prochaska JJ, Rossi JS, Redding CA, Rosen AB, Tsoh JY, et al. Depressed smokers and stage of change: Implications for treatment interventions. *Drug and Alcohol Dependence*, 2004; 76(2):143–51. Available from: <http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/422/CN-00504422/frame.html>

Potenza MN, Steinberg MA, McLaughlin SD, Wu R, Rounsaville BJ, et al. Characteristics of tobacco-smoking problem gamblers calling a gambling helpline. *American Journal on Addictions*, 2004; 13(5):471–93. Available from: <http://www.informaworld.com/smpp/content~content=a713665925~db=all~tab=content~order=page>

Grant BF, Hasin DS, Chou SP, Stinson FS, and Dawson DA. Nicotine dependence and psychiatric disorders in the United States: Results from the national epidemiologic survey on alcohol and related

conditions. Arch Gen Psychiatry, 2004; 61(11):1107–15. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/15520358>

Baker A, Kay-Lamkin F, Bucci S, Haile M, Richmond R, et al. Intervention for tobacco dependence among people with a mental illness, technical report no 192. Sydney: National Alcohol Research Centre, 2004. Available from: <http://catalogue.nla.gov.au/Record/3109098>.

Kisley S, Wise M, Preston N, Malmgren S, and Shannon P. A group intervention to reduce smoking in individuals with psychiatric disorder: Brief report of a pilot study. Australian & New Zealand Journal of Public Health, 2003; 27(1):61–3. Available from:

<http://www3.interscience.wiley.com/journal/118894869/abstract>

Killen JD, Fortmann SP, Schatzberg A, Hayward C, and Varady A. Onset of major depression during treatment for nicotine dependence. Addictive Behaviors, 2003; 28(3):461–70. Available from:

www.ncbi.nlm.nih.gov/pubmed/12628619

Kahler CW, Brown RA, Strong DR, Lloyd-Richardson EE, and Niaura R. History of major depressive disorder among smokers in cessation treatment: Associations with dysfunctional attitudes and coping. Addictive Behaviors, 2003; 28(6):1033–47. Available from:

www.ncbi.nlm.nih.gov/pubmed/12834649

Hulse G, Alcohol and drug problems: A case study workbook. Sydney: Oxford University Press; 2003.

Strasser K, Moeller-Saxone K, Meadows G, Hocking B, Stanton J, et al. Smoking cessation in schizophrenia: General practice guidelines. Australian Family Physician, 2002; 31(1):21–6. Available from: <http://cat.inist.fr/?aModele=afficheN&cpsidt=13510705>

Rigotti NA. Treatment of tobacco use and dependence. The New England Journal of Medicine, 2002; 346(7):506–12.

Petry NM and Oncken C. Cigarette smoking is associated with increased severity of gambling problems in treatment-seeking gamblers. Addiction, 2002; 97(6):745–53. Available from:

www.blackwell-synergy.com/doi/pdf/10.1046/j.1360-0443.2002.00163.x

Hulse G and White J, et al, The management of alcohol and drug problems. Sydney: Oxford University Press; 2002.

Dierker LC, Avenevoli S, Stolar M, and Merikangas KR. Smoking and depression: An examination of mechanisms of comorbidity. The American Journal of Psychiatry, 2002; 159(6):947–53. Available from: <http://ajp.psychiatryonline.org/cgi/content/full/159/6/947>

Burgess ES, Brown RA, Kahler CW, Niaura R, Abrams DB, et al. Patterns of change in depressive symptoms during smoking cessation: Who's at risk for relapse? Journal of Consulting and Clinical Psychology, 2002; 70(2):356–61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11952193>

Ashton M and Weston M. A smoking cessation program for people with mental illness, book of proceedings. in The Australian and New Zealand Mental Health Services Conference. 2002.

Lawrence D, Holman C, and Jablensky A. Duty to care. Preventable physical illness in people with mental illness. Perth: The University of Western Australia; 2001

Wilens TE, Spencer TJ, Biederman J, Girard K, Doyle R, et al. A controlled clinical trial of bupropion for attention deficit hyperactivity disorder in adults. *American Journal of Psychiatry*, 2001; 158(2):282–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11156812>

Meadows G, Strassser K, Moeller-Saxone K, Hocking B, Stanton J, et al. Smoking cessation in schizophrenia: The development of collaborative management guidelines. *Australasian Psychiatry*, 2001; 9(4):340–4. Available from: <http://pt.wkhealth.com/pt/re/ausp/abstract.00127911-200112000-00006.htm;jsessionid=L9yLqrhJj6FRxpyDRnJjMNv5CgJ2QtqMcjVXHkF6XmyjQL96lFXf!353761397!181195628!8091!-1>

McNeill A. Smoking and mental health - a review of the literature. London: Smokefree London Programme, 2001. Available from: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.622.6748&rep=rep1&type=pdf>.

Glassman AH, Covey LS, Stetner F, and Rivelli S. Smoking cessation and the course of major depression: A follow-up study. *Lancet*, 2001; 357(9272):1929–32. Available from: www.ncbi.nlm.nih.gov/pubmed/11425414

George T and Vessichio J. Nicotine addiction and other psychiatric disorders. *Psychiatric Times*, 2001; 18(2). Available from: <http://www.psychiatrictimes.com/display/article/10168/49511>

George T and Vessiccho J. Nicotine addiction and schizophrenia. *Psychiatric Times*, 2001; 18(2):39. Available from: <http://www.psychiatrictimes.com/display/article/10168/49511>

Degenhardt L, Hall W, and Lynskey M. Alcohol, cannabis and tobacco use among australians: A comparison of their associations with other drug use and use disorders, affective and anxiety disorders, and psychosis. *Addiction*, 2001; 96(11):1603–14. Available from: <http://www3.interscience.wiley.com/journal/120188434/abstract>

Coghlan R, Lawrence D, Holman D, and Jablensky A. Duty of care, physical illness in people with mental illness. Perth: Dept of Public Health and Dept. of Psychiatry and Behavioural Science, University of Western Australia, 2001.

Kelly C and McCreddie R. Cigarette smoking and schizophrenia. *Advances in Psychiatric Treatment* 2000; 6:327–31. Available from: <http://apt.rcpsych.org/cgi/content/full/6/5/327>

Jablensky A, McGrath J, Herrman H, Castle D, Gureje O, et al. Psychotic disorders in urban areas: An overview of the study on low prevalence disorders. *Australian & New Zealand Journal of Psychiatry*, 2000; 34(2):221–36. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10789527>

Harris J, Best D, Man L, Welch S, Gossop M, et al. Changes in cigarette smoking among alcohol and drug misusers during inpatient detoxification. *Addiction Biology*, 2000; 5(4):443–50. Available from: <http://www3.interscience.wiley.com/journal/119050468/abstract>

Frosch D, Shoptaw S, Nahom D, and Jarvik M. Associations between tobacco smoking and illicit drug use among methadone-maintained opiate-dependent individuals. *Experimental and Clinical Psychopharmacology*, 2000; 8(1):97–103. Available from: www.apa.org/journals/features/pha8197.pdf

Breslau N and Johnson EO. Predicting smoking cessation and major depression in nicotine-dependent smokers. *American Journal of Public Health*, 2000; 90(7):1122–7. Available from: <http://www.ajph.org/cgi/reprint/90/7/1122>

Ashton M, Weston M, Crocker A, and Read K. Tobacco and mental illness, what can we do and is it our business?. . Book of proceedings. in *The Australian and New Zealand Mental Health Services Conference*. 2000.

Rosen-Chase C and Dyson V. Treatment of nicotine dependence in the chronic mentally ill. *Journal of Substance Abuse Treatment*, 1999; 16(4):315–20. Available from: http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6T90-3WJ6S8K-5&_user=559483&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_version=1&_urlVersion=0&_us_erid=559483&_md5=abc105fe25106621261bd7b8d1c4e3b7

Keating D and Hertzman C, *Developmental health and the wealth of nations: Social, biological, and educational dynamics*. New York: Guilford Press; 1999. Available from: http://eric.ed.gov/ERICWebPortal/custom/portlets/recordDetails/detailmini.jsp?_nfpb=true&_ERID=ED433914&ERICExtSearch_SearchType_0=no&accno=ED433914

Jorm AF, Rodgers B, Jacomb PA, Christensen H, Henderson S, et al. Smoking and mental health: Results from a community survey. *Medical Journal of Australia*, 1999; 170(2):74–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10026688>

Jorm A. Association between smoking and mental disorders: Results from an Australian national prevalence survey. *Australian and New Zealand Journal of Public Health*, 1999; 23(3):245–8. Available from: <http://www3.interscience.wiley.com/journal/120141554/abstract>

Jablensky A, McGrath J, Herrman H, Castle D, Gureje O, et al. *People living with psychotic illness: An Australian study 1997-98*. Canberra: Mental Health Branch, Commonwealth Department of Health and Aged Care, 1999.

Hitsman B, Pingitore R, Spring B, Mahableshwarkar A, Mizes JS, et al. Antidepressant pharmacotherapy helps some cigarette smokers more than others. *Journal of Consulting and Clinical Psychology*, 1999; 67(4):547–54. Available from: <http://psycnet.apa.org/index.cfm?fa=main.landing>

Hayford K, Patten C, Rummans T, Schroeder D, Offord K, et al. Efficacy of bupropion for smoking cessation in smokers with a former history of major depression or alcoholism. *The British Journal of Psychiatry*, 1999; 174:173–8. Available from: <http://bjp.rcpsych.org/cgi/content/abstract/174/2/173>

Covey L. Tobacco cessation among patients with depression. *Primary Care*, 1999; 26(3):691. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10436294>

Wolfiger NH. The effects of parental divorce on adult tobacco and alcohol consumption. *Journal of Health and Social Behavior*, 1998; 39(3):254–69. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9785697>

Wilhelm K. The relevance of smoking and nicotine to clinical psychiatry. *Australasian Psychiatry*, 1998; 6(3):130–2.

Cunningham-Williams RM, Cottler LB, Compton WM, 3rd, and Spitznagel EL. Taking chances: Problem gamblers and mental health disorders--results from the st. Louis epidemiologic catchment area study. *American Journal of Public Health*, 1998; 88(7):1093–6.

SANE. Cessation of smoking in long term psychiatric patients. Melbourne, VIC: SANE, 1997.

Australian Bureau of Statistics. 4326.0 mental health and well-being: Profile of adults. Canberra: ABS, 1997. Available from: <http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4326.01997?OpenDocument>.

Smart RG and Ferris J. Alcohol, drugs and gambling in the ontario adult population, 1994. *The Canadian Journal of Psychiatry*, 1996; 41(1):36–45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8919422>

Polgar S, McGartland M, Borlongon C, Shytle R, and Sandber O. Smoking cessation programs are neglecting the needs of persons with neuropsychiatric disorders. *Australian and New Zealand Journal of Medicine*, 1996; 26(4):572–3. Available from: <http://cat.inist.fr/?aModele=afficheN&cpsidt=3198056>

Hurt RD, Offord KP, Croghan IT, Gomez-Dahl L, Kottke TE, et al. Mortality following inpatient addictions treatment. Role of tobacco use in a community-based cohort. *Journal of the American Medical Association*, 1996; 275(14):1097–103. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/8601929>

Bertram S, B F, and S E. Knowledge and use of alcohol and other drugs among vietnamese-speaking migrants. *Drug and Alcohol Review*, 1996; 15(2):121–6. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/09595239600185761>

Breslau N. Psychiatric comorbidity of smoking and nicotine dependence. *Behavior Genetics*, 1995; 25(2):95–101. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7733862>

Bertram S and Flaherty B. Alcohol and other drug use, attitudes and knowledge amongst spanish-speakers in sydney and wollongong.: DAMEC, NSW, 1993.

Lohr J and Flynn K. Smoking and schizophrenia. *Schizophrenia Research*, 1992; 8(2):93–102. Available from: www.blackwell-synergy.com/doi/pdf/10.1046/j.1365-2850.1997.00056.x

Difranza J and Guerrea M. Alcoholism and smoking. *Journal of Studies on Alcohol*, 1990; 51:130–5.

Quit Victoria. Mental health settings. Available from: <https://www.quit.org.au/resources/mental-health/>

Australian Bureau of Statistics. 4364.0.55.001—national health survey: First results, 2014–15
Available from:

<http://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/4364.0.55.001~2014-15~Main%20Features~Key%20findings~1>

Australian Bureau of Statistics. TableBuilder. Available from:

<http://www.abs.gov.au/websitedbs/d3310114.nsf/home/about+tablebuilder>

9A.3.1 Prevalence of smoking among people with mental health problems

Janssen, E, Vuolo, M, & Airagnes, G. (2024). Daily cigarette smoking among inpatients for substance use disorders in France, 2010-2020: Commonalities and specificities across substances. *Tob Induc Dis*, 22. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39502623>

Datta, BK, Coughlin, SS, Ayadi, J, Majeed, B, & Ansa, BE. (2024). Relationship between social support, life satisfaction, and smoking status among US adults with cardiovascular risks. *Behav Med*, 50(4), 288-297. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39470042>

Airagnes, G, Sanchez-Rico, M, Deguilhem, A, Blanco, C, Olfson, M, Ouazana Vedrines, C et al. (2024). Nicotine dependence and incident psychiatric disorders: prospective evidence from US national study. *Mol Psychiatry*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39261672>

Gaiha, SM, Wang, M, Baiocchi, M, & Halpern-Felsher, B. (2024). Depression screening outcomes among adolescents, young adults, and adults reporting past 30-day tobacco and cannabis use. *Addict Behav*, 156, 108076. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38838604>

Thaker, P, Gartner, C, Kisely, S, & Plevier, S. (2024). Systematic review of tobacco smoking prevalence among young people in treatment for first-episode psychosis. *Int J Ment Health Nurs*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38606655>

Cho, S, Park, JH, Kim, DH, Choi, H, Park, Y, Kim, HJ et al. (2024). Association between Combustible Cigarettes and Noncombustible Nicotine or Tobacco Products and Generalized Anxiety Disorder Based on Data from the 8th Korea National Health and Nutrition Examination Survey 2021. *Korean J Fam Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38523421>

Ferron, JC, Brunette, MF, Aschbrenner, KA, ElSayed, MW, & Pratt, SI. (2024). Tobacco, Alcohol, and Drug Use Among Young Adults with Serious Mental Illness. *Community Ment Health J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38427276>

Raut, A., Pearlman, D. N., Doherty, J., Skelly, K., & Hartzell, H. (2024). Disparities in Tobacco Use Among Individuals With Mental Distress. *R I Med J (2013)*, 107(4), 45-48. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38536141>

Lee, PH, Tervo-Clemmens, B, Liu, RT, Gersten, MB, Jung, JY, Janes, AC, & Gilman, J. (2024). Use of Tobacco Products and Suicide Attempts Among Elementary School-Aged Children. *JAMA Netw Open*, 7(2), e240376. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38407905>

Kjeld, SG, Homberg, T, & Andersen, S. (2024). Relationship between smoking and mental health in educational settings with high smoking rates: a cohort study among Danish youth. *Scand J Public Health*, 14034948241227305. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38342989>

- Albarrak, DA, Alotaibi, AB, Alotaibi, RF, Alramadhan, SH, Bin Muhanna, AI, Aldehan, AM, & Bin Abdulrahman, KA. (2023). The Association Between Nicotine Dependence and Mental Health in the General Population of Saudi Arabia: A Cross-Sectional Analytical Study. *Int J Gen Med*, 16, 5801-5815. . Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38106979>
- Cabrera-Serrano, A, Felici-Giovanini, ME, Diaz-Garcia, RM, Ramos-Colon, MV, & Rodriguez-Ayuso, I. (2023). Tobacco Use and Depression: A Hidden Epidemic among Smokers 18 Years and Older, Puerto Rico, 2018-2020. *P R Health Sci J*, 42(4), 298-303. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38104286>
- Schulz, JA, Gimm, G, West, JC, Kock, L, & Villanti, AC. (2023). Patterns of Alcohol, Marijuana, and Tobacco Use Among U.S. Adolescents and Young Adults by Disability Status: 2015-2019. *J Adolesc Health*. . Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38043040>
- Rajan, S, Mitchell, A, Zavala, GA, Podmore, D, Khali, H, Chowdhury, AH et al. (2023). Tobacco use in people with severe mental illness: Findings from a multi-country survey of mental health institutions in South Asia. *Tob Induc Dis*, 21, 166. . Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38098747>
- Mattingly, DT, Mezuk, B, Elliott, MR, Neighbors, HW, & Fleischer, NL. (2023). Racial and Ethnic Disparities in Mental Health Problems and Tobacco and Cannabis Use Among US Emerging Adults. *J Racial Ethn Health Disparities*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37828404>
- Parker, MA, Cruz-Cano, R, Streck, JM, Ballis, E, & Weinberger, AH. (2023). Incidence of opioid misuse by cigarette smoking status in the United States. *Addict Behav*, 147, 107837. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37659270>
- Streck, JM, Potter, K, Pachas, G, Cather, C, Nielsen, L, & Evins, AE. (2023). Alternative combusted tobacco product and multiple tobacco product use among individuals with serious mental illness enrolled in a large pragmatic randomized controlled trial. *Addict Behav*, 148, 107844. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37678006>
- Cuomo, RE, Mackey, TK, & Purushothaman, V. (2023). Tobacco/nicotine dependence as a risk factor for substance use disorders and related mental health conditions among cancer patients. *Psychooncology*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37409875>
- Hwang, JW, Eun, Y, & Song, CH. (2023). Associations of Alcohol Consumption and Smoking Behaviors with Depressed Mood According to Gender in Korean Young Adults. *Korean J Fam Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37423254>
- Watson, ML, McKnight, ER, Groner, JA, Manos, BE, Ebersole, AM, & Bonny, AE. (2023). Patterns and perceptions of nicotine use among U.S. adolescents and young adults receiving medication treatment for opioid use disorder. *Prev Med Rep*, 35, 102303. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37449006>
- De Micheli, A, Provenzani, U, Solmi, M, Van Pabst, AVL, Youssef, E, McGuire, P, & Fusar-Poli, P. (2023). Prevalence of tobacco smoking in people at clinical high-risk for psychosis: Systematic review

and meta-analysis. *Schizophr Res*, 254, 109-115. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36827942>

Loretan, CG, Wang, TW, Watson, CV, & Jamal, A. (2022). Disparities in Current Cigarette Smoking Among US Adults With Mental Health Conditions. *Prev Chronic Dis*, 19, E87. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36548524>

Akbari, M, Seydavi, M, Chasson, GS, Leventhal, AM, & Lockwood, MI. (2022). Global Prevalence of Smoking Among Individuals with Obsessive-Compulsive Disorder and Symptoms: A Systematic Review and Meta-Analysis. *Health Psychol Rev*, 1-36. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36173036>

Galiatsatos, P, Oluyinka, M, Min, J, Schreiber, R, Lansey, DG, Ikpe, R et al. (2022). Prevalence of Mental Health and Social Connection among Patients Seeking Tobacco Dependence Management: A Pilot Study. *Int J Environ Res Public Health*, 19(18). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36142029>

Goodwin, RD, & Davidson, L. (2022). Prevalence of Cigarette Smoking Among US Adults With Major Depression or Substance Use Disorders. *JAMA*, 328(6), 585. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35943476>

Han, B, Tipperman, D, & Compton, WM. (2022). Prevalence of Cigarette Smoking Among US Adults With Major Depression or Substance Use Disorders-Reply. *JAMA*, 328(6), 585-586. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35943474>

Asharani, PV, Seet, V, Devi, F, Wang, P, Roystonn, K, & Subramaniam, M. (2022). Electronic nicotine delivery systems: prevalence and perception of risk/harm in individuals with mental illness. *Singapore Med J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35851648>

Cotton, SM, Sharmin, S, Gao, CX, Brown, E, Menssink, J M, Rickwood, D et al. (2022). Prevalence and Correlates of Tobacco Use in Young People Presenting to Australian Primary Mental Health Services. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35665823>

Lee B, Levy DE, Macy JT, Elam KK, Bidulescu A, et al. Smoking trajectories from adolescence to early adulthood as a longitudinal predictor of mental health in adulthood: Evidence from 21 years of a nationally representative cohort. *Addiction*, 2022; 117(6):1727–36. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34817100>

Kastaun S, Brose LS, Scholz E, Viechtbauer W, and Kotz D. Mental health symptoms and associations with tobacco smoking, dependence, motivation, and attempts to quit: Findings from a population survey in germany (debra study). *Eur Addict Res*, 2022:1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35358964>

Iatrou G, Gourgoulisanis KI, Kotrotsiou E, and Gouva M. Assessment of greek smokers' psychological characteristics and empathy while smoking in enclosed public spaces and near nonsmokers. *Cureus*, 2022; 14(3):e22910. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35399413>

Han B, Volkow ND, Blanco C, Tipperman D, Einstein EB, et al. Trends in prevalence of cigarette smoking among US adults with major depression or substance use disorders, 2006-2019. *Journal of*

the American Medical Association, 2022; 327(16):1566–76. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/35471512>

Greenhalgh EM, Brennan E, Segan C, and Scollo M. Monitoring changes in smoking and quitting behaviours among australians with and without mental illness over 15 years. *Australian and New Zealand Journal of Public Health*, 2022; 46(2):223–9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34821438>

Wei Y, Lv Y, Zhou J, Gao X, Duan J, et al. Smoking cessation in late life is associated with increased risk of all-cause mortality amongst oldest old people: A community-based prospective cohort study. *Age Ageing*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33492360>

Tomioka K, Shima M, and Saeki K. Association between heaviness of cigarette smoking and serious psychological distress is stronger in women than in men: A nationally representative cross-sectional survey in Japan. *Harm Reduct J*, 2021; 18(1):27. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33663543>

Sanchez-Villegas A, Gea A, Lahortiga-Ramos F, Martinez-Gonzalez J, Molero P, et al. Bidirectional association between tobacco use and depression risk in the sun cohort study. *Adicciones*, 2021; 0(0):1725. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34882246>

Public Health England. Smoking prevalence in adults: Current smokers aged over 18 years by wellbeing group and region. PHE, 2021. Available from:

<https://www.gov.uk/government/publications/smoking-prevalence-in-adults-current-smokers-aged-over-18-years-by-wellbeing-group-and-region/smoking-prevalence-in-adults-commentary-current-smokers-aged-over-18-years-by-wellbeing-group-and-region>.

Peckham E, Allgar V, Crosland S, Heron P, Johnston G, et al. Investigating smoking and nicotine dependence among people with severe mental illness during the COVID-19 pandemic: Analysis of linked data from a UK closing the gap cohort. *BJPsych Open*, 2021; 7(3):e86. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33888178>

Lopes J and Silva I. Prevalence of tobacco smoking in patients with schizophrenia and bipolar disorder in a portuguese hospital. *Acta Med Port*, 2021; 34(4):321-2. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34214427>

Lloyd EC, Reed ZE, and Wootton RE. The absence of association between anorexia nervosa and smoking: Converging evidence across two studies. *Eur Child Adolesc Psychiatry*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34939143>

Lee J, Salloum RG, Lindstrom K, and Kathryn McHugh R. Benzodiazepine misuse and cigarette smoking status in US adults: Results from the national survey on drug use and health, 2015-2018. *Addictive Behaviors*, 2021; 123:107058. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34315094>

Khan MR, Ban K, Caniglia EC, Edelman JE, Gaither J, et al. Brief original report: Does smoking status provide information relevant to screening for other substance use among US adults? *Preventive*

Medicine Reports, 2021; 23:101483. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34345578>

Huh Y and Cho HJ. Associations between the type of tobacco products and suicidal behaviors: A nationwide population-based study among Korean adolescents. *International Journal of Environmental Research and Public Health*, 2021; 18(2). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33418893>

Hahad O, Daiber A, Michal M, Kuntic M, Lieb K, et al. Smoking and neuropsychiatric disease-associations and underlying mechanisms. *Int J Mol Sci*, 2021; 22(14). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34298890>

Guggenheim FG, Lieberman PB, and Farris SG. Cigarette smoking in an acute partial hospital program. *J Nerv Ment Dis*, 2021; 209(6):415-20. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33966016>

Greenhalgh E and Scollo M. Alcohol and tobacco use in victoria and Australia: Results from the 2019 National Drug Strategy Household Survey. Melbourne: Centre for Behavioural Research in Cancer, Cancer Council Victoria, 2021.

Gold AK, Albury EA, Stromberg A, French NJ, Peters AT, et al. Cigarette smokers with a mood symptom history enrolled in a physical activity trial. *Bipolar Disorders*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34606143>

Friedman AS, Tam J, and Busch SH. US smoking trends among young adults with serious psychological distress: 1997-2018. *Journal of General Internal Medicine*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34378116>

Fornaro M, Carvalho AF, De Prisco M, Mondin AM, Billeci M, et al. The prevalence, odds, predictors, and management of tobacco use disorder or nicotine dependence among people with severe mental illness: Systematic review and meta-analysis. *Neurosci Biobehav Rev*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34838527>

Fan Z, Gong X, Xu H, Wang H, Zeng N, et al. Gender differences in the associations between tobacco smoke exposure and depressive symptoms among u.S. Adults: Nhanes 2007-2018. *Journal of Psychiatric Research*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34799126>

Erinoso O, Oyapero A, Osoba M, Amure M, Osibogun O, et al. Association between anxiety, alcohol, poly-tobacco use and waterpipe smoking: A cross-sectional study in lagos, Nigeria. *Niger Postgrad Med J*, 2021; 28(2):117-25. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34494598>

Chido-Amajuoyi OG, Mantey DS, Omega-Njemnobi O, Yu RK, and Shete S. Association of dual and poly tobacco use with depressive symptoms and use of antidepressants. *Addictive Behaviors*, 2021; 115:106790. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33387979>

Alzahrani HA, Alghamdi MS, Alzahrani AA, and Alzahrani AA. Tobacco smoking and depressive symptoms among male medical students in al-baha university. *J Family Med Prim Care*, 2021; 10(6):2235-40. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34322418>

Ali AM, Hori H, Kim Y, and Kunugi H. Predictors of nutritional status, depression, internet addiction, facebook addiction, and tobacco smoking among women with eating disorders in Spain. *Front Psychiatry*, 2021; 12:735109. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34899416>

Tzu-Hsuan Chen D. The psychosocial impact of the COVID-19 pandemic on changes in smoking behavior: Evidence from a nationwide survey in the UK. *Tob Prev Cessat*, 2020; 6:59. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33163705>

Tam J, Taylor GMJ, Zivin K, Warner KE, and Meza R. Modeling smoking-attributable mortality among adults with major depression in the United States. *Preventive Medicine*, 2020; 140:106241. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32860820>

Smith PH, Chhipa M, Bystrick J, Roy J, Goodwin RD, et al. Cigarette smoking among those with mental disorders in the US population: 2012-2013 update. *Tobacco Control*, 2020; 29(1):29-35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30377242>

Parker MA and Villanti AC. Relationship between comorbid drug use disorders, affective disorders, and current smoking. *Substance Use and Misuse*, 2020;1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33143491>

Guerin N and White V. *Assad 2017 statistics & trends: Australian secondary students' use of tobacco, alcohol, over-the-counter drugs, and illicit substances in 2017. Second edition.* Cancer Council Victoria, 2020. Available from: <https://www.health.gov.au/resources/publications/secondary-school-students-use-of-tobacco-alcohol-and-other-drugs-in-2017>.

Brose LS, Brown J, Robson D, and McNeill A. Mental health, smoking, harm reduction and quit attempts - a population survey in England. *BMC Public Health*, 2020; 20(1):1237. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32795286>

Australian Institute of Health and Welfare. *National Drug Strategy Household Survey 2019. Drug Statistics series no. 32 Cat. no. PHE 270* Canberra: AIHW, 2020. Available from: <https://www.aihw.gov.au/getmedia/3564474e-f7ad-461c-b918-7f8de03d1294/aihw-phe-270-NDSHS-2019.pdf.aspx?inline=true>.

Australian Institute of Health and Welfare. *Data tables: National Drug Strategy Household Survey 2019 - 2. Tobacco smoking chapter, supplementary data tables.* Canberra: AIHW, 2020. Available from: <https://www.aihw.gov.au/reports/illicit-use-of-drugs/national-drug-strategy-household-survey-2019/data>.

Sharma P, Virani S, Mitra S, Ahmed S, and Desousa A. Trends of tobacco consumption among women admitted to and visiting the psychiatric inpatient unit of a tertiary care center in Mumbai: A cross-sectional study. *Prim Care Companion CNS Disord*, 2019; 21(3). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31087823>

Richardson S, McNeill A, and Brose LS. Smoking and quitting behaviours by mental health conditions in Great Britain (1993-2014). *Addictive Behaviors*, 2019; 90:14–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30352340>

Parker MA, Sigmon SC, and Villanti AC. Higher smoking prevalence among United States adults with co-occurring affective and drug use diagnoses. *Addictive Behaviors*, 2019; 99:106112. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31476691>

Oliveira RM, Santos JLF, and Furegato ARF. Prevalence and smokers' profile: Comparisons between the psychiatric population and the general population. *Rev Lat Am Enfermagem*, 2019; 27:e3149. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31038640>

Murray EG, Isbister GK, McCrabb S, Halpin SA, and Bonevski B. An examination of factors associated with tobacco smoking amongst patients presenting with deliberate self-poisoning. *J Affect Disord*, 2019; 260:544-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31539691>

Ferreira VR, Jardim TV, Sousa ALL, Rosa BMC, and Jardim PCV. Smoking, alcohol consumption and mental health: Data from the brazilian study of cardiovascular risks in adolescents (erica). *Addictive Behaviors Reports*, 2019; 9:100147. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31193769>

Al-Mrayat YD, Okoli CTC, Studts CR, Rayens MK, and Hahn EJ. The psychometric properties of the minnesota tobacco withdrawal scale among patients with mental illness. *Biol Res Nurs*, 2019;1099800419895573. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31854206>

Abd Rashid R, Kanagasundram S, Danaee M, Abdul Majid H, Sulaiman AH, et al. The prevalence of smoking, determinants and chance of psychological problems among smokers in an urban community housing project in malaysia. *International Journal of Environmental Research and Public Health*, 2019; 16(10). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31109033>

Petty D and Aubry T. Tobacco use and smoking behaviors of individuals with a serious mental illness. *Psychiatric Rehabilitation Journal*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29781639>

Lores L, Monje A, Bergada M, Arellano E, Rodriguez-Larrea J, et al. Prevalence of smoking in a psychiatric hospital and its relationship with respiratory symptoms and the prevalence of copd. *Int J Chron Obstruct Pulmon Dis*, 2018; 13:2797-804. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30233170>

Fujiwara M, Inagaki M, Nakaya N, Fujimori M, Higuchi Y, et al. Smoking among adults with serious psychological distress: Analysis of anonymized data from a national cross-sectional survey in Japan. *J Affect Disord*, 2018; 239:131-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30005326>

Dickerson F, Schroeder J, Katsafanas E, Khushalani S, Origoni AE, et al. Cigarette smoking by patients with serious mental illness, 1999-2016: An increasing disparity. *Psychiatric Services*, 2018; 69(2):147-53. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28945183>

Al Risi K, Al Maqbali M, Asiri WA, and Al Sinawi H. Rate of tobacco smoking among psychiatric outpatients attending a tertiary care hospital in Oman: A cross-sectional study. *Oman Med J*, 2018; 33(3):224-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29896330>

Quickstats: Percentage* of adults aged ≥ 18 years with or without psychological distress(dagger) who were current smokers,(section sign) by age group and level of distress - national health

interview survey,(paragraph sign) 2014-2016. Morbidity and Mortality Weekly Report, 2018; 67(23):672. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29902163>

Matcham F, Carroll A, Chung N, Crawford V, Galloway J, et al. Smoking and common mental disorders in patients with chronic conditions: An analysis of data collected via a web-based screening system. *General Hospital Psychiatry*, 2017; 45:12-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28274333>

Luiz Ribeiro A, Satoshi N, and Catherine MO. Heartbeat: Coronary heart disease, obesity, smoking and long-lasting psychological distress. *Heart*, 2017; 103(9):644. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28396464>

Burns A, Strawbridge JD, Clancy L, and Doyle F. Exploring smoking, mental health and smoking-related disease in a nationally representative sample of older adults in Ireland - a retrospective secondary analysis. *Journal of Psychosomatic Research*, 2017; 98:78-86. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28554376>

Schapir L, Lahav T, Zalsman G, Krivoy A, Sever J, et al. Cigarette smoking, alcohol and cannabis use in patients with pervasive developmental disorders. *Substance Use and Misuse*, 2016:1–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27248185>

McElroy SL. The dual epidemic of tobacco dependence and obesity among those with severe mental illness. *Journal of Clinical Psychiatry*, 2016; 77(3):e327–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27046321>

Lawrence D and Williams JM. Trends in smoking rates by level of psychological distress-time series analysis of US national health interview survey data 1997-2014. *Nicotine & Tobacco Research*, 2016; 18(6):1463–70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26705303>

Guydish J, Passalacqua E, Pagano A, Martinez C, Le T, et al. An international systematic review of smoking prevalence in addiction treatment. *Addiction*, 2016; 111(2):220–30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26392127>

Flanagan JC, Hakes JK, McClure EA, Snead AL, and Back SE. Effects of intimate partner violence, PTSD, and alcohol use on cigarette smoking in a nationally representative sample. *American Journal on Addictions*, 2016; 25(4):283–90. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27196699>

Steinberg ML, Williams JM, and Li Y. Poor mental health and reduced decline in smoking prevalence. *American Journal of Preventive Medicine*, 2015; 49(3):362–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26071864>

Kotecha S. Mental health patients 'smoke three times as much'. *BBC News*, 2015. Available from: <http://www.bbc.com/news/health-32982669>

Jackson JG, Diaz FJ, Lopez L, and de Leon J. A combined analysis of worldwide studies demonstrates an association between bipolar disorder and tobacco smoking behaviors in adults. *Bipolar Disorders*, 2015; 17(6):575–97. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26238269>

Forbes MK, Flanagan JC, Barrett EL, Crome E, Baillie AJ, et al. Smoking, posttraumatic stress disorder, and alcohol use disorders in a nationally representative sample of Australian men and women. *Drug and Alcohol Dependence*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26386825>

Dell'Osso B, Nicolini H, Lanzagorta N, Benatti B, Spagnolin G, et al. Cigarette smoking in patients with obsessive compulsive disorder: A report from the international college of obsessive compulsive spectrum disorders (icocs). *CNS Spectr*, 2015; 20(5):469–73. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26349811>

Chao AM, White MA, and Grilo CM. Smoking status and psychosocial factors in binge eating disorder and bulimia nervosa. *Eat Behav*, 2015; 21:54–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26741260>

Cao XL, Li Y, Zhong BL, Chiu HF, Ungvari GS, et al. Current cigarette smoking in chinese female patients with schizophrenia: A meta-analysis. *Psychiatry Res*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26688158>

Wijesundera H, Hanwella R, and de Silva VA. Antipsychotic medication and tobacco use among outpatients with schizophrenia: A cross-sectional study. *Annals of General Psychiatry*, 2014; 13(1):7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24642279>

Lipari RN and Van Horn S. Smoking and mental illness among adults in the United States, in The cbhsq report. Rockville (MD): 2013. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28459516>.

Leung J, Gartner C, Hall W, Lucke J, and Dobson A. A longitudinal study of the bi-directional relationship between tobacco smoking and psychological distress in a community sample of young Australian women. *Psychological Medicine*, 2011:1–10. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22011372>

Clarke D, Eaton W, Petronis K, Ko J, Chatterjee A, et al. Increased risk of suicidal ideation in smokers and former smokers compared to never smokers: Evidence from the baltimore eca follow-up study. *Suicide & Life-Threatening Behavior*, 2010; 40(4):307–18. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2936702/?tool=pubmed>

Ischaki E and Gratziau C. Smoking and depression: Is smoking cessation effective? *Therapeutic Advances in Respiratory Disease*, 2009; 3(1):31–8. Available from: <http://tar.sagepub.com/cgi/reprint/3/1/31>

Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, et al. Smoking and mental illness: A population-based prevalence study. *Journal of the American Medical Association*, 2000; 284(20):2606–10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11086367>

9A.3.1.2 Smoking prevalence over time

Streck, JM, Parker, MA, Cruz, R, Rosen, RL, Baker, TB, Piper, ME, & Weinberger, AH. (2024). Prevalence and Trends in Cigarette Smoking With and Without Tobacco Use Disorder Among Adults in the United States: 2010–2021. *J Clin Psychiatry*, 85(3). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38874573>

Kiviruusu, O, Berg, N, Piirtola, M, Viertio, S, Suvisaari, J, Korhonen, T, & Marttunen, M. (2024). Life course associations between smoking and depressive symptoms. A 30-year Finnish follow-up study. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38243907>

Branas, M, Frankenburg, FR, Temes, CM, Fitzmaurice, GM, & Zanarini, MC. (2023). Longitudinal Description and Prediction of Smoking Among Borderline Patients: An 18-Year Follow-Up Study. *J Clin Psychiatry*, 84(6). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37943989>

Saloner, B, Li, W, Flores, M, Progovac, AM, & Le Cook, B. (2023). A Widening Divide: Cigarette Smoking Trends Among People With Substance Use Disorder And Criminal Legal Involvement. *Health Aff (Millwood)*, 42(2), 187-196. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36745833>

Wu MH, Wang WE, Wang TN, Lin NC, Lu ML, et al. Asthma and early smoking associated with high risk of panic disorder in adolescents and young adults. *Soc Psychiatry Psychiatr Epidemiol*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34279695>

Wang Y, Liu Y, Waldron M, Houston-Ludlam AN, McCutcheon VV, et al. Temporal trends in smoking and nicotine dependence in relation to co-occurring substance use in the United States, 2005-2016. *Drug and Alcohol Dependence*, 2021; 226:108903. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34304125>

Stamates AL, Roberts R, and Lau-Barraco C. Alcohol, cannabis, and tobacco polysubstance use: A latent profile analysis of age of onset. *Substance Abuse*, 2021;1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34283703>

Schermitzler B, Miley K, Vinogradov S, and Ramsay IS. Smoking is related to reduced motivation, but not global cognition, in the first two years of treatment for first episode psychosis. *J Clin Med*, 2021; 10(8). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33920376>

Safarini OA, Taya H, Abu Elhija Y, Qadous M, Farhoud A, et al. Assessment of the relationship of depression with tobacco and caffeine use among university students: A cross-sectional study. *Cureus*, 2021; 13(10):e19098. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34868751>

Quittschalle J, Pabst A, Lobner M, Lupp M, Hesel K, et al. Association of alcohol and tobacco consumption with depression severity in the oldest old. Results from the age different old age cohort platform. *International Journal of Environmental Research and Public Health*, 2021; 18(15). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34360253>

Higuchi Y, Fujiwara M, Nakaya N, Fujimori M, Yamada Y, et al. Trends in smoking rates among individuals with serious psychological distress: Analysis of data from a Japanese national survey, 2007-2016. *Psychiatry Res*, 2020; 291:113225. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32599444>

Zvolensky MJ, Jardin C, Wall MM, Gbedemah M, Hasin D, et al. Psychological distress among smokers in the United States: 2008-2014. *Nicotine & Tobacco Research*, 2018; 20(6):707-13. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28482108>

Weinberger AH, Streck JM, Pacek LR, and Goodwin RD. Nondaily cigarette smoking is increasing among people with common mental health and substance use problems in the United States: Data

from representative samples of US adults, 2005-2014. *Journal of Clinical Psychiatry*, 2018; 79(5). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30153404>

Streck JM, Weinberger AH, Pacek LR, Gbedemah M, and Goodwin RD. Cigarette smoking quit rates among persons with serious psychological distress in the United States from 2008-2016: Are mental health disparities in cigarette use increasing? *Nicotine & Tobacco Research*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30351429>

Mishina K, Tiiri E, Lempinen L, Sillanmaki L, Kronstrom K, et al. Time trends of finnish adolescents' mental health and use of alcohol and cigarettes from 1998 to 2014. *Eur Child Adolesc Psychiatry*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29704065>

Zvolensky MJ, Jardin C, Wall MM, Gbedemah M, Hasin D, et al. Psychological distress among smokers in the United States: 2008 to 2014. *Nicotine & Tobacco Research*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28482108>

Hartz SM, Horton AC, Hancock DB, Baker TB, Caporaso NE, et al. Genetic correlation between smoking behaviors and schizophrenia. *Schizophr Res*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28285025>

Young-Wolff KC, Kline-Simon AH, Das S, Mordecai DJ, Miller-Rosales C, et al. Smoking trends among adults with behavioral health conditions in integrated health care: A retrospective cohort study. *Psychiatric Services*, 2016; appips201500337. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27079992>

Larson A, Bovbjerg V, and Luck J. State clean indoor air laws and smoking among adults with poor mental health. *J Public Health Policy*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27146608>

Forman-Hoffman VL, Hedden SL, Miller GK, Brown K, Teich J, et al. Trends in cigarette use, by serious psychological distress status in the United States, 1998-2013. *Addictive Behaviors*, 2016; 64:223-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27690139>

Jasek JP, Williams JM, Mandel-Ricci J, and Johns M. Trends in smoking among adults with serious psychological distress during comprehensive tobacco control in new york city, 2003–2012. *Tobacco Control*, 2015; 24(6):622–3. Available from: <http://tobaccocontrol.bmj.com/content/24/6/622.short>

9A.3.1.1 Latest estimates of prevalence

Hanafin, J, Sunday, S, & Clancy, L. (2024). Inequalities in Smoking and E-Cigarette Use in Young Adults With Mental Ill-Health, 20 years After Ireland's Smoking Ban. *Tob Use Insights*, 17, 1179173X241283744. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39290597>

9A.3.2 Contribution of smoking to health outcomes and social inequality

Pietschnig, J, Oberleiter, S, & Kohler, MD. (2024). Smoking behavior is associated with suicidality in individuals with psychosis and bipolar disorder: a systematic quantitative review and meta-analysis. *Front Psychol*, 15, 1369669. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39328818>

- Liu, Y, Zhang, L, Fu, S, Wei, S, Jin, Z, & He, L. (2024). Gender differences in the relationship between nicotine exposure and symptoms of depression. *Pharmacol Biochem Behav*, 244, 173857. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39154790>
- Qiu, L, Liang, C, Kochunov, P, Hutchison, KE, Sui, J, Jiang, R et al. (2024). Associations of alcohol and tobacco use with psychotic, depressive and developmental disorders revealed via multimodal neuroimaging. *Transl Psychiatry*, 14(1), 326. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39112461>
- Ren, J, Zhang, Y, Song, H, Gou, H, Zhao, Q, Hong, W et al. (2024). The interaction of oxytocin and nicotine addiction on psychosocial stress: an fMRI study. *Transl Psychiatry*, 14(1), 348. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39214996>
- Zhu, X, Ye, R, Jiang, X, & Zhang, J. (2024). Smoking as a mediator in the association between major depressive disorder and schizophrenia on lung cancer risk: a bidirectional/multivariable and mediation Mendelian randomization study. *Front Psychiatry*, 15, 1367858. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39176232>
- Wilkinson, AV, Chen, B, Swann, AC, Graham, DP, Nielsen, DA, Kosten, TR et al. (2024). Use of cigarettes and e-cigarettes, impulsivity and anxiety: Influences on suicidal ideation among youth and young adults in Texas. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39028574>
- Ghaderi, A, Khoshakhlagh, AH, Gruszecka-Kosowska, A, Askari-Lemjiri, F, Alemi, F, Molavi, N et al. (2024). Heavy metal concentrations and clinical symptoms in patients diagnosed with schizophrenia related to cigarette smoking. *Sci Rep*, 14(1), 15074. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38956098>
- Liu, LJ, Lin, EM, Tsao, SL, Wang, HY, & Ho, MC. (2024). Attentional Bias for Opioids in Taiwanese Heavy Smokers with Chronic Noncancer Pain. *Medicina (Kaunas)*, 60(7). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39064536>
- Velioglu, HA, Yildiz, S, Ozdemir-Oktem, E, Cankaya, S, Lundmark, AK, Ozsimsek, A et al. (2024). Smoking affects global and regional brain entropy in depression patients regardless of depression: Preliminary findings. *J Psychiatr Res*, 177, 147-152. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39018709>
- Gaiha, SM, Wang, M, Baiocchi, M, & Halpern-Felsher, B. (2024). Depression screening outcomes among adolescents, young adults, and adults reporting past 30-day tobacco and cannabis use. *Addict Behav*, 156, 108076. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38838604>
- Johnson, EC, Austin-Zimmerman, I, Thorpe, HHA, Levey, DF, Baranger, DAA, Colbert, SMC et al. (2024). Cross-ancestry genetic investigation of schizophrenia, cannabis use disorder, and tobacco smoking. *Neuropsychopharmacology*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38906991>
- Koster, M, Mannsdorfer, L, van der Pluijm, M, de Haan, L, Ziermans, T, van Wingen, G, & Vermeulen, J. (2024). The Association Between Chronic Tobacco Smoking and Brain Alterations in Schizophrenia: A Systematic Review of Magnetic Resonance Imaging Studies. *Schizophr Bull*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38824451>

Urun Unal, B, & Marakoglu, K. (2024). Smoking and Depression among Medical School Students: A Cross-Sectional Study from Turkey's Largest Province. *Healthcare (Basel)*, 12(11). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38891206>

Azeem, T, Khan, S, & Samiullah, F. (2024). Effects of smoking behaviour changes on depression in older people: A retrospective study. *Australas J Ageing*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38742245>

Park, SK, Oh, CM, Kim, E, Ryoo, JH, & Jung, JY. (2024). The longitudinal analysis for the association between smoking and the risk of depressive symptoms. *BMC Psychiatry*, 24(1), 364. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38750466>

Qu, Y, Du, Z, Shen, Y, Zhou, Q, Zhou, Z, Jiang, Y, & Zhu, H. (2024). Smoking may increase the usage of antidepressant: evidence from genomic perspective analysis. *Eur Arch Psychiatry Clin Neurosci*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38702554>

Rantanen, AT, Kautiainen, H, Ekblad, MO, & Korhonen, PE. (2024). Depressive symptoms and smoking: Effect on mortality in a primary care cohort. *J Psychosom Res*, 182, 111690. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38704926>

Kwon, M, Nam, E, & Lee, J. (2024). Poly-tobacco use and mental health in South Korean adolescents. *Tob Induc Dis*, 22. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38779296>

Chen, Y, Chaudhary, S, Li, G, Fucito, LM, Bi, J, & Li, CR. (2024). Deficient sleep, altered hypothalamic functional connectivity, depression and anxiety in cigarette smokers. *Neuroimage Rep*, 4(1). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38605733>

Ma, Z, & Ma, Z. (2024). Relationship between stable smoking behavior and cognitive function in male schizophrenia patients. *J Psychiatr Res*, 173, 296-301. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38555677>

Kondracki, AJ, Attia, JR, Valente, MJ, Roth, KB, Akin, M, McCarthy, CA, & Barkin, JL. (2024). Exploring a Potential Interaction Between the Effect of Specific Maternal Smoking Patterns and Comorbid Antenatal Depression in Causing Postpartum Depression. *Neuropsychiatr Dis Treat*, 20, 795-807. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38586309>

Li, W, Yue, L, & Xiao, S. (2024). Demographic, biochemical, clinical, and cognitive symptom differences between smokers and non-smokers in Chinese older male patients with chronic schizophrenia. *Eur Arch Psychiatry Clin Neurosci*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38462585>

Ma, B, Zhao, Q, Song, J, Li, Y, Bian, Y, Wang, Z, & Yang, F. (2024). Smoking Behavior and its Effect on Psychiatric Symptoms of Patients with Stable Schizophrenia. *Altern Ther Health Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38430154>

Mohan, P, Chatterjee, K, Sinha, S, Saini, RK, Sharma, AK, & Waikole, S. (2024). Smoking blunts sertraline response in depression: A prospective observational cohort study. *Med J Armed Forces India*, 80(2), 145-152. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38525466>

- Mu, X, Wu, W, Wang, S, Su, X, Guan, H, Guan, X et al. (2024). Smoking affects symptom improvement in schizophrenia: a prospective longitudinal study of male patients with first-episode schizophrenia. *Schizophrenia (Heidelb)*, 10(1), 34. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38491003>
- Lee, PH, Tervo-Clemmens, B, Liu, RT, Gersten, MB, Jung, JY, Janes, AC, & Gilman, J. (2024). Use of Tobacco Products and Suicide Attempts Among Elementary School-Aged Children. *JAMA Netw Open*, 7(2), e240376. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38407905>
- Blaes, SL, Lewis, B, Teitelbaum, S, Reisfeld, G, & Boissoneault, J. (2024). Associations between smoking and pain in early recovery in residential substance use treatment-seekers. *J Subst Use Addict Treat*, 160, 209311. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38336263>
- Mouro Ferraz Lima, T, Castaldelli-Maia, JM, Apter, G, & Leopoldo, K. (2023). Neurobiological associations between smoking and internalizing disorders. *Int Rev Psychiatry*, 35(5-6), 486-495. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38299645>
- Freibott, CE, Biondi, BE, Rao, SR, Blokhina, E, Dugas, JN, Patts, G et al. (2024). Is Abstinence from Alcohol and Smoking Associated with Less Anxiety and Depressive Symptoms Among People with HIV? *AIDS Behav*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38285292>
- Johnstad, PG. (2023). Unhealthy behaviors associated with mental health disorders: a systematic comparative review of diet quality, sedentary behavior, and cannabis and tobacco use. *Front Public Health*, 11, 1268339. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38249418>
- Sprenger, S, & Anderson, JS. (2024). Dying to Quit: Understanding the Burden of Tobacco in Psychiatric Patients-A Clinical Review. *J Psychiatr Pract*, 30(1), 23-31. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38227724>
- Chen, SJ, Lu, MY, Lai, JC, Bair, MJ, & Cheng, HY. (2023). Sex and age disparities in the influence of tobacco smoking on depression: Evidence from the Taiwan Biobank. *Tob Induc Dis*, 21, 155. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38026502>
- Matsuyama, Y, & Tabuchi, T. (2023). Does Tobacco Smoking Increase Social Isolation? A Mendelian Randomization Study. *Am J Epidemiol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37981720>
- Zhang, R, Yan, W, Manza, P, Shokri-Kojori, E, Demiral, SB, Schwandt, M et al. (2023). Disrupted brain state dynamics in opioid and alcohol use disorder: attenuation by nicotine use. *Neuropsychopharmacology*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37935861>
- Gao, Z, Xiu, M, Liu, J, Wu, F, & Zhang, X. (2023). Smoking, Symptoms Improvement, and Total Antioxidant Capacity in Patients with Drug-naïve First-episode Schizophrenia: A Prospective Cohort Study. *Curr Neuropharmacol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37859307>
- Adil, N, Jamal, A, & Soha, S. (2023). Mental Health Issue Accelerates Aging Faster Than Smoking - Study Rings Alarm For Pakistan Where It Still Remains A Taboo! *J Pak Med Assoc*, 73(10), 2133. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37876096>

- Zvolska, K, Tichopad, A, Stepankova, L, Pankova, A, Adamcekova, Z, Donin, G et al. (2023). Increasing prevalence of mental disorders in smokers seeking treatment of tobacco dependence: a retrospective observational study. *BMC Psychiatry*, 23(1), 621. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37620772>
- Balbuena, L, Peters, E & Speed, D. (2023). Using polygenic risk scores to investigate the evolution of smoking and mental health outcomes in UK biobank participants. *Acta Psychiatr Scand*, e13601. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37607129>
- Vlasceanu, AM, Gradinaru, D, Stan, M, Nitescu, VG, & Baconi, DL. (2023). Relationships between Serum Biomarkers of Oxidative Stress and Tobacco Smoke Exposure in Patients with Mental Disorders. *Antioxidants (Basel)*, 12(6). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37372029>
- Cambron, C. (2022). E-Cigarette Use Is Associated with Increased Psychological Distress among Youth: A Pooled Cross-Sectional Analysis of State-Level Data from 2019 and 2021. *Int J Environ Res Public Health*, 19(18). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36142001>
- Chang, YC, Lee, YH, Chiang, T, & Liu, CT. (2022). Associations of Smoking and Alcohol Consumption with Loneliness, Depression, and Loss of Interest Among Chinese Older Males and Females. *Int J Ment Health Addict*, 1-16. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36101657>
- Du, X, Wu, R, Kang, L, Zhao, L, & Li, C. (2022). Tobacco smoking and depressive symptoms in Chinese middle-aged and older adults: Handling missing values in panel data with multiple imputation. *Front Public Health*, 10, 913636. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36091567>
- Galan, D, Perry, B I, Warriar, V, Davidson, CC, Stupart, O, Easton, D et al. (2022). Applying Mendelian randomization to appraise causality in relationships between smoking, depression and inflammation. *Sci Rep*, 12(1), 15041. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36057695>
- Li, Y, Wu, F, Mu, Q, Xu, K, Yang, S, Wang, P et al. (2022). Metal ions in cerebrospinal fluid: Associations with anxiety, depression, and insomnia among cigarette smokers. *CNS Neurosci Ther*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36168907>
- Schirmbeck, F, van der Ven, E, Boyette, LL, McGuire, P, Valmaggia, LR, Kempton, MJ et al. (2022). Differential trajectories of tobacco smoking in people at ultra-high risk for psychosis: Associations with clinical outcomes. *Front Psychiatry*, 13, 869023. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35942478>
- Forman-Hoffman VL, Hedden SL, Glasheen C, Davies C, and Colpe LJ. The role of mental illness on cigarette dependence and successful quitting in a nationally representative, household-based sample of U.S. Adults. *Annals of Epidemiology*, 2016; 26(7):447–54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27247163>
- Patnode CD, Henderson JT, Thompson JH, Senger CA, Fortmann SP, et al. Behavioral counseling and pharmacotherapy interventions for tobacco cessation in adults, including pregnant women: A review of reviews for the US preventive services task force. *Annals of Internal Medicine*, 2015; 163(8):608–21. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26389650>

Mendelsohn CP, Kirby DP, and Castle DJ. Smoking and mental illness. An update for psychiatrists. *Australas Psychiatry*, 2015; 23(1):37–43. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25512967>

Stockings EA, Bowman JA, Bartlem KM, McElwaine KM, Baker AL, et al. Implementation of a smoke-free policy in an inpatient psychiatric facility: Patient-reported adherence, support, and receipt of nicotine-dependence treatment. *International Journal of Mental Health Nursing*, 2015; 24(4):342–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25970237>

Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, et al. Change in mental health after smoking cessation: Systematic review and meta-analysis. *British Medical Journal*, 2014; 348:g1151. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24524926>

Lawrence D, Hancock KJ, and Kisely S. The gap in life expectancy from preventable physical illness in psychiatric patients in western Australia: Retrospective analysis of population based registers. *British Medical Journal*, 2013; 346:f2539. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23694688>

Williams JM, Steinberg ML, Griffiths KG, and Cooperman N. Smokers with behavioral health comorbidity should be designated a tobacco use disparity group. *American Journal of Public Health*, 2013; 103(9):1549–55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/23865661>

Wye PM, Bowman JA, Wiggers JH, Baker A, Knight J, et al. Smoking restrictions and treatment for smoking: Policies and procedures in psychiatric inpatient units in Australia. *Psychiatric Services*, 2009; 60(1):100–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19114578>

Ratschen E, Britton J, Doody GA, Leonardi-Bee J, and McNeill A. Tobacco dependence, treatment and smoke-free policies: A survey of mental health professionals' knowledge and attitudes. *General Hospital Psychiatry*, 2009; 31(6):576–82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19892217>

Olivier D, Lubman D, and Fraser R. Tobacco smoking within psychiatric inpatient settings: Biopsychosocial perspective. *Australia and New Zealand Journal of Psychiatry*, 2007; 41(7):572–80. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/00048670701392809>

Guydish J, Passalacqua E, Tajima B, and Manser ST. Staff smoking and other barriers to nicotine dependence intervention in addiction treatment settings: A review. *J Psychoactive Drugs*, 2007; 39(4):423–33. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18303699>

Reilly P, Murphy L, and Alderton D. Challenging the smoking culture within a mental health service supportively. *International Journal of Mental Health Nursing*, 2006; 15(4):272–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/17064324/>

Baker A, Ivers R, Bowman J, Butler T, Kay-Lambkin F, et al. Where there's smoke, there's fire: High prevalence of smoking among some sub-populations and recommendations for intervention. *Drug and Alcohol Review*, 2006; 25:85–96. Available from: <http://www.informaworld.com/smpp/content~content=a741424195~db=all~order=page>

de Leon J and Diaz FJ. A meta-analysis of worldwide studies demonstrates an association between schizophrenia and tobacco smoking behaviors. *Schizophr Res*, 2005; 76(2-3):135–57. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15949648>

Ashton M. Policy within mental health services- pro-actively addressing tobacco. in *The Mental Health Service Conference*. Sydney. 2005.

Williams J and Ziedonis D. Addressing tobacco among individuals with a mental illness or an addiction. *Addictive Behaviors*, 2004; 29(6):1067–83. Available from: <https://pubmed.ncbi.nlm.nih.gov/15236808/>

US Department of Health and Human Services. The health consequences of smoking: A report of the surgeon general. Atlanta, Georgia: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2004. Available from: http://www.cdc.gov/tobacco/data_statistics/sgr/index.htm.

Prochaska JJ, Delucchi K, and Hall SM. A meta-analysis of smoking cessation interventions with individuals in substance abuse treatment or recovery. *Journal of Consulting and Clinical Psychology*, 2004; 72(6):1144–56. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15612860>

Lawn S. Systematic barriers to quitting smoking among institutionalised public mental health service populations: A comparison of two Australian sites. *The International Journal of Social Psychiatry*, 2004; 50(3):201–15. Available from: <https://pubmed.ncbi.nlm.nih.gov/15511114/>

Lawn S and Pols R. Nicotine withdrawal: Pathway to aggression and assault in the locked psychiatric ward? *Australasian Psychiatry*, 2003; 11(2):199–203. Available from: <https://journals.sagepub.com/doi/abs/10.1046/j.1039-8562.2003.00548.x>

Degenhardt L and Hall W. Patterns of co-morbidity between alcohol use and other substance use in the Australian population. *Drug and Alcohol Review*, 2003; 22(1):7–13. Available from: <https://pubmed.ncbi.nlm.nih.gov/12745353/>

Bowman J and Walsh R. Smoking intervention within alcohol and other drug treatment services: A selective review with suggestions for practical management. *Drug and Alcohol Review*, 2003; 22(1):73–82. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1080/0959523021000059857>

Reichler H, Baker A, Lewin T, and Carr V. Smoking among in-patients with drug-related problems in an Australian psychiatric hospital. *Drug and Alcohol Review*, 2001; 20(2):231–7. Available from: <http://www.informaworld.com/smpp/content~content=a713659508~db=all~order=page>

McNeill A. Smoking and mental health: A review of the literature. London: Smokefree London Programme, 2001. Available from: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.622.6748&rep=rep1&type=pdf>.

Lawrence D, Holman C, and Jablensky A. Duty to care. Preventable physical illness in people with mental illness. Perth: The University of Western Australia, 2001.

Lawn S. Australians with mental illness who smoke. *The British Journal of Psychiatry*, 2001; 178(1):85. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11136222>

Degenhardt L and Hall W. The relationship between tobacco use, substance-use disorders and mental health: Results from the national survey of mental health and well-being. *Nicotine & Tobacco Research*, 2001; 3(3):225–34. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200110050457>

Ashton CH. Adverse effects of cannabis and cannabinoids. *Br J Anaesth*, 1999; 83(4):637–49. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/10673884>

9A.3.3 Explanations for higher smoking prevalence

Lin, W, Krebs, NM, Zhu, J, Horn, K, Foulds, J, Evins, AE, & Muscat, JE. (2024). Racial Differences in Nicotine Reduction: Pooled Results from Two Double-Blind Randomized Controlled Trials. *J Racial Ethn Health Disparities*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39230651>

Xiong, H, Ma, F, Tang, D, & Liu, D. (2024). Correlations among nicotine dependence, health-related quality of life, and depression in current smokers: a cross-sectional study with a mediation model. *Front Psychiatry*, 15, 1455918. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39257561>

Hanafin, J, Sunday, S, & Clancy, L. (2024). Inequalities in Smoking and E-Cigarette Use in Young Adults With Mental Ill-Health, 20 years After Ireland's Smoking Ban. *Tob Use Insights*, 17, 1179173X241283744. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39290597>

Lamela, D, Pinto, TM, & Jongenelen, I. (2024). Cigarette Smoking in Women Victims of Police-Reported Intimate Partner Violence: The Role of Childhood Maltreatment, Type of Partner Abuse, and Psychological Distress Symptoms. *J Trauma Dissociation*, 1-13. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39140216>

Johnson, EC, Austin-Zimmerman, I, Thorpe, HHA, Levey, DF, Baranger, DAA, Colbert, SMC et al. (2024). Cross-ancestry genetic investigation of schizophrenia, cannabis use disorder, and tobacco smoking. *Neuropsychopharmacology*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38906991>

Koster, M, Mannsdorfer, L, van der Pluijm, M, de Haan, L, Ziermans, T, van Wingen, G, & Vermeulen, J. (2024). The Association Between Chronic Tobacco Smoking and Brain Alterations in Schizophrenia: A Systematic Review of Magnetic Resonance Imaging Studies. *Schizophr Bull*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38824451>

Urun Unal, B, & Marakoglu, K. (2024). Smoking and Depression among Medical School Students: A Cross-Sectional Study from Turkey's Largest Province. *Healthcare (Basel)*, 12(11). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38891206>

Chenoweth, MJ. (2024). Cracking the chicken and egg problem of schizophrenia and substance use: Genetic interplay between schizophrenia, cannabis use disorder, and tobacco smoking. *Neuropsychopharmacology*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38862577>

de With, J, van der Heijden, HS, van Amelsvoort, T, Daemen, M, Simons, C, Alizadeh, B et al. (2024). Correction: The association between childhood trauma and tobacco smoking in patients with

psychosis, unaffected siblings, and healthy controls. *Eur Arch Psychiatry Clin Neurosci*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38753192>

Honeycutt, SC, Lichte, DD, Gilles-Thomas, EA, Mukherjee, A, & Loney, GC. (2024). Acute nicotine administration reduces the efficacy of punishment in curbing remifentanyl consumption in a seeking-taking chain schedule of reinforcement. *Psychopharmacology (Berl)*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38775944>

Voorhies, K, Hecker, J, Lee, S, Hahn, G, Prokopenko, D, McDonald, ML et al. (2024). Examining the Effect of Genes on Depression as Mediated by Smoking and Modified by Sex. *Genes (Basel)*, 15(5). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38790194>

Shevorykin, A, Hyland, BM, Robles, D, Ji, M, Vantucci, D, Bensch, L et al. (2024). Tobacco use, trauma exposure and PTSD: a systematic review. *Health Psychol Rev*, 1-32. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38711288>

Audrain-McGovern, J, Klapac, O, Koita, F, Manikandan, D, & Stone, MD. (2024). The interacting effects of depression symptoms and sweet flavoring on the rewarding and reinforcing value of cigarillo use among young adults. *Drug Alcohol Depend Rep*, 11, 100234. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38646014>

Blaes, SL, Lewis, B, Teitelbaum, S, Reisfeld, G, & Boissoneault, J. (2024). Associations between smoking and pain in early recovery in residential substance use treatment-seekers. *J Subst Use Addict Treat*, 160, 209311. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38336263>

Mouro Ferraz Lima, T, Castaldelli-Maia, JM, Apter, G, & Leopoldo, K. (2023). Neurobiological associations between smoking and internalizing disorders. *Int Rev Psychiatry*, 35(5-6), 486-495. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38299645>

Chirokoff, V, Dupuy, M, Abdallah, M, Fatseas, M, Serre, F, Auriacombe, M et al. (2023). Craving dynamics and related cerebral substrates predict timing of use in alcohol, tobacco, and cannabis use disorders. *Addict Neurosci*, 9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38389954>

Zvolensky, MJ, Shepherd, JM, Clausen, BK, Robison, J, Cano, MA, de Dios, M, & Correa-Fernandez, V. (2024). Posttraumatic stress and probable post traumatic stress disorder as it relates to smoking behavior and beliefs among trauma exposed hispanic persons who smoke. *J Behav Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38409553>

Johnstad, PG. (2023). Unhealthy behaviors associated with mental health disorders: a systematic comparative review of diet quality, sedentary behavior, and cannabis and tobacco use. *Front Public Health*, 11, 1268339. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38249418>

de With, J, van der Heijden, S, van Amelsvoort, T, Daemen, M, Simons, C, Alizadeh, B et al. (2024). The association between childhood trauma and tobacco smoking in patients with psychosis, unaffected siblings, and healthy controls. *Eur Arch Psychiatry Clin Neurosci*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38231398>

Farris, SG, Smith, JE, Steinberg, DR, Altman, BR, Lambert-Messerlian, GM, Dunsiger, SI et al. (2023). Methodological approach for an integrated female-specific study of anxiety and smoking comorbidity. *Front Psychiatry*, 14, 1267753. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38076702>

van de Weijer, MP, Vermeulen, J, Schranter, A, Munafo, MR, Verweij, KJH, & Treur, JL. (2024). The potential role of gray matter volume differences in the association between smoking and depression: A narrative review. *Neurosci Biobehav Rev*, 156, 105497. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38100958>

Zvolensky, MJ, Shepherd, JM, Clausen, BK, de Dios, M, Cano, MA, & Redmond, BY. (2023). Differences in smoking behavior and beliefs about abstinence among Latinx individuals with and without depression who smoke cigarettes. *J Ethn Subst Abuse*, 1-19. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38108308>

Chen, SJ, Lu, MY, Lai, JC, Bair, MJ, & Cheng, HY. (2023). Sex and age disparities in the influence of tobacco smoking on depression: Evidence from the Taiwan Biobank. *Tob Induc Dis*, 21, 155. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38026502>

Gollan, JK, Liverant, G, Jao, NC, Lord, KA, Whitton, AE, Hogarth, L et al. (2023). Depression Severity Moderates Reward Learning Among Smokers with Current or Past Major Depressive Disorder in a Smoking Cessation Randomized Clinical Trial. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37943674>

Hahn, B, & Holzel, ES. (2023). Nicotine insensitivity and cue-controlled smoking behavior in people with schizophrenia. *Schizophr Res*, 262, 51-52. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37922844>

Lund, MK, Hojlund, M, & Lund, LC. (2023). Smoking and severe mental illness: Drawing inference on the directionality of the association requires careful consideration of the temporality of the data. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37933520>

Youssef, L, Salameh, P, Sacre, H, Akel, M, Hallit, S, & Obeid, S. (2023). Association between alexithymia, emotional intelligence, smoking addiction, and alcohol use disorder among a sample of Lebanese adults. *PLoS One*, 18(11), e0295114. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38033004>

He, M, Zhou, J, Li, X, & Wang, R. (2023). Investigating the causal effects of smoking, sleep, and BMI on major depressive disorder and bipolar disorder: a univariable and multivariable two-sample Mendelian randomization study. *Front Psychiatry*, 14, 1206657. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37900287>

Lunding, SH, Ueland, T, Aas, M, Hoegh, MC, Werner, MCF, Rodevand, L et al. (2023). Tobacco smoking related to childhood trauma mediated by cognitive control and impulsiveness in severe mental disorders. *Schizophr Res*, 261, 236-244. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37806047>

Luo, Z, Xu, W, Jiang, S, Zhou, Q, Guan, Y, Li, L et al. (2023). The mediating role of negative emotions in the relationship between smoking and health-related quality of life among Chinese individuals: A cross-sectional study. *Tob Induc Dis*, 21, 135. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37849528>

North, CS, & Pfefferbaum, B. (2023). Tobacco and Illicit Drug Use and Drug Use Disorders Among Employees of Businesses Affected by the 9/11 Attacks on the World Trade Center in New York City. *J Dual Diagn*, 19(4), 180-188. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37796633>

Papadosifaki, G, Psarra, V, Touloumis, C, Tzavara, C, Farsalinos, K, Sakellari, E et al. (2023). Perceptions and attitudes of people with severe mental disorders towards smoking in Greece. *Psychiatriki*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37793036>

Thomas, JE, Pasch, KE, Nathan Marti, C, & Loukas, A. (2024). Depressive symptoms prospectively increase risk for new onset cigarette and ENDS dependence symptoms. *Addict Behav*, 148, 107870. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37776758>

Ariaratnam, S, Kee, CC, Krishnapillai, AD, Sanaudi, R, Tohit, NM, Ho, KB et al. (2023). Smoking status and its relationship with depression among the elderly population in Malaysia: Findings from the National Health and Morbidity Survey 2018. *Tob Induc Dis*, 21, 109. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37654502>

Dawes, MH, Estave, PM, Albertson, SE, Wallace, CW, Holleran, KM, & Jones, SR. (2023). Nicotine modifies cocaine responding in a concurrent self-administration model. *Drug Alcohol Depend*, 251, 110960. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37703771>

Nguyen, N, Peyser, ND, Olgin, JE, Pletcher, MJ, Beatty, AL, Modrow, MF et al (2023). Associations between tobacco and cannabis use and anxiety and depression among adults in the United States: Findings from the COVID-19 citizen science study. *PLoS One*, 18(9), e0289058. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37703257>

Pineles, SL, Ni, P, Pandey, S, Japuntich, SJ, Cesare, N, Shor, R et al. (2023). Tobacco withdrawal-induced changes in sensorimotor filtering as a predictor of smoking lapse in trauma-exposed individuals. *Addict Behav*, 148, 107868. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37774527>

Tsukahara, M, So, R, Yoshimura, Y, Yamashita, R, Yada, Y, Kodama, M et al. (2023). Effect of smoking habits and concomitant valproic acid use on relapse in patients with treatment-resistant schizophrenia receiving clozapine: A 1-year retrospective cohort study. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37681448>

Zhu, T, Wang, W, Chen, Y, Kranzler, HR, Li, CR, & Bi, J. (2023). Machine Learning of Functional Connectivity to Biotype Alcohol and Nicotine Use Disorders. *Biol Psychiatry Cogn Neurosci Neuroimaging*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37696489>

Balbuena, L, Peters, E & Speed, D. (2023). Using polygenic risk scores to investigate the evolution of smoking and mental health outcomes in UK biobank participants. *Acta Psychiatr Scand*, e13601. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37607129>

Pagano, A, Le, T, Fong, T, Reid, R, Delucchi, K, & Guydish, J. (2023). Gambling, tobacco use, and health among individuals in substance use disorder treatment. *Am J Addict*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37543853>

Zhang, M, Tang, J, Li, W, Xue, K, Wang, Z, Chen, Y L. (2023). Schizophrenia mediating the effect of smoking phenotypes on antisocial behavior: A Mendelian randomization analysis. *CNS Neurosci Ther*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37650156>

DeAtley, T, Stone, MD, Johnson, AC, Melissa, M, Audrain-McGovern, J, & Strasser, AA. (2023). Differences in biobehavioral measures of cigarette smoking by depression symptomology. *Addict Behav*, 146, 107800. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37437421>

Yang, X, Cheng, S, Li, C, Pan, C, Liu, L, Meng, P et al. (2023). Evaluating the interaction between 3'aQTL and alcohol consumption/smoking on anxiety and depression: 3'aQTL-by-environment interaction study in UK Biobank cohort. *J Affect Disord*, 338, 518-525. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37390921>

Pang, RD, Tucker, C. J., Guillot, C. R., Belcher, B., & Kirkpatrick, M. G. (2023). Associations of DHEA(S) with negative and positive affect in people who smoke daily with elevated and low depression symptoms: A pilot laboratory study. *Addict Behav*, 146, 107801. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37423068>

Schroder, B, & Muhlberger, A. (2023). Measuring attentional bias in smokers during and after psychosocial stress induction with a Trier Social Stress Test in virtual reality via eye tracking. *Front Psychol*, 14, 1129422. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37063522>

Klymova, S, & Cavanaugh, C. (2023). The Impact of Posttraumatic Stress Disorder, Ethnic Discrimination, and Nativity on Nicotine Dependence among Female Hispanic Survivors of Physical Intimate Partner Violence. *Subst Use Misuse*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36798007>

Firth, J, Wootton, RE, Sawyer, C, & Taylor, GM. (2023). Clearing the air: clarifying the causal role of smoking in mental illness. *World Psychiatry*, 22(1), 151-152. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36640385>

Grabowska, K, Ziemichod, W, & Biala, G. (2023). Recent Studies on the Development of Nicotine Abuse and Behavioral Changes Induced by Chronic Stress Depending on Gender. *Brain Sci*, 13(1). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36672102>

Kascakova, N, Petrikova, M, Furstova, J, Hasto, J, Geckova, AM, & Tavel, P. (2022). Associations of childhood trauma with long-term diseases and alcohol and nicotine use disorders in Czech and Slovak representative samples. *BMC Public Health*, 22(1), 1769. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36123641>

Compton, MT, Zern, A, Langlois, S, & Ashekun, O. (2022). Associations Between Adverse Childhood Experiences and Tobacco, Alcohol, and Drug Use Among Individuals with Serious Mental Illnesses in Public-Sector Treatment Settings. *Community Ment Health J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35976478>

Qiu, X, Han, X, Wang, Y, Ding, W, Sun, Y, Lei, H et al. (2022). Reciprocal modulation between cigarette smoking and internet gaming disorder on participation coefficient within functional brain networks. *Brain Imaging Behav*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36018530>

Diaz-Martinez, J, Delgado-Enciso, I, Campa, A, Tamargo, JA, Martin, HR, Johnson, A et al. (2022). Stress Increases the Association between Cigarette Smoking and Mental Disorders, as Measured by the COVID-19-Related Worry Scale, in the Miami Adult Studies on HIV (MASH) Cohort during the Pandemic. *Int J Environ Res Public Health*, 19(13). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35805866>

Farooqui, M, Shoaib, S, Afaq, H, Quadri, S, Zaina, F, Baig, A et al. (2022). Bidirectionality of Smoking and Depression in Adolescents: a Systemic Review. *Trends Psychiatry Psychother*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35738567>

Stevenson, J, Miller, CL, Martin, K, Mohammadi, L & Lawn, S. (2022). Investigating the reciprocal temporal relationships between tobacco consumption and psychological disorders for youth: an international review. *BMJ Open*, 12(6), e055499. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35697442>

Wilkinson AV, Swann AC, Graham DP, Patriquin MA, Salas R, et al. Emotional self-regulation, impulsivity, 5-HTT and tobacco use behavior among psychiatric inpatients. *J Affect Disord*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35623482>

Wei S, Wang D, Zhou H, Xia L, Tian Y, et al. Demographics, clinical characteristics and cognitive symptoms of heavy smokers and non-heavy smokers in chinese male patients with chronic schizophrenia. *European Archives of Psychiatry and Clinical Neuroscience*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35474549>

Munafo M. Growing evidence for a causal role for smoking in mental health. *Nicotine & Tobacco Research*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35100432>

Del Casale A, Paolini M, Gentile G, Borro M, Zocchi C, et al. Dopamine drd2 and drd3 polymorphisms involvement in nicotine dependence in patients with treatment-resistant mental disorders. *Journal of Personalized Medicine*, 2022; 12(4). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35455685>

Andreasen JT. [tobacco dependence, anxiety, and depression]. *Ugeskr Laeger*, 2022; 184(14). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35410654>

van der Heijden HS, Schirmbeck F, McGuire P, Valmaggia LR, Kempton MJ, et al. Association between tobacco use and symptomatology in individuals at ultra-high risk to develop a psychosis: A longitudinal study. *Schizophr Res*, 2021; 236:48-53. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34390981>

Tindimwebwa LK, Ajayi AI, and Adeniyi OV. Prevalence and determinants of tobacco use amongst south african adults with mental illness in the eastern cape. *S Afr J Psychiatr*, 2021; 27:1637. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34230866>

Snell M, Harless D, Shin S, Cunningham P, and Barnes A. A longitudinal assessment of nicotine dependence, mental health, and attempts to quit smoking: Evidence from waves 1-4 of the population assessment of tobacco and health (path) study. *Addictive Behaviors*, 2021; 115:106787. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33383566>

Pivac N, Nedic Erjavec G, Sagud M, Nikolac Perkovic M, Tudor L, et al. The association between bdnf c270t genetic variants and smoking in patients with mental disorders and in healthy controls. *Prog Neuropsychopharmacol Biol Psychiatry*, 2021; 113:110452. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34637871>

Piirtola M, Kaprio J, Baker TB, Piasecki TM, Piper ME, et al. The associations of smoking dependence motives with depression among daily smokers. *Addiction*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33629475>

Pang RD, Tucker CJ, D'Orazio LM, Weinberger AH, and Guillot CR. Affect and subjective cognitive functioning by depression symptom levels during naturalistic cigarette smoking in premenopausal females who smoke daily. *Psychol Addict Behav*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33844567>

Mason TB, Tackett AP, Smith CE, and Leventhal AM. Tobacco product use for weight control as an eating disorder behavior: Recommendations for future clinical and public health research. *Int J Eat Disord*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34866222>

Mamey MR, Burns GL, Barbosa-Leiker C, Smith CL, and McPherson S. Parallel growth modeling to better understand smoking with stimulant use outcomes during an integrated treatment trial. *Experimental and Clinical Psychopharmacology*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33570975>

Hu H, Liu W, Zhang S, Pan J, and Zheng X. Depression mediates the relationship between smoking and pain: Evidence from a nationally representative study in a low- and middle-income country. *Addictive Behaviors*, 2021; 119:106937. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33848759>

De Angelis F, Wendt FR, Pathak GA, Tylee DS, Goswami A, et al. Drinking and smoking polygenic risk is associated with childhood and early-adulthood psychiatric and behavioral traits independently of substance use and psychiatric genetic risk. *Transl Psychiatry*, 2021; 11(1):586. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34775470>

Barkhuizen W, Dudbridge F, and Ronald A. Genetic overlap and causal associations between smoking behaviours and mental health. *Sci Rep*, 2021; 11(1):14871. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34290290>

Al-Soufi L and Costas J. Colocalization of association signals at nicotinic acetylcholine receptor genes between schizophrenia and smoking traits. *Drug and Alcohol Dependence*, 2021; 220:108517. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33454625>

Wootton RE, Richmond RC, Stuijzand BG, Lawn RB, Sallis HM, et al. Evidence for causal effects of lifetime smoking on risk for depression and schizophrenia: A mendelian randomisation study. *Psychological Medicine*, 2020; 50(14):2435–43. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31689377>

Okubo R and Tabuchi T. Smoking and drinking among patients with mental disorders: Evidence from a nationally representative Japanese survey. *J Affect Disord*, 2020; 279:443-50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33120245>

Masiero M, Cropley M, and Pravettoni G. Increasing smoking cessation adherence: Do we need to consider the role of executive function and rumination? *Eur J Psychol*, 2020; 16(1):1-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33680166>

Konjevod M, Nikolac Perkovic M, Svob Strac D, Uzun S, Nedic Erjavec G, et al. Significant association of mu-opioid receptor 1 haplotype with tobacco smoking in healthy control subjects but not in patients with schizophrenia and alcohol dependence. *Psychiatry Res*, 2020; 291:113278. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32763540>

Friedman AS. Smoking to cope: Addictive behavior as a response to mental distress. *J Health Econ*, 2020; 72:102323. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32505043>

Brady KT. Social determinants of health and smoking cessation: A challenge. *American Journal of Psychiatry*, 2020; 177(11):1029-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33135467>

Borges AM, Yang MJ, Farris SG, Zvolensky M, and Leyro TM. Examining the role of emotion regulation in the bidirectional relation between physiological and subjective stress response among daily cigarette smokers. *Pers Individ Dif*, 2020; 155. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32863505>

Ben Salah A, Nakajima M, DeAngelis BN, and al'Absi M. Effects of tobacco addiction on links between early life adversities, sleep disturbance, and depression: A moderated mediation approach. *Preventive Medicine Reports*, 2020; 20:101225. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33194536>

White AT, Williams P, Anand V, Benjamin J, and Kim S. Cigarettes and straws: Late positive potential modulation in mental illness and nicotine addiction. *Conf Proc IEEE Eng Med Biol Soc*, 2019; 2019:1-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31945829>

Seng S, Otachi JK, and Okoli CTC. Reasons for tobacco use and perceived tobacco-related health risks in an inpatient psychiatric population. *Issues Ment Health Nurs*, 2019:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31398069>

Qurashi I, Stephenson P, Nagaraj C, Chu S, Drake R, et al. Changes in smoking status, mental state and plasma clozapine concentration: Retrospective cohort evaluation. *BJPsych Bull*, 2019:1-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31391141>

Han JW and Lee H. The longitudinal relationships between depression and smoking in hardcore smokers using autoregressive cross-lagged modeling. *J Korean Acad Nurs*, 2019; 49(1):69-79. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30837444>

Chang LH, Whitfield JB, Liu M, Medland SE, Hickie IB, et al. Associations between polygenic risk for tobacco and alcohol use and liability to tobacco and alcohol use, and psychiatric disorders in an

independent sample of 13,999 Australian adults. *Drug and Alcohol Dependence*, 2019; 205:107704. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31731259>

MacCabe JH. It is time to start taking tobacco seriously as a risk factor for psychosis: Self-medication cannot explain the association. *Acta Psychiatrica Scandinavica*, 2018; 138(1):3-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29956305>

Li H, Chen J, Chen C, Xu Z, Xu J, et al. Csf glutamate level decreases in heavy smokers and negatively correlates with bdi scores. *Psychiatry Res*, 2018; 270:627-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30384282>

Harvey PD. Cigarette smoking, cognitive performance, and severe mental illness: Quitting smoking really does seem to matter. *American Journal of Psychiatry*, 2018; 175(11):1054-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30380939>

Hammett PJ, Lando HA, Taylor BC, Widome R, Erickson DJ, et al. The relationship between smoking cessation and binge drinking, depression, and anxiety symptoms among smokers with serious mental illness. *Drug and Alcohol Dependence*, 2018; 194:128-35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30439609>

Epperson AE, Anzai NE, and Prochaska JJ. Natural american spirit brand preference among smokers with mental illness. *Tobacco Induced Diseases*, 2018; 16. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31210981>

Clark V, Conrad AM, Lewin TJ, Baker AL, Halpin SA, et al. Addiction vulnerability: Exploring relationships among cigarette smoking, substance misuse, and early psychosis. *Journal of Dual Diagnosis*, 2018; 14(2):78–88. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29261427>

Boksa P. Smoking, psychiatric illness and the brain. *J Psychiatry Neurosci*, 2017; 42(3):147-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28440208>

Bhavsar V, Jauhar S, Murray RM, Hotopf M, Hatch SL, et al. Tobacco smoking is associated with psychotic experiences in the general population of south london. *Psychological Medicine*, 2017:1-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28655360>

Tidey JW. A behavioral economic perspective on smoking persistence in serious mental illness. *Preventive Medicine*, 2016; 92:31-5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27196141>

Sundgren E, Hallqvist J, and Fredriksson L. Health for smokers with schizophrenia - a struggle to maintain a dignified life. *Disability and Rehabilitation*, 2016; 38(5):416–22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25958996>

Robson D, Yates M, Craig TJ, Healey A, and McNeill A. Time to smoke: Facilitating smoking breaks in mental health inpatient settings. *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27085082>

Potvin S, Lungu O, Lipp O, Lalonde P, Zaharieva V, et al. Increased ventro-medial prefrontal activations in schizophrenia smokers during cigarette cravings. *Schizophr Res*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27005897>

Peckham E, Bradshaw TJ, Brabyn S, Knowles S, and Gilbody S. Exploring why people with smi smoke and why they may want to quit: Baseline data from the scimitar rct. *J Psychiatr Ment Health Nurs*, 2016; 23(5):282–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26147943>

Pal A and Balhara YP. A review of impact of tobacco use on patients with co-occurring psychiatric disorders. *Tob Use Insights*, 2016; 9:7–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26997871>

Garey L, Farris SG, Schmidt NB, and Zvolensky MJ. The role of smoking-specific experiential avoidance in the relation between perceived stress and tobacco dependence, perceived barriers to cessation, and problems during quit attempts among treatment-seeking smokers. *J Contextual Behav Sci*, 2016; 5(1):58–63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27231668>

Garey L, Bakhshaie J, Brandt CP, Langdon KJ, Kauffman BY, et al. Interplay of dysphoria and anxiety sensitivity in relation to emotion regulatory cognitions of smoking among treatment-seeking smokers. *American Journal on Addictions*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27122303>

Farris SG, Paulus DJ, Gonzalez A, Mahaffey BL, Bromet EJ, et al. Posttraumatic stress symptoms and body mass index among world trade center disaster-exposed smokers: A preliminary examination of the role of anxiety sensitivity. *Psychiatry Res*, 2016; 241:135–40. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27173658>

Zhang XY, Rao WW, Yu Q, Yu Y, Kou C, et al. Association of the manganese superoxide dismutase gene ala-9val polymorphism with age of smoking initiation in male schizophrenia smokers. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26544677>

Peechatka AL, Whitton AE, Farmer SL, Pizzagalli DA, and Janes AC. Cigarette craving is associated with blunted reward processing in nicotine-dependent smokers. *Drug and Alcohol Dependence*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26233484>

Mahaffey BL, Gonzalez A, Farris SG, Zvolensky MJ, Bromet EJ, et al. Smoking to regulate negative affect: Disentangling the relationship between posttraumatic stress and emotional disorder symptoms, nicotine dependence, and cessation-related problems. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26304694>

Farris SG, Leventhal AM, Schmidt NB, and Zvolensky MJ. Anxiety sensitivity and pre-cessation smoking processes: Testing the independent and combined mediating effects of negative affect-reduction expectancies and motives. *J Stud Alcohol Drugs*, 2015; 76(2):317–25. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25785807>

Braithwaite RS, Fang Y, Tate J, Mentor SM, Bryant KJ, et al. Do alcohol misuse, smoking, and depression vary concordantly or sequentially? A longitudinal study of hiv-infected and matched

uninfected veterans in care. *AIDS Behav*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26187007>

Bakhshaie J, Zvolensky MJ, Salazar A, Vujanovic AA, and Schmidt NB. Anxiety sensitivity and smoking behavior among trauma-exposed daily smokers: The explanatory role of smoking-related avoidance and inflexibility. *Behav Modif*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26538275>

Audrain-McGovern J, Leventhal AM, and Strong DR. The role of depression in the uptake and maintenance of cigarette smoking. *Int Rev Neurobiol*, 2015; 124:209–43. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26472531>

Ameringer KJ, Chou CP, Sussman S, Unger JB, and Leventhal AM. Identifying shared latent dimensions of psychological symptoms: Implications for the psychological correlates of smoking. *J Psychopathol Behav Assess*, 2015; 37(3):454–68. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26478654>

Choi N and Dinitto D. Drinking, smoking, and psychological distress in middle and late life. *Aging & Mental Health*, 2011; 15(6):720–31. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/21562984>

9A.3.4 Barriers to be overcome when quitting

Yang, T, Cottrell, RR, & Wu, D. (2024). Is unassisted smoking cessation choice and success associated with high mental stress? Evidence from six cities in China. *Tob Induc Dis*, 22. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/39403276>

Mahreen, ZSH, Harjit Singh, KK, Ng, CWL, & Low, LTK. (2024). Challenges to smoking cessation in patients with substance use disorders. *Ann Acad Med Singap*, 53(2), 124-126. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/38920238>

Castaldelli-Maia, JM, & Apter, G. (2023). Breaking Down Barriers to Smoking Cessation for Mental Health Patients in France. *Psychiatr Serv*, 74(11), 1212. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/37908082>

Branas, M, Frankenburg, FR, Temes, CM, Fitzmaurice, GM, & Zanarini, MC. (2023). Longitudinal Description and Prediction of Smoking Among Borderline Patients: An 18-Year Follow-Up Study. *J Clin Psychiatry*, 84(6). Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/37943989>

Abufarsakh, B, Scarduzio, J, & Okoli, C. (2023). Smoking Cessation Barriers among Individuals with Mental Illnesses: A Qualitative Study of Patients' and Healthcare Providers' Perspectives. *Issues Ment Health Nurs*, 1-8. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/37819773>

Iyehen, EO, Omoruyi, OO, Rowa-Dewar, N, & Dobbie, F. (2023). Exploring the barriers and facilitators to the uptake of smoking cessation services for people in treatment or recovery from problematic drug or alcohol use: A qualitative systematic review. *PLoS One*, 18(7), e0288409. Retrieved from

<https://www.ncbi.nlm.nih.gov/pubmed/37440505>

Yang, X, Zhang, L, Lin, H, Lin, H, Cao, W, & Chang, C. (2023). Psychiatric symptoms and intentions to quit smoking: How regularity and volume of cigarette consumption moderate the relationship. *Tob Induc Dis*, 21, 78. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37323507>

Addicott, MA, Oliveto, AH, & Daughters, SB. (2023). Smoking status affects cognitive, emotional and neural-connectivity response to distress-inducing auditory feedback. *Drug Alcohol Depend*, 246, 109855. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37028104>

Kock, L, Brown, J, Cox, S, McNeill, A, Robson, D, Shahab, L et al. (2023). Association of psychological distress with smoking cessation, duration of abstinence from smoking, and use of non-combustible nicotine-containing products: A cross-sectional population survey in Great Britain. *Addict Behav*, 138, 107570. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36493683>

Abarno, CN, Chapman, HM, & Copeland, AL. (2022). The impact of depression on opioid withdrawal symptoms among smokers with opioid use disorder receiving medication-assisted opioid detoxification. *Exp Clin Psychopharmacol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36074624>

Nagawa, CS, Pbert, L, Wang, B, Cutrona, SL, Davis, M, Lemon, SC, & Sadasivam, RS. (2022). Association between family or peer views towards tobacco use and past 30-day smoking cessation among adults with mental health problems. *Prev Med Rep*, 28, 101886. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35855923>

Xie S, Minami H, Kumar DS, Hecht J, Bloom EL, et al. Readiness to quit smoking among smokers in substance use treatment: Associations with stress, substance use severity, relapse concerns and gender. *J Subst Use*, 2021; 26(6):669-76. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34899051>

Shpigel DM, Gittleman JM, Estey D, Birchwale JT, Rosensweig SR, et al. Psychosocial and psychiatric-related stress and cigarette smoking among black and latinx adults with psychiatric disorders. *J Ethn Subst Abuse*, 2021:1-25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34139960>

Selva Kumar D, Nahvi S, Rivera-Mindt M, Arnsten J, and Minami H. The impact of coping with stressful events on negative affect and cravings among smokers with mood disorders. *Nicotine & Tobacco Research*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34918163>

Quinn MH, Olonoff M, Bauer AM, Fox E, Jao N, et al. History and correlates of smoking cessation behaviors among individuals with current or past major depressive disorder enrolled in a smoking cessation trial. *Nicotine & Tobacco Research*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34259871>

Kertes J, Neumark Y, Grunhaus L, and Stein-Reisner O. Comparison of perceptions and smoking cessation experiences between smokers with and without serious mental illness in a large health maintenance organization. *Journal of Dual Diagnosis*, 2021:1-12. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34651553>

Johnston J, Xia J, Yau MTK, Wang JC, Okoli CTC, et al. The impact of psychiatric disorder diagnosis on motivation to quit and stage of change among patients at a hospital-based outpatient smoking

cessation clinic. *Journal of Dual Diagnosis*, 2021;1-22. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33600740>

Correa JB, Lawrence D, McKenna BS, Gaznick N, Saccone PA, et al. Psychiatric comorbidity and multimorbidity in the eagles trial: Descriptive correlates and associations with neuropsychiatric adverse events, treatment adherence, and smoking cessation. *Nicotine & Tobacco Research*, 2021; 23(10):1646–55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33788933>

Schoetsanitis G, Haen E, Conca A, Piacentino D, Ridders F, et al. Lack of smoking effects on pharmacokinetics of oral paliperidone-analysis of a naturalistic therapeutic drug monitoring sample. *Pharmacopsychiatry*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32767297>

Olando Y, Kuria MW, Mathai M, and Huffman MD. Barriers and facilitators to cessation among tobacco users with concomitant mental illness attending group behavioral tobacco cessation: A qualitative study. *Tob Prev Cessat*, 2020; 6:46. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32954059>

Jaen-Moreno MJ, Sarramea F, and Balanza-Martinez V. Smoking cessation in severe mental illness: Challenges and opportunities in the COVID-19 times. *Adicciones*, 2020; 0(0):1579. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33338250>

Ho RKS, Lee GMT, Fok PWY, Chan HCH, and Ching JKW. Characteristics of chinese smokers with psychotic disorders and their predictors on smoking cessation in hong kong. *Tob Prev Cessat*, 2020; 6:7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32548344>

Garey L, Rogers AH, Manning K, Smit T, Derrick JL, et al. Effects of smoking cessation treatment attendance on abstinence: The moderating role of psychologically based behavioral health conditions. *Journal of Substance Abuse Treatment*, 2020; 109:1-7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31856945>

Carstens C and Linley J. Desire to quit smoking in an outpatient population of persons with serious mental illness. *J Behav Health Serv Res*, 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32342438>

Vermeulen JM, van den Brink W, and de Haan L. The unhealthy chicken or the unhealthy egg: Quitting smoking matters: Response to hajek et al. *American Journal of Psychiatry*, 2019; 176(7):575-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31256619>

Scherf-Clavel M, Deckert J, Menke A, and Unterecker S. Smoking is associated with lower dose-corrected serum concentrations of escitalopram. *J Clin Psychopharmacol*, 2019. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31343441>

Sarramea Crespo F, Jaen-Moreno MJ, Gutierrez-Rojas L, Balanza-Martinez V, Garcia-Alvarez L, et al. "Readiness to change" predicts efficacy of reduction among smokers with severe mental illness. *Eur Addict Res*, 2019:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31163435>

Okoli CTC and Seng S. Factors associated with the initial sensations of smoking among smokers with mental illness. *Issues Ment Health Nurs*, 2019:1-7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31188699>

Metse AP, Stockings E, Bailey J, Regan T, Bartlem K, et al. Rates of retention of persons with a mental health disorder in outpatient smoking cessation and reduction trials, and associated factors: Protocol for a systematic review and meta-analysis. *BMJ Open*, 2019; 9(9):e030646. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31488491>

Lubitz SF, Flitter A, Wileyto EP, Ziedonis D, Stevens N, et al. History and correlates of smoking cessation behaviors among smokers with serious mental illness. *Nicotine & Tobacco Research*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31816049>

Keller-Hamilton B, Moe AM, Breitborde NJK, Lee A, and Ferketich AK. Reasons for smoking and barriers to cessation among adults with serious mental illness: A qualitative study. *J Community Psychol*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31102293>

Keizer I, Gex-Fabry M, Croquette P, Humair JP, and Khan AN. Tobacco craving and withdrawal symptoms in psychiatric patients during a motivational enhancement intervention based on a 26-hour smoking abstinence period. *Tob Prev Cessat*, 2019; 5:22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32411885>

Kalkhoran S, Thorndike AN, Rigotti NA, Fung V, and Baggett TP. Cigarette smoking and quitting-related factors among US adult health center patients with serious mental illness. *Journal of General Internal Medicine*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30783880>

Scherf-Clavel M, Samanski L, Hommers LG, Deckert J, Menke A, et al. Analysis of smoking behavior on the pharmacokinetics of antidepressants and antipsychotics: Evidence for the role of alternative pathways apart from cyp1a2. *Int Clin Psychopharmacol*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30557209>

Brunette MF, Ferron JC, Aschbrenner KA, Pratt SI, Geiger P, et al. Attitudes about smoking cessation treatment, intention to quit, and cessation treatment utilization among young adult smokers with severe mental illnesses. *Addictive Behaviors*, 2018; 89:248-55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30343187>

Aschbrenner KA, Bobak C, Schneider EJ, Naslund JA, Brunette MF, et al. Egocentric social networks and smoking among adults with serious mental illness. *Transl Behav Med*, 2018; 8(4):531-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30016519>

Farris SG, Aston ER, Zvolensky MJ, Abrantes AM, and Metrik J. Psychopathology and tobacco demand. *Drug and Alcohol Dependence*, 2017; 177:59-66. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28575783>

Christiansen BA, Carbin J, TerBeek E, and Fiore MC. Helping smokers with severe mental illness who do not want to quit. *Substance Use and Misuse*, 2017;1-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29161166>

Thorndike AN, Achtyes ED, Cather C, Pratt S, Pachas GN, et al. Weight gain and 10-year cardiovascular risk with sustained tobacco abstinence in smokers with serious mental illness: A subgroup analysis of a randomized trial. *Journal of Clinical Psychiatry*, 2016; 77(3):e320–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27046320>

Sheals K, Tombor I, McNeill A, and Shahab L. A mixed-method systematic review and meta-analysis of mental health professionals' attitudes toward smoking and smoking cessation among people with mental illnesses. *Addiction*, 2016; 111(9):1536–53. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/27003925>

Sharma R, Gartner CE, and Hall WD. The challenge of reducing smoking in people with serious mental illness. *The Lancet Respiratory Medicine*, 2016; 4(10):835-44. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27707462>

Powers J, Carroll AJ, Veluz-Wilkins AK, Blazekovic S, Gariti P, et al. Is the effect of anhedonia on smoking cessation greater for women versus men? *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27287389>

Penberthy JK, Penberthy JM, Harris MR, Nanda S, Ahn J, et al. Are smoking cessation treatments associated with suicidality risk? An overview. *Subst Abuse*, 2016; 10:19–30. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27081311>

Metse AP, Wiggers J, Wye P, Moore L, Clancy R, et al. Smoking and environmental characteristics of smokers with a mental illness, and associations with quitting behaviour and motivation; a cross sectional study. *BMC Public Health*, 2016; 16(1):332. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27080019>

Farris SG, Zvolensky MJ, and Schmidt NB. Difficulties with emotion regulation and psychopathology interact to predict early smoking cessation lapse. *Cognit Ther Res*, 2016; 40(3):357–67. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27239081>

Dedert EA, Hicks TA, Dennis PA, Calhoun PS, and Beckham JC. Roles of inter-individual differences and intra-individual acute elevations in early smoking lapse in people with posttraumatic stress disorder. *Addictive Behaviors*, 2016; 60:171–6. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27155244>

Buckner JD, Langdon KJ, Jeffries ER, and Zvolensky MJ. Socially anxious smokers experience greater negative affect and withdrawal during self-quit attempts. *Addictive Behaviors*, 2016; 55:46–9.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26790139>

Zale EL, Maisto SA, and Ditre JW. Anxiety and depression in bidirectional relations between pain and smoking: Implications for smoking cessation. *Behav Modif*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26467214>

Wong DC, Chan SS, and Lam TH. Depressive symptoms delayed quit attempts and shortened abstinence in young smokers of the hong kong youth quitline. *Nicotine & Tobacco Research*, 2015.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25818111>

Tulloch HE, Pipe AL, Clyde MJ, Reid RD, and Els C. The quit experience and concerns of smokers with psychiatric illness. *American Journal of Preventive Medicine*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26711162>

Sullivan TP, Flanagan JC, Dudley DN, Holt LJ, Mazure CM, et al. Correlates of smoking status among women experiencing intimate partner violence: Substance use, posttraumatic stress, and coping.

American Journal on Addictions, 2015; 24(6):546–53. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26249027>

Simioni N and Cottencin O. Resurgence of anorexic symptoms during smoking cessation in patients with a history of anorexia nervosa: An unseen problem?-report of two cases. *Int J Eat Disord*, 2015.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26016609>

Pawlina MM, Rondina Rde C, Espinosa MM, and Botelho C. Depression, anxiety, stress, and motivation over the course of smoking cessation treatment. *J Bras Pneumol*, 2015; 41(5):433–9.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26578135>

McLeish AC, Johnson AL, Avallone KM, and Zvolensky MJ. Evaluating the role of anxiety sensitivity in barriers to cessation and reasons for quitting among smokers with asthma. *Psychol Health Med*,

2015:1–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26033273>

Ditre JW, Langdon KJ, Kosiba JD, Zale EL, and Zvolensky MJ. Relations between pain-related anxiety, tobacco dependence, and barriers to quitting among a community-based sample of daily smokers.

Addictive Behaviors, 2015; 42:130–5. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25462660>

Brown-Johnson CG, Cataldo JK, Orozco N, Lisha NE, Hickman NJ, 3rd, et al. Validity and reliability of the internalized stigma of smoking inventory: An exploration of shame, isolation, and discrimination in smokers with mental health diagnoses. *American Journal on Addictions*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25930661>

Brown CH, Medoff D, Dickerson FB, Fang LJ, Lucksted A, et al. Factors influencing implementation of smoking cessation treatment within community mental health centers. *Journal of Dual Diagnosis*,

2015; 11(2):145–50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25985201>

Anzai N, Young-Wolff KC, and Prochaska JJ. Symptom severity and readiness to quit among hospitalized smokers with mental illness. *Psychiatric Services*, 2015; 66(4):443–4. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25828988>

Albanese BJ, Allan NP, Boffa JW, Chavarria J, Raines AM, et al. Suicidality prospectively predicts greater urges to smoke following a cessation attempt: Mediation through perceived barriers to cessation. *J Affect Disord*, 2015; 190:221–6. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26519643>

Tidey JW, Colby SM, and Xavier EM. Effects of smoking abstinence on cigarette craving, nicotine withdrawal, and nicotine reinforcement in smokers with and without schizophrenia. *Nicotine & Tobacco Research*, 2014; 16(3):326–34. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24113929>

Short NA, Raines AM, Oglesby ME, Zvolensky MJ, and Schmidt NB. Insomnia and emotion dysregulation: Independent and interactive associations with posttraumatic stress symptoms among trauma-exposed smokers. *J Affect Disord*, 2014; 165:159–65. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24882194>

Ahnallen C and Tidey J. Personalized smoking environment cue reactivity in smokers with schizophrenia and controls: A pilot study. *Psychiatry Research*, 2011; 188(2):286–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21550670>

Ratschen E, Britton J, Doody G, and McNeill A. Smoking attitudes, behaviour and nicotine dependence among mental health acute inpatients: An exploratory study. *The International Journal of Social Psychiatry*, 2010; 56(2):107–18. Available from: <http://isp.sagepub.com/cgi/reprint/56/2/107>

Radek R, Kohlhaas K, Rueter L, and Mohler E. Treating the cognitive deficits of schizophrenia with alpha4beta2 neuronal nicotinic receptor agonists. *Current Pharmaceutical Design*, 2010; 16(3):309–22. Available from: <http://www.benthamdirect.org/pages/content.php?CPD/2010/00000016/00000003/0008B.SGM>

Piper M, Smith S, Schlam T, Fleming M, Bittrich A, et al. Psychiatric disorders in smokers seeking treatment for tobacco dependence: Relations with tobacco dependence and cessation. *Journal of Consulting and Clinical Psychology*, 2010; 78(1):13–23. Available from: <http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2010-00910-005&CFID=27951098&CFTOKEN=69967056>

MacPherson L, Tull M, Matusiewicz A, Rodman S, Strong D, et al. Randomized controlled trial of behavioral activation smoking cessation treatment for smokers with elevated depressive symptoms. *Journal of Consulting and Clinical Psychology*, 2010; 78(1):55–61. Available from: <http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2010-00910-009&CFID=27962030&CFTOKEN=19031561>

Lawrence D and Mitrou F. Common mental disorders: Missed opportunity for smoking prevention and cessation. *The Australian and New Zealand Journal of Psychiatry*, 2010; 44(1):96–7. Available from: <http://informahealthcare.com/doi/full/10.3109/00048670903393712>

Cole M, Trigoboff E, Demler T, and Opler L. Impact of smoking cessation on psychiatric inpatients treated with clozapine or olanzapine. *Journal of Psychiatric Practice*, 2010; 16(2):75–81. Available from: <http://journals.lww.com/pages/login.aspx?ReturnUrl=%2fpracticalpsychiatry%2fsecure%2fpages%2fpurchase.aspx%3fan%3d00131746-201003000-00002>

Weinberger A, Desai R, and McKee S. Nicotine withdrawal in u.S. Smokers with current mood, anxiety, alcohol use, and substance use disorders. *Drug and Alcohol Dependence*, 2009; 108(1–2):7–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20006451>

Vickers K, Patten C, Lewis B, Clark M, Ussher M, et al. Feasibility of an exercise counseling intervention for depressed women smokers. *Nicotine & Tobacco Research*, 2009; 11(8):985–95. Available from: <http://ntr.oxfordjournals.org/cgi/content/abstract/11/8/985>

Schroeder SA. Antidepressant treatment and smoking cessation in bipolar disorder—reply. *Journal of the American Medical Association*, 2009; 301(20):2093. Available from: <http://jama.ama-assn.org/cgi/content/full/301/20/2093-a>

Schroeder SA. Moving forward in smoking cessation: Issues for psychiatric nurses Journal of the American Psychiatric Nurses Association, 2009; 15(1):68–72. Available from: <http://jap.sagepub.com/cgi/reprint/15/1/68>

Schroeder S. A 51-year-old woman with bipolar disorder who wants to quit smoking. Journal of the American Medical Association, 2009; 301(5):522–31. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19126801

Sarna L, Bialous SA, Wells MJ, and Kotlerman J. Smoking among psychiatric nurses: Does it hinder tobacco dependence treatment? Journal of the American Psychiatric Nurses Association, 2009; 15(1):59–67. Available from: <http://jap.sagepub.com/cgi/reprint/15/1/59>

Sacco K, Creedon C, Reutenauer E, Vessicchio J, Weinberger A, et al. Effects of atomoxetine on cognitive function and cigarette smoking in schizophrenia. Schizophrenia Research, 2009; 107(2–3):332–3. Available from: http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6TC2-4TVY5GJ-2&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&view=c&_acct=C000050221&_version=1&_urlVersion=0&_userid=10&md5=376a94e7653595579f6e1fa8dcf5c394

Prochaska J, Hall S, and Hall S. Stage-tailored tobacco cessation treatment in inpatient psychiatry. Psychiatric Services (Washington, D.C), 2009; 60(6):848. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19487360

Aubin HJ. Management of emergent psychiatric symptoms during smoking cessation. Current Medical Research and Opinion, 2009; 25(2):519–25. Available from: <http://www.informapharmascience.com/doi/abs/10.1185/03007990802707600>

Ziedonis D, Hitsman B, Beckham J, Zvolensky M, Adler L, et al. Tobacco use and cessation in psychiatric disorders: National institute of mental health report. Nicotine & Tobacco Research, 2008; 10(12):1691–715. Available from: <http://www.informaworld.com/smpp/content~db=all?content=10.1080/14622200802443569>

Prochaska J, Hall S, Tsoh J, Eisendrath S, Rossi J, et al. Treating tobacco dependence in clinically depressed smokers: Effect of smoking cessation on mental health functioning. American Journal of Public Health, 2008; 98(3):446–8. Available from: <http://www.ajph.org/cgi/reprint/98/3/446>

Hitsman B, Borrelli B, McChargue DE, Spring B, and Niaura R. History of depression and smoking cessation outcome: A meta-analysis. Journal of Consulting and Clinical Psychology, 2003; 71(4):657–63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12924670>

9A.3.5 Interventions for reducing smoking

Leutwyler, H, Hubbard, E, Bussell, T, Balestra, D, Cooper, B, Souza, RB, & Humfleet, G. (2024). A Pilot Randomized Controlled Trial of a Multicomponent Smoking Cessation Intervention for Adults with Serious Mental Illness. *Games Health J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39587943>

- Lee, S, Li, M, Le, GH, Teopiz, KM, Vinberg, M, Ho, R et al. (2024). Glucagon-like peptide-1 receptor agonists (GLP-1RAs) as treatment for nicotine cessation in psychiatric populations: a systematic review. *Ann Gen Psychiatry*, 23(1), 45. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39529123>
- Abufarsakh, B, Seng, S, Cline, MS, Scarduzio, J, & Okoli, CTC. (2024). Examining the Receptivity to Tobacco Treatment Information for Individuals Living With Mental Illnesses: A Mixed Method Study. *Int J Ment Health Nurs*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39381847>
- Huddleston, L, Shoesmith, E, Pervin, J, Stevens, R, Gilbody, S, & Ratschen, E. (2024). Effectiveness and experience of implementing digital interventions to promote smoking cessation among adults with severe mental illness: A systematic review and meta-analysis. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39382401>
- Lee, B, Rosen, LV, Mulroy, NM, Qian, Y, Shebl, FM, Becker, JE et al. (2024). Disparities in tobacco smoking and risk of cardiovascular disease in people with low socioeconomic status or serious psychological distress: A simulation analysis. *Am J Prev Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39419234>
- Okoli, CTC, Abufarsakh, B, Seng, S, Xie, W, McGovern, C, Arrows, T et al. (2024). A Pilot Feasibility Study of Delivering a Quit & Win Tobacco-Free Contest in Community Mental Health Programs. *J Am Psychiatr Nurses Assoc*, 10783903241281074. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39367784>
- Pleaver, S, Kisely, SR, Bonevski, B, Siskind, D, Guillaumier, A, McCarter, K, & Gartner, C E. (2024). Interventions for smoking cessation in inpatient psychiatry settings. *Cochrane Database Syst Rev*, 9(9), CD015934. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39229858>
- Talukder, S, Lappin, JM, Boland, VC, Weaver, N, McRobbie, H, & Courtney, RJ. (2024). Receipt of the 5As intervention for smoking cessation among people with and without mental health disorders. *J Psychiatr Res*, 179, 1-7. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39213719>
- Smith, RW, Holt-Lunstad, J, & Kawachi, I. (2024). Response to "Letter re: Benchmarking social isolation, loneliness and smoking: challenges and opportunities for public health.". *Am J Epidemiol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38775293>
- VanderWeele, TJ, & Kim, ES. (2024). Letter re: Benchmarking social isolation, loneliness, and smoking: challenges and opportunities for public health. *Am J Epidemiol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38702997>
- Castaldelli-Maia, JM, Camargos de Oliveira, V, Irber, FM, Blaas, IK, Angerville, B, Sousa Martins-da-Silva, A et al. (2023). Psychopharmacology of smoking cessation medications: focus on patients with mental health disorders. *Int Rev Psychiatry*, 35(5-6), 397-417. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38299651>
- Lima, DR, Davanzo, LC, Carvalho, CFC, Guimaraes-Pereira, BBS, Loreto, AR et al. (2023). Is self-reported 7-day point prevalence abstinence an useful measure for smoking cessation effectiveness among individuals with mental disorders? *Int Rev Psychiatry*, 35(5-6), 468-474. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38299649>

- Farris, SG, Smith, JE, Steinberg, DR, Altman, BR, Lambert-Messerlian, GM, Dunsiger, SI et al. (2023). Methodological approach for an integrated female-specific study of anxiety and smoking comorbidity. *Front Psychiatry*, 14, 1267753. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38076702>
- Davis, A. (2023). Pharmacotherapy for Treatment of Tobacco Use Disorder. *J Psychosoc Nurs Ment Health Serv*, 61(11), 6-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37909890>
- Swong, S, Nicholson, A, Smelson, D, Rogers, ES, El-Shahawy, O, & Sherman, SE. (2023). The effectiveness of a telephone smoking cessation program in mental health clinic patients by level of mental well-being and functioning: a secondary data analysis of a randomized clinical trial. *BMC Public Health*, 23(1), 2190. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37936218>
- Swong, S, Nicholson, A, Smelson, D, Rogers, ES, El-Shahawy, O, & Sherman, SE. (2023). The Effectiveness of a Telephone Smoking Cessation Program in Mental Health Clinic Patients by Level of Mental Well-Being and Functioning: A Secondary Data Analysis of a Randomized Clinical Trial. *Res Sq*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37674733>
- Tavakoli-Ardakani, M, Gholamzadeh Sani, Z, Beyraghi, N, Najarimoghadam, S, & Kheradmand, A. (2023). Comparison between cytisine and Nicotine Replacement Therapy in smoking cessation among inpatient psychiatric patients. *J Addict Dis*, 1-8. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37565489>
- Bennett, ME, Graham, AL, & Dickerson, F. (2023). Connecting Hospitalized Tobacco Smokers With Serious Mental Illness to a Digital Cessation Intervention. *Psychiatr Serv*, appips23074017. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37731347>
- Vojjala, M, Wysota, CN, Oketunbi, O, King, Q, & Rogers, ES. (2023). Integrating the "Quit and Stay Quit Monday" Model into Smoking Cessation Services for Smokers with Mental Health Conditions: A Pilot Randomized Controlled Trial. *J Smok Cessat*, 2023, 8165232. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37521160>
- Wojnar, M, Wierzbinski, P, Samochowiec, J, Rymaszewska, J, Filipiak, KJ, Wichniak, A et al. (2023). Management of nicotine dependence in patients with psychiatric disorders - recommendations of the Polish Psychiatric Association - part I. *Psychiatr Pol*, 1-16. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37370218>
- Daumit, GL, Evins, AE, Cather, C, Dalcin, AT, Dickerson, FB, Miller, ER et al. (2023). Effect of a Tobacco Cessation Intervention Incorporating Weight Management for Adults With Serious Mental Illness: A Randomized Clinical Trial. *JAMA Psychiatry*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37378972>
- Ajnakina, O, & Steptoe, A. (2023). The shared genetic architecture of smoking behaviours and psychiatric disorders: evidence from a population-based longitudinal study in England. *BMC Genom Data*, 24(1), 31. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37254052>
- Dickerson, F, Goldsholl, S, Yuan, CT, Dalcin, A, Eidman, B, Minahan, E et al. (2023). Promoting Evidence-Based Tobacco Cessation Treatment in Community Mental Health Clinics: Protocol for a

Prepost Intervention Study. *JMIR Res Protoc*, 12, e44787. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37171851>

Rubenstein, D, Pacek, LR, Smith, C, McClernon, FJ, Enyioha, C, & Vilardaga, R. (2023). Stagnant daily smoking prevalence between 2008 and 2019 among Black and Hispanic adults with serious psychological distress. *Drug Alcohol Depend*, 248, 109943. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37247521>

Nagawa, CS, Lane, IA, Davis, M, Wang, B, Pbert, L, Lemon, SC, & Sadasivam, RS. (2022). Experiences Using Family or Peer Support for Smoking Cessation and Considerations for Support Interventions: A Qualitative Study in Persons With Mental Health Conditions. *J Dual Diagn*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36576889>

Lodge, S, Bartlem, K, Gibson, L, Fehily, C, Bradley, T, McKeon, E et al. (2022). Characteristics and service use of NSW Quitline callers with and without mental health conditions. [MS Top Pick]. *Front Psychiatry*, 13, 868084. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36545036>

White, SA, Stone, E, Murphy, KA, Daumit, GL, & McGinty, EE. (2022). Smoking Cessation Medication Prescribing for Smokers With and Without Mental Illness. [MS Top Pick]. *Psychiatr Serv*, appips202100690. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36349496>

Morar, T, & Robertson, L. (2022). Smoking cessation among people with mental illness: A South African perspective. *S Afr Fam Pract (2004)*, 64(1), e1-e9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36073100>

McBrayer, K, Ouyang, F, Adams, Z, Hulvershorn, L, & Aalsma, MC. (2022). Rates of Tobacco Use Disorder, Pharmacologic Treatment, and Associated Mental Health Disorders in a Medicaid Claim Review Among Youth in Indiana, USA. *Tob Use Insights*, 15, 1179173X221119133. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36052177>

Nagawa, CS, Wang, B, Davis, M, Pbert, L, Cutrona, SL, Lemon, SC, & Sadasivam, RS. (2022). Examining pathways between family or peer factors and smoking cessation in a nationally representative US sample of adults with mental health conditions who smoke: a structural equation analysis. *BMC Public Health*, 22(1), 1566. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35978318>

Hart, JT, Boeckman, LM, & Beebe, LA. (2022). Unique cessation tools in the box: Quitline utilization and effectiveness trends among a large sample of tobacco users reporting mental health disorders. *Front Psychiatry*, 13, 869802. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35928774>

-Gonzalez, IL, Rodriguez-Donate, MC, & Guirao-Perez, G. (2022). Tobacco Consumption and Mental Health in the Canary Islands: A Quantitative Analysis. *Front Public Health*, 10, 959442. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35899167>

Baker, AL, McCarter, K, Brophy, L, Castle, D, Kelly, P J, Cocks, N et al. (2022). Adapting Peer Researcher Facilitated Strategies to Recruit People Receiving Mental Health Services to a Tobacco Treatment Trial. *Front Psychiatry*, 13, 869169. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35722563>

- Gonzalez-Roz, A, & Secades-Villa, R. (2022). Contingency management for smokers with mental health disorders and smoking-sensitive conditions caused or exacerbated by tobacco use: A review of existing studies, intervention parameters, and research priorities. *Exp Clin Psychopharmacol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35737555>
- Mahase, E. (2022). Government tobacco plan must tackle link between mental health and smoking, say psychiatrists. *BMJ*, 377, o1521. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35728820>
- Sanford, BT, Toll, BA, Palmer, AM, Foster, MG, Cummings, KM, Stansell, S& Rojewski, AM. (2022). Tobacco Treatment Outcomes for Hospital Patients With and Without Mental Health Diagnoses. *Front Psychiatry*, 13, 853001. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35722554>
- Schottl, SE, Niedermeier, M, Kopp-Wilfling, P, Fruhauf, A, Bichler, CS, Edlinger, M et al. (2022). Add-on exercise interventions for smoking cessation in people with mental illness: a systematic review and meta-analysis. *BMC Sports Sci Med Rehabil*, 14(1), 115. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35729669>
- Spanakis P, Peckham E, Young B, Heron P, Bailey D, et al. A systematic review of behavioural smoking cessation interventions for people with severe mental ill health-what works? *Addiction*, 2022; 117(6):1526–42. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34697848>
- Okoli CTC, Otachi JK, Seng S, Abufarsakh B, and Williams LB. Evaluating simulation-based tobacco treatment scenarios for providers delivering treatment for people living with mental illnesses. *Front Psychiatry*, 2022; 13:868550. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35463520>
- Jackson MA, Baker AL, Gould GS, Brown AL, Dunlop AJ, et al. Smoking cessation interventions for pregnant women attending treatment for substance use disorders: A systematic review. *Addiction*, 2022; 117(4):847–60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34374145>
- Hogan TH, Quisenberry A, Breitborde N, Moe A, and Ferketich A. An evaluation of the feasibility of implementing a novel tobacco dependence treatment program for high-risk individuals into clinical practice within a community mental health center. *Int J Ment Health Syst*, 2022; 16(1):15. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35184758>
- El-Sherbiny NA and Elsary AY. Smoking and nicotine dependence in relation to depression, anxiety, and stress in egyptian adults: A cross-sectional study. *J Family Community Med*, 2022; 29(1):8-16. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35197723>
- DeAtley T, Sokolovsky AW, Snell ML, and Tidey J. Mediation pathways of tobacco use among adult daily smokers with psychiatric symptoms in the population assessment of tobacco and health (path) survey. *Addictive Behaviors*, 2022; 129:107249. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35104739>
- Talukder SR, Lappin JM, Boland V, McRobbie H, and Courtney RJ. Inequity in smoking cessation clinical trials testing pharmacotherapies: Exclusion of smokers with mental health disorders. *Tobacco Control*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34862325>

Sharma-Kumar R, Puljevic C, Morphet K, Meurk C, and Gartner C. The acceptability and effectiveness of videos promoting smoking cessation among australians experiencing mental illness. *Health Educ Behav*, 2021;10901981211034738. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34496656>

Schnitzer K, Maravic MC, Arntz D, Phillips NL, Pachas G, et al. Integrated smoking cessation for smokers with serious mental illness: Protocol for a convergent mixed methods implementation evaluation study. *JMIR Res Protoc*, 2021; 10(7):e25390. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34313603>

Li Y, Zhang C, Ding S, Li J, Li L, et al. Physical activity, smoking, alcohol consumption and depressive symptoms among young, early mature and late mature people: A cross-sectional study of 76,223 in China. *J Affect Disord*, 2021; 299:60-6. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34822919>

Hernandez-Perez A, Vargas-Nunez I, Moreno-Jimenez B, Perez-Padilla R, and Ramirez-Venegas A. Affective comorbidity associated with symptoms, lung function, and differences between patients with copd for biomass and tobacco smoke exposure. *J Clin Psychol Med Settings*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34618283>

Hawes MR, Roth KB, and Cabassa LJ. Systematic review of psychosocial smoking cessation interventions for people with serious mental illness. *Journal of Dual Diagnosis*, 2021; 17(3):216–35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34281493>

Guo K, Li J, Li J, Chen N, Li Y, et al. The effects of pharmacological interventions on smoking cessation in people with alcohol dependence: A systematic review and meta-analysis of nine randomized controlled trials. *International Journal of Clinical Practice*, 2021; 75(11):e14594. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34228852>

Gilbody S, Peckham E, Bailey D, Arundel C, Heron P, et al. Smoking cessation in severe mental illness: Combined long-term quit rates from the UK scimitar trials programme. *The British Journal of Psychiatry*, 2021; 218(2):95–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31685048>

Gao N, Solomon P, Clay Z, and Swarbrick P. A pilot study of wellness coaching for smoking cessation among individuals with mental illnesses. *J Ment Health*, 2021:1-7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33999729>

Busch AM, Nederhoff DM, Dunsiger SI, Japuntich SJ, Chrastek M, et al. Chronic care treatment for smoking cessation in patients with serious mental illness: A pilot randomized trial. *BMC Psychiatry*, 2021; 21(1):104. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33593332>

Browne J, Halverson TF, and Vilardaga R. Erratum to: Engagement with a digital therapeutic for smoking cessation designed for persons with psychiatric illness fully mediates smoking outcomes in a pilot randomized controlled trial. *Transl Behav Med*, 2021. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34875095>

Brown E, O'Donoghue B, White SL, Chanen A, Bedi G, et al. Tobacco smoking in young people seeking treatment for mental ill-health: What are their attitudes, knowledge and behaviours towards

quitting? *Ir J Psychol Med*, 2021; 38(1):30–9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32317033>

Billingsley BE and Steinberg ML. Motivational interviewing produces change talk in smokers with serious mental illness. *Journal of Dual Diagnosis*, 2021:1-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33725471>

Anso-Diego G, Gonzalez-Roz A, Krotter A, Garcia-Perez A, and Secades-Villa R. Contingency management for smoking cessation among individuals with substance use disorders: In-treatment and post-treatment effects. *Addictive Behaviors*, 2021; 119:106920. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33798921>

Alghzawi HM and Storr CL. Gender differences in the interrelations among social support, stressful life events, and smoking cessation in people with severe mental illnesses. *Journal of the American Psychiatric Nurses Association*, 2021:10783903211008248. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33926296>

Vilardaga R, Rizo J, Palenski PE, Mannelli P, Oliver JA, et al. Erratum to: Pilot randomized controlled trial of a novel smoking cessation app designed for individuals with co-occurring tobacco dependence and serious mental illness. *Nicotine & Tobacco Research*, 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32761194>

Tinghino B, Lugoboni F, Amatulli A, Biasin C, Bramani Araldi M, et al. The fodrat study (food addiction, drugs, alcohol and tobacco): First data on food addiction prevalence among patients with addiction to drugs, tobacco and alcohol. *Eat Weight Disord*, 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32072572>

Taylor GMJ, Itani T, Thomas KH, Rai D, Jones T, et al. Prescribing prevalence, effectiveness, and mental health safety of smoking cessation medicines in patients with mental disorders. *Nicotine & Tobacco Research*, 2020; 22(1):48–57. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31289809>

Rogers ES, Wysota C, Prochaska JJ, Tenner C, Dognin J, et al. A behavioral economic intervention to increase psychiatrist adherence to tobacco treatment guidelines: A provider-randomized study protocol. *Implement Sci Commun*, 2020; 1. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32617528>

Myers MG, Strong DR, Chen TC, and Linke SE. Enhancing engagement in evidence-based tobacco cessation treatment for smokers with mental illness: A pilot randomized trial. *Journal of Substance Abuse Treatment*, 2020; 111:29-36. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32087836>

Lightfoot K, Panagiotaki G, and Nobes G. Effectiveness of psychological interventions for smoking cessation in adults with mental health problems: A systematic review. *British Journal of Health Psychology*, 2020; 25(3):615–38. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32678937>

Jimenez-Murcia S, Gimenez M, Granero R, Masuet-Aumatell C, Ramon JM, et al. Gambling disorder seeking treatment patients and tobacco use in relation to clinical profiles. *Addictive Behaviors*, 2020:106723. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33203595>

Japuntich SJ, Hammett PJ, Rogers ES, Fu S, Burgess DJ, et al. Effectiveness of proactive tobacco cessation outreach in smokers with serious mental illness. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31957794>

Jaen-Moreno MJ, Garcia-Portilla MP, Sarramea F, Bobes Bascaran T, Galvan G, et al. The multi-component smoking cessation support programme (mcscsp) is effective in patients with severe mental disorder without gender differences. *Adicciones*, 2020; 0(0):1481. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33338247>

DeAtley T, Denlinger-Apte RL, Cioe PA, Colby SM, Cassidy RN, et al. Biopsychosocial mechanisms associated with tobacco use in smokers with and without serious mental illness. *Preventive Medicine*, 2020:106190. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32622776>

Chan HY, Chen JJ, Pan YJ, and Hsu CC. Nicotine replacement therapy and healthy lifestyle psychoeducation for smoking reduction in acute psychiatric inpatients: A cluster-randomized parallel study. *J Clin Psychopharmacol*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32032137>

Bilevicius E, Single A, Baumgartner C, Bui V, Kempe T, et al. Developing and testing the effectiveness of a novel online integrated treatment for problem gambling and tobacco smoking: A protocol for an open-label randomized controlled trial. *Trials*, 2020; 21(1):937. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33213506>

Alghzawi H, Trinkoff A, Zhu S, and Storr C. Remission from nicotine dependence among people with severe mental illness who received help/services for tobacco/nicotine use. *Int J Methods Psychiatr Res*, 2020:e1845. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32945054>

Vilardaga R, Rizo J, Palenski P, Mannelli P, Oliver JA, et al. Pilot randomized controlled trial of a novel smoking cessation app designed for individuals with co-occurring tobacco dependence and serious mental illness. *Nicotine & Tobacco Research*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31667501>

Sweeney R, Moodie M, Baker AL, Borland R, Castle D, et al. Protocol for an economic evaluation of the quitlink randomized controlled trial for accessible smoking cessation support for people with severe mental illness. *Front Psychiatry*, 2019; 10:618. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31551827>

Steinberg ML, Weinberger AH, and Tidey JW. Non-pharmacological treatments for tobacco users with mental health symptoms. *Nicotine & Tobacco Research*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30768202>

Peckham E, Arundel C, Bailey D, Crosland S, Fairhurst C, et al. A bespoke smoking cessation service compared with treatment as usual for people with severe mental ill health: The scimitar+ rct. *Health*

Technology Assessment, 2019; 23(50):1–116. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31549622>

Olando Y, Kuria M, Mathai M, and Huffman MD. Efficacy of a group tobacco cessation behavioral intervention among tobacco users with concomitant mental illness in kenya: Protocol for a controlled clinical trial. BMC Public Health, 2019; 19(1):1700. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31852536>

Klein P, Lawn S, Tsourtos G, and van Agteren J. Tailoring of a smartphone smoking cessation app (kick.It) for serious mental illness populations: Qualitative study. JMIR Hum Factors, 2019; 6(3):e14023. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31482850>

Johnson AL, Kaye J, Baker TB, Fiore MC, Cook JW, et al. Psychiatric comorbidities in a comparative effectiveness smoking cessation trial: Relations with cessation success, treatment response, and relapse risk factors. Drug and Alcohol Dependence, 2019; 207:107796. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31864163>

Jaen-Moreno MJ, Feu N, Redondo-Ecija J, Montiel FJ, Gomez C, et al. Smoking cessation opportunities in severe mental illness (tobacco intensive motivational and estimate risk - timer-): Study protocol for a randomized controlled trial. Trials, 2019; 20(1):47. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30642365>

Heron P, McCloud T, Arundel C, Bailey D, Ker S, et al. Standard smoking cessation services in sites participating in the scimitar+ trial for people with severe mental ill health. BJPsych Bull, 2019:1-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31162002>

Hammett PJ, Lando HA, Erickson DJ, Widome R, Taylor BC, et al. Proactive outreach tobacco treatment for socioeconomically disadvantaged smokers with serious mental illness. Journal of Behavioral Medicine, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31363948>

Evins AE, Benowitz NL, West R, Russ C, McRae T, et al. Neuropsychiatric safety and efficacy of varenicline, bupropion, and nicotine patch in smokers with psychotic, anxiety, and mood disorders in the eagles trial. J Clin Psychopharmacol, 2019; 39(2):108–16. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30811371>

Cohn AM, Zhou Y, Cha S, Perreras L, and Graham AL. Treatment engagement mediates the links between symptoms of anxiety, depression, and alcohol use disorder with abstinence among smokers registered on an internet cessation program. Journal of Substance Abuse Treatment, 2019; 98:59-65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30665605>

Ballbe M, Martinez C, Feliu A, Torres N, Nieva G, et al. Effectiveness of a telephone-based intervention for smoking cessation in patients with severe mental disorders: Study protocol for a randomized controlled trial. Trials, 2019; 20(1):38. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30635072>

Vilardaga R, Rizo J, Ries RK, Kientz JA, Ziedonis DM, et al. Formative, multimethod case studies of learn to quit, an acceptance and commitment therapy smoking cessation app designed for people

with serious mental illness. *Transl Behav Med*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30445507>

Rasmussen M, Klinge M, Krogh J, Nordentoft M, and Tonnesen H. Effectiveness of the gold standard programme (gsp) for smoking cessation on smokers with and without a severe mental disorder: A danish cohort study. *BMJ Open*, 2018; 8(6):e021114. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29895653>

Lawn S, Van Agteren J, Zabeen S, Bertossa S, Barton C, et al. Adapting, pilot testing and evaluating the kick.It app to support smoking cessation for smokers with severe mental illness: A study protocol. *International Journal of Environmental Research and Public Health*, 2018; 15(2). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30720772>

Curtis J, Zhang C, McGuigan B, Pavel-Wood E, Morell R, et al. Y-quit: Smoking prevalence, engagement, and effectiveness of an individualized smoking cessation intervention in youth with severe mental illness. *Front Psychiatry*, 2018; 9:683. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30618864>

Clyde M, Pipe A, Els C, Reid R, Fu A, et al. Nicotine metabolite ratio and smoking outcomes using nicotine replacement therapy and varenicline among smokers with and without psychiatric illness. *J Psychopharmacol*, 2018:269881118773532. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29788791>

Castaldelli-Maia JM, Loreto AR, Guimaraes-Pereira BBS, Carvalho CFC, Gil F, et al. Smoking cessation treatment outcomes among people with and without mental and substance use disorders: An observational real-world study. *European Psychiatry*, 2018; 52:22-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29609056>

Bullen C, Verbiest M, Galea-Singer S, Kurdziel T, Laking G, et al. The effectiveness and safety of combining varenicline with nicotine e-cigarettes for smoking cessation in people with mental illnesses and addictions: Study protocol for a randomised-controlled trial. *BMC Public Health*, 2018; 18(1):596. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29728074>

Aschbrenner KA, Patten CA, and Brunette MF. Feasibility of a support person intervention to promote smoking cessation treatment use among smokers with mental illness. *Transl Behav Med*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29385555>

Trainor K and Leavey G. Barriers and facilitators to smoking cessation among people with severe mental illness: A critical appraisal of qualitative studies. *Nicotine & Tobacco Research*, 2017; 19(1):14–23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27613905>

Sharma R, Alla K, Pfeffer D, Meurk C, Ford P, et al. An appraisal of practice guidelines for smoking cessation in people with severe mental illness. *Australian and New Zealand Journal of Psychiatry*, 2017; 51(11):1106–20. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28859486>

Schwindt R, Hudmon KS, Knisely M, Davis L, and Pike C. Impact of tobacco quitlines on smoking cessation in persons with mental illness: A systematic review. *J Drug Educ*, 2017; 47(1-2):68-81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29534595>

- Rogers ES, Friedes R, Jakes A, Grossman E, Link A, et al. Long-term abstinence and predictors of tobacco treatment uptake among hospitalized smokers with serious mental illness enrolled in a smoking cessation trial. *Journal of Behavioral Medicine*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28349344>
- Metse AP, Wiggers JH, Wye PM, Wolfenden L, Prochaska JJ, et al. Smoking and mental illness: A bibliometric analysis of research output over time. *Nicotine & Tobacco Research*, 2017; 19(1):24-31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27980040>
- Knudsen HK. Implementation of smoking cessation treatment in substance use disorder treatment settings: A review. *American Journal of Drug and Alcohol Abuse*, 2017; 43(2):215–25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27314884>
- Brunette MF, Pratt SI, Bartels SJ, Scherer EA, Sigmon SC, et al. Randomized trial of interventions for smoking cessation among medicaid beneficiaries with mental illness. *Psychiatric Services*, 2017:appips201700245. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29137560>
- Brunette MF, Ferron JC, Robinson D, Coletti D, Geiger P, et al. Brief web-based interventions for young adult smokers with severe mental illnesses: A randomized, controlled pilot study. *Nicotine & Tobacco Research*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29059417>
- Zawertailo L. Safety of smoking cessation drugs for mentally ill patients. *Lancet*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27116919>
- Wu Q, Gilbody S, Peckham E, Brabyn S, and Parrott S. Varenicline for smoking cessation and reduction in people with severe mental illnesses: Systematic review and meta-analysis. *Addiction*, 2016; 111(9):1554–67. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27043328>
- Weinberger AH. Smoking cessation and adults with serious mental illness: The need for more research at every step of the quit process. *Nicotine & Tobacco Research*, 2016; 18(3):227–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26718906>
- Vilardaga R, Rizo J, Kientz JA, McDonell MG, Ries RK, et al. User experience evaluation of a smoking cessation app in people with serious mental illness. *Nicotine & Tobacco Research*, 2016; 18(5):1032-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26581430>
- Verplaetse TL, Smith PH, Pittman BP, Mazure CM, and McKee SA. Associations of gender, smoking, and stress with transitions in major depression diagnoses. *Yale J Biol Med*, 2016; 89(2):123–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27354839>
- Tzilos GK, Reddy MK, Caviness CM, Anderson BJ, and Stein MD. Mostly smokers: Identifying subtypes of emerging adult substance users. *Substance Use and Misuse*, 2016:1-6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27484392>
- Tong L, Shi HJ, Zhang Z, Yuan Y, Xia ZJ, et al. Mediating effect of anxiety and depression on the relationship between attention-deficit/hyperactivity disorder symptoms and smoking/drinking. *Sci Rep*, 2016; 6:21609. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26923609>

Tidey JW, Cassidy RN, and Miller ME. Smoking topography characteristics of very low nicotine content cigarettes, with and without nicotine replacement, in smokers with schizophrenia and controls. *Nicotine & Tobacco Research*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26995794>

Tedeschi GJ, Cummins SE, Anderson CM, Anthenelli RM, Zhuang YL, et al. Smokers with self-reported mental health conditions: A case for screening in the context of tobacco cessation services. *PLoS One*, 2016; 11(7):e0159127. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27391334>

Smits JA, Powers MB, Rosenfield D, Zvolensky MJ, Jacquart J, et al. Bdnf val66met polymorphism as a moderator of exercise enhancement of smoking cessation treatment in anxiety vulnerable adults. *Ment Health Phys Act*, 2016; 10:73–7. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27453731>

Smits JA, Kauffman BY, Lee-Furman E, Zvolensky MJ, Otto MW, et al. Enhancing panic and smoking reduction treatment with d-cycloserine: Study protocol for a randomized controlled trial. *Contemporary Clinical Trials*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27015966>

Smith RC, Amiaz R, Si TM, Maayan L, Jin H, et al. Varenicline effects on smoking, cognition, and psychiatric symptoms in schizophrenia: A double-blind randomized trial. *PLoS One*, 2016; 11(1):e0143490. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26730716>

Schmidt NB, Raines AM, Allan NP, and Zvolensky MJ. Anxiety sensitivity risk reduction in smokers: A randomized control trial examining effects on panic. *Behav Res Ther*, 2016; 77:138–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26752327>

Sankaranarayanan A, Clark V, Baker A, Palazzi K, Lewin TJ, et al. Reducing smoking reduces suicidality among individuals with psychosis: Complementary outcomes from a healthy lifestyles intervention study. *Psychiatry Res*, 2016; 243:407–12. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/27450743>

Rohsenow DJ, Martin RA, Tidey JW, Colby SM, and Monti PM. Treating smokers in substance treatment with contingent vouchers, nicotine replacement and brief advice adapted for sobriety settings. *Journal of Substance Abuse Treatment*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27658756>

Rogers ES, Smelson DA, Gillespie CC, Elbel B, Poole S, et al. Telephone smoking-cessation counseling for smokers in mental health clinics: A patient-randomized controlled trial. *American Journal of Preventive Medicine*, 2016; 50(4):518–27. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/26711163>

Reid HH and Ledgerwood DM. Depressive symptoms affect changes in nicotine withdrawal and smoking urges throughout smoking cessation treatment: Preliminary results. *Addict Res Theory*, 2016; 24(1):48-53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27547173>

Poulet E, Galvao F, Haffen E, Szekely D, Brault C, et al. Effects of smoking status and madrs retardation factor on response to low frequency repetitive transcranial magnetic stimulation for

depression. *European Psychiatry*, 2016; 38:40-4. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27657664>

Patten CA, Bronars CA, Vickers Douglas KS, Ussher MH, Levine JA, et al. Supervised, vigorous intensity exercise intervention for depressed female smokers: A pilot study. *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27613946>

Murphy KG and Flory K. Motives underlying smoking in college students with adhd. *American Journal of Drug and Alcohol Abuse*, 2016;1-10. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27494646>

Metse AP, Wiggers J, Wye P, Clancy R, Moore L, et al. Uptake of smoking cessation aids by smokers with a mental illness. *Journal of Behavioral Medicine*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27357297>

Mendelsohn CP and Wodak Am A. Smoking cessation in people with alcohol and other drug problems. *Australian Family Physician*, 2016; 45(8):569-73. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27610446>

Mathew AR, Hogarth L, Leventhal AM, Cook JW, and Hitsman B. Cigarette smoking and depression comorbidity: Systematic review & proposed theoretical model. *Addiction*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27628300>

Mathai AJ, Kanwar J, Okusaga O, Fuchs D, Lowry CA, et al. Blood levels of monoamine precursors and smoking in patients with schizophrenia. *Front Public Health*, 2016; 4:182. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27626030>

Lembke A and Humphreys K. A call to include people with mental illness and substance use disorders alongside 'regular' smokers in smoking cessation research. *Tobacco Control*, 2016; 25(3):261–2. Available from:

<http://tobaccocontrol.bmj.com/content/25/3/261.abstract>

Knowles S, Planner C, Bradshaw T, Peckham E, Man MS, et al. Making the journey with me: A qualitative study of experiences of a bespoke mental health smoking cessation intervention for service users with serious mental illness. *BMC Psychiatry*, 2016; 16:193. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27278101>

Khanna P, Clifton AV, Banks D, and Tosh GE. Smoking cessation advice for people with serious mental illness. *Cochrane Database of Systematic Reviews*, 2016; 1:CD009704. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26816385>

Jeon DW, Shim JC, Kong BG, Moon JJ, Seo YS, et al. Adjunctive varenicline treatment for smoking reduction in patients with schizophrenia: A randomized double-blind placebo-controlled trial. *Schizophr Res*, 2016; 176(2-3):206–11. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/27543252>

Garey L, Cheema MK, Otal TK, Schmidt NB, Neighbors C, et al. The sequential pathway between trauma-related symptom severity and cognitive-based smoking processes through perceived stress and negative affect reduction expectancies among trauma exposed smokers. *American Journal on Addictions*, 2016; 25(7):565-72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27658236>

Garey L, Bakhshaie J, Vujanovic AA, Reitzel LR, Schmidt NB, et al. Posttraumatic stress symptom severity and cognitive-based smoking processes among trauma-exposed treatment-seeking smokers: The role of perceived stress. *Addictive Behaviors*, 2016; 60:84–9. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27100473>

Garcia-Portilla MP, Garcia-Alvarez L, Sarramea F, Galvan G, Diaz-Mesa E, et al. It is feasible and effective to help patients with severe mental disorders to quit smoking: An ecological pragmatic clinical trial with transdermal nicotine patches and varenicline. *Schizophr Res*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27237598>

Ferron JC, Devitt T, McHugo GJ, J AJ, Cook JA, et al. Abstinence and use of community-based cessation treatment after a motivational intervention among smokers with severe mental illness. *Community Ment Health J*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26932324>

Dickerson FB, Savage CL, Schweinfurth LA, Medoff DR, Goldberg RW, et al. The use of peer mentors to enhance a smoking cessation intervention for persons with serious mental illnesses. *Psychiatric Rehabilitation Journal*, 2016; 39(1):5-13. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/26461436>

Dennis PA, Kimbrel NA, Dedert EA, Beckham JC, Dennis MF, et al. Supplemental nicotine preloading for smoking cessation in posttraumatic stress disorder: Results from a randomized controlled trial. *Addictive Behaviors*, 2016; 59:24–9. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27046670>

Deng H, Wang J, Zhang X, Ma M, Domingo C, et al. Smoking reduction and quality of life in chronic patients with schizophrenia in a chinese population: A pilot study. *American Journal on Addictions*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26824336>

Dedert EA, Resick PA, McFall ME, Dennis PA, Olsen M, et al. Pilot cases of combined cognitive processing therapy and smoking cessation for smokers with posttraumatic stress disorder. *Behavior Therapy*, 2016; 47(1):54–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26763497>

Dahne J, Murphy JG, and MacPherson L. Depressive symptoms and cigarette demand as a function of induced stress. *Nicotine & Tobacco Research*, 2016. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27245238>

Brody AL, Zorick T, Hubert R, Helleman GS, Balali S, et al. Combination extended smoking cessation treatment plus home visits for smokers with schizophrenia: A randomized controlled trial. *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27613888>

Akimoto H, Oshima S, Negishi A, Ohara K, Ohshima S, et al. Assessment of the risk of suicide-related events induced by concomitant use of antidepressants in cases of smoking cessation treatment with varenicline and assessment of latent risk by the use of varenicline. *PLoS One*, 2016; 11(9):e0163583. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/27657721>

Quit smoking help for people with mental health problems. *Nursing Standard*, 2016; 30(34):10.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27097176>

Martinez C, Guydish J, Le T, Tajima B, and Passalacqua E. Predictors of quit attempts among smokers enrolled in substance abuse treatment. *Addictive Behaviors*, 2015; 40:1–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25218064>

Covey LS, Hu MC, Winhusen T, Lima J, Berlin I, et al. Anxiety and depressed mood decline following smoking abstinence in adult smokers with attention deficit hyperactivity disorder. *Journal of Substance Abuse Treatment*, 2015; 59:104–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26272693>

Wu BJ and Lan TH. Predictors of smoking reduction outcomes in a sample of 287 patients with schizophrenia spectrum disorders. *European Archives of Psychiatry and Clinical Neuroscience*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26310877>

Tidey JW and Miller ME. Smoking cessation and reduction in people with chronic mental illness. *British Medical Journal*, 2015; 351:h4065. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26391240>

Stubbs B, Vancampfort D, Bobes J, De Hert M, and Mitchell AJ. How can we promote smoking cessation in people with schizophrenia in practice? A clinical overview. *Acta Psychiatrica Scandinavica*, 2015; 132(2):122–30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25754402>

Steinberg ML, Williams JM, Stahl NF, Budsock PD, and Cooperman NA. An adaptation of motivational interviewing increases quit attempts in smokers with serious mental illness. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25744954>

Secades-Villa R, Vallejo-Seco G, Garcia-Rodriguez O, Lopez-Nunez C, Weidberg S, et al. Contingency management for cigarette smokers with depressive symptoms. *Experimental and Clinical Psychopharmacology*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26280589>

Roberts E, Evins AE, McNeill A, and Robson D. Efficacy and acceptability of pharmacotherapy for smoking cessation in adults with serious mental illness: A systematic review and network meta-analysis. *Addiction*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26594837>

Prochaska JJ, Fromont SC, Ramo DE, Young-Wolff KC, Delucchi K, et al. Gender differences in a randomized controlled trial treating tobacco use among adolescents and young adults with mental health concerns. *Nicotine & Tobacco Research*, 2015; 17(4):479–85. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25762759>

Peckham E, Man MS, Mitchell N, Li J, Becque T, et al. Smoking cessation intervention for severe mental ill health trial (scimitar): A pilot randomised control trial of the clinical effectiveness and cost-effectiveness of a bespoke smoking cessation service. *Health Technology Assessment*, 2015; 19(25):1–148. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25827850>

O'Brien B, Knight-West O, Walker N, Parag V, and Bullen C. E-cigarettes versus nrt for smoking reduction or cessation in people with mental illness: Secondary analysis of data from the ascend trial. *Tobacco Induced Diseases*, 2015; 13(1):5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25814920>

- Michopoulos I, Rizos E, Gournellis R, Karvouni A, Kotsioumpa I, et al. Smoking reduction in psychiatric inpatients is feasible: Results from a 12-month prospective study. *Annals of General Psychiatry*, 2015; 14:4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25685172>
- Matsuda A, Hasegawa J, Wang X, Tsuno S, and Miura N. Psychological characteristics in patients during treatment for tobacco dependence. *Yonago Acta Med*, 2015; 58(2):81–4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26306058>
- Ker S and Reilly J. From efficacy to effectiveness in smoking cessation in severe mental illness. *British Medical Journal*, 2015; 351:h5681. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26507687>
- Kelly MM, Jensen KP, and Sofuoglu M. Co-occurring tobacco use and posttraumatic stress disorder: Smoking cessation treatment implications. *American Journal on Addictions*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26584242>
- Heffner JL, McClure JB, Mull KE, Anthenelli RM, and Bricker JB. Acceptance and commitment therapy and nicotine patch for smokers with bipolar disorder: Preliminary evaluation of in-person and telephone-delivered treatment. *Bipolar Disorders*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25912192>
- Gilbody S, Peckham E, Man MS, Mitchell N, Li J, et al. Bespoke smoking cessation for people with severe mental ill health (scimitar): A pilot randomised controlled trial. *Lancet Psychiatry*, 2015; 2(5):395–402. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26360283>
- Gaalema DE, Miller ME, and Tidey JW. Predicted impact of nicotine reduction on smokers with affective disorders. *Tob Regul Sci*, 2015; 1(2):154–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26236765>
- Foster DW, Schmidt NB, and Zvolensky MJ. Influences of barriers to cessation and reasons for quitting on substance use among treatment-seeking smokers who report heavy drinking. *Journal of Addiction Research & Therapy*, 2015; 6(4). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26949566>
- Farris SG, Allan NP, Morales PC, Schmidt NB, and Zvolensky MJ. Does successful smoking cessation reduce anxious arousal among treatment-seeking smokers? *J Anxiety Disord*, 2015; 36:92–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26460537>
- Evins AE, Cather C, and Laffer A. Treatment of tobacco use disorders in smokers with serious mental illness: Toward clinical best practices. *Harv Rev Psychiatry*, 2015; 23(2):90–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25747922>
- Evins AE and Cather C. Effective cessation strategies for smokers with schizophrenia. *Int Rev Neurobiol*, 2015; 124:133–47. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26472528>
- Dickerson F, Savage CL, Schweinfurth LA, Goldberg RW, Bennett M, et al. The experience of peer mentors in an intervention to promote smoking cessation in persons with psychiatric illness. *Community Ment Health J*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26602772>

Clyde M, Tulloch H, Reid R, Els C, and Pipe A. Task and barrier self-efficacy among treatment-seeking smokers with current, past or no psychiatric diagnosis. *Addictive Behaviors*, 2015; 46:65–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25813271>

Bernard P, Ninot G, Cyprien F, Courtet P, Guillaume S, et al. Exercise and counseling for smoking cessation in smokers with depressive symptoms: A randomized controlled pilot trial. *Journal of Dual Diagnosis*, 2015; 11(3-4):205–16. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26683252>

Bennett ME, Brown CH, Li L, Himelhoch S, Bellack A, et al. Smoking cessation in individuals with serious mental illness: A randomized controlled trial of two psychosocial interventions. *Journal of Dual Diagnosis*, 2015:0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26457385>

Barnett PG, Wong W, Jeffers A, Hall SM, and Prochaska JJ. Cost-effectiveness of smoking cessation treatment initiated during psychiatric hospitalization: Analysis from a randomized, controlled trial. *Journal of Clinical Psychiatry*, 2015; 76(10):e1285–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26528651>

Barnett PG, Jeffers A, Smith MW, Chow BK, McFall M, et al. Cost-effectiveness of integrating tobacco cessation into post-traumatic stress disorder treatment. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25943761>

Baker HM, Ranney LM, and Goldstein AO. Pilot implementation of a wellness and tobacco cessation curriculum in north carolina group homes. *Community Ment Health J*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26711097>

Baker AL, Richmond R, Kay-Lambkin FJ, Filia SL, Castle D, et al. Randomized controlled trial of a healthy lifestyle intervention among smokers with psychotic disorders. *Nicotine & Tobacco Research*, 2015; 17(8):946–54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25744962>

Ashton M, Rigby A, and Galletly C. Evaluation of a community-based smoking cessation programme for people with severe mental illness. *Tobacco Control*, 2015; 24(3):275–80. Available from: <http://tobaccocontrol.bmj.com/content/24/3/275.abstract>

Aschbrenner KA, Brunette MF, McElvery R, Naslund JA, Scherer EA, et al. Cigarette smoking and interest in quitting among overweight and obese adults with serious mental illness enrolled in a fitness intervention. *J Nerv Ment Dis*, 2015; 203(6):473–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26034872>

Annamalai A, Singh N, and O'Malley SS. Smoking use and cessation among people with serious mental illness. *Yale J Biol Med*, 2015; 88(3):271–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26339210>

Chengappa KN, Perkins KA, Brar JS, Schlicht PJ, Turkin SR, et al. Varenicline for smoking cessation in bipolar disorder: A randomized, double-blind, placebo-controlled study. *Journal of Clinical Psychiatry*, 2014; 75(7):765–72. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25006684>

Drummond MB, Astemborski J, Lambert AA, Goldberg S, Stitzer ML, et al. A randomized study of contingency management and spirometric lung age for motivating smoking cessation among

injection drug users. BMC Public Health, 2014; 14:761. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25074396>

Tantirangsee N, Assanangkornchai S, and Marsden J. Effects of a brief intervention for substance use on tobacco smoking and family relationship functioning in schizophrenia and related psychoses: A randomised controlled trial. Journal of Substance Abuse Treatment, 2014. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25468004>

Strong DR, Uebelacker L, Fokas K, Saritelli J, Matsko S, et al. Utilization of evidence-based smoking cessation treatments by psychiatric inpatient smokers with depression. Journal of Addiction Medicine, 2014; 8(2):77–83. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24562400>

Stockings EA, Bowman JA, Baker AL, Terry M, Clancy R, et al. Impact of a postdischarge smoking cessation intervention for smokers admitted to an inpatient psychiatric facility: A randomized controlled trial. Nicotine & Tobacco Research, 2014; 16(11):1417–28. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24939916>

Schoenfelder EN, Faraone SV, and Kollins SH. Stimulant treatment of adhd and cigarette smoking: A meta-analysis. Pediatrics, 2014; 133(6):1070–80. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24819571>

Ruther T, Bobes J, De Hert M, Svensson TH, Mann K, et al. EPA guidance on tobacco dependence and strategies for smoking cessation in people with mental illness. European Psychiatry, 2014; 29(2):65–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24485753>

Rogers ES, Fu SS, Krebs P, Noorbaloochi S, Nugent SM, et al. Proactive outreach for smokers using vha mental health clinics: Protocol for a patient-randomized clinical trial. BMC Public Health, 2014; 14(1):1294. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25518878>

Metse AP, Bowman JA, Wye P, Stockings E, Adams M, et al. Evaluating the efficacy of an integrated smoking cessation intervention for mental health patients: Study protocol for a randomised controlled trial. Trials, 2014; 15:266. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24996596>

Mee S. Self-efficacy: A mediator of smoking behavior and depression among college students.

Pediatr Nurs, 2014; 40(1):9–15, 37. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24757914>

McDonell M, McPherson S, Vilardaga R, Srebnik D, Angelo FN, et al. Preliminary findings: Contingency management targeting psycho-stimulant use results in secondary decreases in smoking for severely mentally ill adults. American Journal on Addictions, 2014; 23(4):407–10. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24961363>

Leventhal A. Review: Psychosocial mood management improves smoking cessation in people with current and past depression. Evid Based Ment Health, 2014; 17(2):e3. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24591546>

Bittoun R, Barone M, Mendelsohn CP, Elcombe EL, and Glozier N. Promoting positive attitudes of tobacco-dependent mental health patients towards nrt-supported harm reduction and smoking

cessation. Australian and New Zealand Journal of Psychiatry, 2014; 48(10):954–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24907132>

Baiardini I, Sorino C, Di Marco F, and Facchini F. Smoking cessation, anxiety, mood and quality of life: Reassuring evidences. Minerva Med, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25396688>

Vujanovic AA, Farris SG, Harte CB, Smits JA, and Zvolensky MJ. Smoking status and exercise in relation to ptsd symptoms: A test among trauma-exposed adults. Ment Health Phys Act, 2013; 6(2). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24273598>

DeHay T, Morris C, May MG, Devine K, and Waxmonsky J. Tobacco use in youth with mental illnesses. Journal of Behavioral Medicine, 2012; 35(2):139–48. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21479646>

Adams CE, Baillie LE, and Copeland AL. The smoking-related weight and eating episodes test (sweet): Development and preliminary validation. Nicotine & Tobacco Research, 2011; 13(11):1123–31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21849410>

Gonzalez-Pinto A, Alberich S, Ruiz de Azua S, Martinez-Cengotitabengoa M, Fernandez M, et al. Psychosis and smoking cessation: Difficulties in quitting associated with sex and substance abuse. Psychiatry Research, 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21885130>

Baker A, Kay-Lambkin F, Richmond R, Filia S, Castle D, et al. Study protocol: A randomised controlled trial investigating the effect of a healthy lifestyle intervention for people with severe mental disorders. BMC Public Health, 2011; 11(1):10. Available from: <http://www.biomedcentral.com/1471-2458/11/10>

Arbour-Nicitopoulos K, Faulkner G, Cohn T, and Selby P. Smoking cessation in women with severe mental illness: Exploring the role of exercise as an adjunct treatment. Archives of Psychiatric Nursing, 2011; 25(1):43–52. Available from: <http://www.psychiatricnursing.org/article/PIIS0883941710000592/fulltext>

Abrams DB, Graham AL, Levy DT, Mabry PL, and Orleans CT. Boosting population quits through evidence-based cessation treatment and policy. American Journal of Preventive Medicine, 2010; 38(3 Suppl):S351–63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20176308>

[No authors listed]. An official position statement of the association of women’s health, obstetric & neonatal nursing. Journal of Obstetric, Gynecologic, and Neonatal Nursing, 2010; 39(5):611–3. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1552-6909.2010.01178.x/full>

Ashton M, Miller CL, Bowden JA, and Bertossa S. People with mental illness can tackle tobacco. Australian and New Zealand Journal of Psychiatry, 2010; 44(11):1021–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21034185>

Abrantes A, Strong D, Lloyd-Richardson E, Niaura R, Kahler C, et al. Regular exercise as a protective factor in relapse following smoking cessation treatment. American Journal on Addictions, 2009; 18(1):100–1. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19219672>

Hashibe M, Straif K, Tashkin D, Morgenstern H, Greenland S, et al. Epidemiological review of marijuana use and cancer risk. *Alcohol*, 2005; 35(3):265–75.

Kalant H. Adverse effects of cannabis on health: An update of the literature since 1996. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 2004; 28(849-63).

Kassel JD, Stroud LR, and Paronis CA. Smoking, stress, and negative affect: Correlation, causation, and context across stages of smoking. *Psychol Bull*, 2003; 129(2):270–304. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12696841>

Jablensky A, McGrath J, Herrman H, Castle D, Gureje O, et al. People living with psychotic illness: An Australian study 1997-98. Mental Health Branch, Commonwealth Department of Health and Aged Care, Canberra, 1999.

Australian Bureau of Statistics. 4364.0.55.001 - national health survey: First results, 2017-18 Available from: <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/national-health-survey-first-results/latest-release>.

9A.3.5.1 Depression

Ranjan, R, Yadav, J, Ranjan, V, Venkateswaran, C, & John, D. (2024). Association of Self-reported Depressed Mood and Alcohol or Tobacco Use Among Older Adults in India: A Study Based on Longitudinal Aging Study India Wave-1. *Indian J Psychol Med*, 02537176241253338. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39564331>

Balsara, K, Iftikhar, A, Galiatsatos, P, DiClemente, C, Mattingly, B, & Kanarek, NF. (2024). Cigarette Smoking, Mental Health, Depression, Maryland Behavioral Risk Factor Surveillance System Survey, 2020. *Tob Use Insights*, 17, 1179173X241285351. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39372423>

Son, JS. (2024). The Interconnected Nature of Smoking, Depression, and Obesity in Behavioral Medicine. *Korean J Fam Med*, 45(4), 181-182. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39054837>

Blackwell, AKM, Daryan, S, Roy, D, Duffy, D, Hisler, G, Sawyer, K et al. (2024). Integrating smoking cessation treatment into usual online psychological care for people with common mental illness: Protocol for an online randomised feasibility and pilot study (ESCAPE digital). *Contemp Clin Trials*, 141, 107541. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38643854>

Hitsman, B, Papandonatos, GD, Fox, EN, Bauer, AM, Gollan, JK, Huffman, MD et al. (2024). Effect of early medication adherence on behavioral treatment utilization and smoking cessation among individuals with current or past major depressive disorder. *Addict Behav*, 151, 107952. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38199093>

Marchal-Mateos, MI, Lopez-Nunez, C, & Fernandez-Artamendi, S. (2024). Effectiveness of Contingency Management in Tobacco Smokers with Depressive Symptoms: A Systematic Review. *Subst Use Misuse*, 1-13. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38268117>

Krishnamoorthy, NK, Roohi, TF, Peddha, MS, Kinattingal, N, Wani, SUD, Krishna, KL et al. (2023). Evaluation of Antidepressant Activity of Capsaicin Nanoemulsion in Nicotine Withdrawal-Induced Depression in Mice. *Brain Sci*, 13(12). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38137115>

van de Weijer, MP, Vermeulen, J, Schranter, A, Munafo, MR, Verweij, KJH, & Treur, JL. (2024). The potential role of gray matter volume differences in the association between smoking and depression: A narrative review. *Neurosci Biobehav Rev*, 156, 105497. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38100958>

Cabrera-Serrano, A, Felici-Giovanini, ME, Diaz-Garcia, RM, Ramos-Colon, MV, & Rodriguez-Ayuso, I. (2023). Tobacco Use and Depression: A Hidden Epidemic among Smokers 18 Years and Older, Puerto Rico, 2018-2020. *P R Health Sci J*, 42(4), 298-303. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38104286>

Schnoll, R, Wileyto, EP, Bauer, AM, Fox, EN, Blumenthal, D, Quinn, MH et al. (2023). Seeing Through the Blind: Belief about Treatment Randomization and Smoking Cessation Outcome among People with Current or Past Major Depressive Disorder who Smoke in a Placebo-Controlled Trial of Varenicline. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37934573>

Gollan, JK, Liverant, G, Jao, NC, Lord, KA, Whitton, AE, Hogarth, L et al. (2023). Depression Severity Moderates Reward Learning Among Smokers with Current or Past Major Depressive Disorder in a Smoking Cessation Randomized Clinical Trial. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37943674>

Dahne, J, Wahlquist, AE, Kustanowitz, J, Natale, N, Fahey, M, Graboyes, EM et al. (2023). Behavioral Activation-Based Digital Smoking Cessation Intervention for Individuals With Depressive Symptoms: Randomized Clinical Trial. *J Med Internet Res*, 25, e49809. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37910157>

Steinberg, ML, Rosen, RL, Ganz, O, Wackowski, OA, Jeong, M, & Delnevo, CD. (2024). Communicating the benefits of quitting smoking on mental health increases motivation to quit in people with anxiety and/or depression. *Addict Behav*, 149, 107903. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37924583>

Floyd, J, Mallow, J, Wang, K, Davis, SM, Carpenter, R, & Theeke, L. (2023). Differences in smoking behaviors and readiness to change for patients with COPD and differing categories of depressive symptoms: a descriptive cross-sectional design. *BMC Pulm Med*, 23(1), 335. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37684585>

Wiener, RC. (2023). Nicotine Dependence and Postpartum Depression: A Multicenter Research Network Study. *Subst Use Misuse*, 1-5. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37545109>

Pang, RD, Tucker, C. J., Guillot, C. R., Belcher, B., & Kirkpatrick, M. G. (2023). Associations of DHEA(S) with negative and positive affect in people who smoke daily with elevated and low depression symptoms: A pilot laboratory study. *Addict Behav*, 146, 107801. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37423068>

Gravelly, S, Driezen, P, McClure, EA, Smith, DM, & Fong, GT. (2023). Prevalence of depressive symptoms and cannabis use among adult cigarette smokers in Canada: cross-sectional findings from the 2020 International Tobacco Control Policy Evaluation Project Canada Smoking and Vaping Survey. *CMAJ Open*, 11(3), E516-E526. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37311596>

McLeish, AC, Walker, KL, Keith, RJ, & Hart, JL. (2023). The Role of Perceived Neighborhood Cohesion in the Association between Depressive Symptoms and Cigarette Smoking. *Subst Use Misuse*, 1-7. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37232382>

Wu, AD, Gao, M, Aveyard, P, & Taylor, G. (2023). Smoking Cessation and Changes in Anxiety and Depression in Adults With and Without Psychiatric Disorders. *JAMA Netw Open*, 6(5), e2316111. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37256615>

Bataineh, B S, Wilkinson, AV, Sumbe, A, Clendennen, SL, Chen, B, Messiah, SE, & Harrell, MB. (2023). The Association Between Tobacco and Cannabis Use and the Age of Onset of Depression and Anxiety Symptoms Among Adolescents and Young Adults. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37042355>

Al Bataineh, MT, Kunstner, A, Dash, NR, Abdulsalam, RM, Al-Kayyali, RZA., Adi, MB et al. (2023). Corrigendum: Altered composition of the oral microbiota in depression among cigarette smokers: A pilot study. *Front Psychiatry*, 14, 1175044. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37009095>

Carroll, AJ, Huffman, MD, Wileyto, EP, Khan, SK, Fox, E, Smith, JD et al. (2023). Change in cardiovascular health among adults with current or past major depressive disorder enrolled in intensive smoking cessation treatment. *J Affect Disord*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37119868>

Hitsman, B, Papandonatos, GD, Gollan, JK, Huffman, MD, Niaura, R, Mohr, DC et al. (2023). Efficacy and Safety of Combination Behavioral Activation for Smoking Cessation and Varenicline for Treating Tobacco Dependence among Individuals with Current or Past Major Depressive Disorder: A 2x2 Factorial, Randomized, Placebo-Controlled Trial. *Addiction*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37069490>

Krotter, A, Aonso-Diego, G, Garcia-Perez, A, Garcia-Fernandez, G, & Secades-Villa, R. (2023). Post-Cessation Weight Gain among Smokers with Depression Predicts Smoking Relapse. *J Dual Diagn*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37015070>

Schnoll, R, Barrila, GM, Dalsimer, S, Hosie Quinn, M, Bauer, AM, Fox, E et al. (2023). Treatment adherence in a smoking cessation clinical trial for individuals with current or past major depressive disorder: Predictors and association with cessation. *Addict Behav*, 143, 107686. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36893514>

Steinberg, ML, Rosen, RL, Ganz, O, Wackowski, OA, Jeong, M, & Delnevo, CD. (2023). Communicating the benefits of quitting smoking on mental health increases motivation to quit in people with anxiety and/or depression: A randomized trial of two health messages. *medRxiv*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36865337>

- Baldassarri, SR, Asch, RH, Hillmer, AT, Pietrzak, RH, DellaGioia, N, Esterlis, I, & Davis, MT. (2023). Nicotine Use and Metabotropic Glutamate Receptor 5 in Individuals With Major Depressive and Posttraumatic Stress Disorders. *Chronic Stress (Thousand Oaks)*, 7, 24705470231154842. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36843572>
- Brinkman, H. R., Hoyt, D. L., Fedorenko, E. J., Mendes, W. B., & Leyro, T. M. (2023). Cardiac Vagal Control Among Community Cigarette Smokers with Low to Moderate Depressive Symptoms. *Appl Psychophysiol Biofeedback*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36732418>
- Quaioto, B. R., Borcoi, A. R., Mendes, S. O., Doblaz, P. C., Dos Santos Vieira, T., Arantes Moreno, I. A., . . . Alvares-da-Silva, A. M. (2023). Tobacco use modify exon IV BDNF gene methylation levels in depression. *J Psychiatr Res*, 159, 240-248. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36753898>
- Wu, Z., Yue, Q., Zhao, Z., Wen, J., Tang, L., Zhong, Z., . . . Zhang, X. (2023). A cross-sectional study of smoking and depression among US adults: NHANES (2005-2018). *Front Public Health*, 11, 1081706. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36794066>
- Correction: Time trends in access to smoking cessation support for people with depression or severe mental illness: a cohort study in English primary care. (2023). *BMJ Open*, 13(2), e048341corr048341. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36792340>
- Fredman Stein, K, Sawyer, K, Daryan, S, Allen, J, & Taylor, G. (2022). Service-user experiences of an integrated psychological intervention for depression or anxiety and tobacco smoking in improving access to psychological therapies services: A qualitative investigation into mechanisms of change in quitting smoking. *Health Expect*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36482862>
- Zhang, H, Gilbert, E, Hussain, SVeldhuizen, S, Le Foll, B, Selby, P, & Zawertailo, L. (2022). Effectiveness of bupropion and varenicline for smokers with baseline depressive symptoms. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36520964>
- Komiyama, M, Ozaki, Y, Wada, H, Yamakage, H, Satoh-Asahara, N, Yasoda, A et al. (2022). Randomized double-blind placebo-controlled multicenter trial for the effects of a polyherbal remedy, Yokukansan (YiganSan), in smokers with depressive tendencies. *BMC Complement Med Ther*, 22(1), 311. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36434692>
- Liu, Q, Leng, P, Gu, Y, Shang, X, Zhou, Y, Zhang, H et al. (2022). The dose-effect relationships of cigarette and alcohol consumption with depressive symptoms: a multiple-center, cross-sectional study in 5965 Chinese middle-aged and elderly men. *BMC Psychiatry*, 22(1), 657. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36284280>
- Mesdom, P, Colle, R, Becquemont, L, Chappell, K, David, DJ, Mendez-David, I et al. (2022). Tobacco use is associated with low peripheral beta-arrestin 1 levels in major depression: A preliminary report. *Drug Alcohol Depend*, 240, 109653. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36209675>
- Mildrum Chana, S, Palenski, PE, Hawes, ES, Wolford-Clevenger, C, Thomas, SJ, Gamble, KL, & Cropsey, KL. (2022). Discrimination and insomnia: Examining depressive symptoms and nicotine withdrawal through a serial mediation model in a sample of smokers. *Addict Behav*, 137, 107506. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36244244>

Denche-Zamorano, A, Mendoza-Munoz, DM, Pastor-Cisneros, R, Adsuar, JC, Carlos-Vivas, J, Franco-Garcia, JM et al. (2022). A Cross-Sectional Study on the Associations between Physical Activity Level, Depression, and Anxiety in Smokers and Ex-Smokers. *Healthcare (Basel)*, 10(8). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36011059>

Han, JW, Lee, H, Min, S, & Lee, B. (2022). Depression with Chronic Disease Is Associated with Increased Use of Medical Services and Medical Expenses in Hardcore Smokers. *Healthcare (Basel)*, 10(8). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36011061>

Hahad, O, Beutel, M, Gilan, D A, Michal, M, Schulz, A, Pfeiffer, N et al. (2022). The association of smoking and smoking cessation with prevalent and incident symptoms of depression, anxiety, and sleep disturbance in the general population. *J Affect Disord*, 313, 100-109. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35777492>

Liu, S, Jiang, H, Zhang, D, Luo, J, & Zhang, H. (2022). The Association between Smoking Cessation and Depressive Symptoms: Diet Quality Plays a Mediating Role. *Nutrients*, 14(15). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35893901>

Wang, Y, Duan, Z, Romm, KF, Ma, Y, Douglas Evans, W, Bennett, B et al. (2022). Bidirectional associations between depressive symptoms and cigarette, e-cigarette, cannabis, and alcohol use: Cross-lagged panel analyses among young adults before and during COVID-19. *Addict Behav*, 134, 107422. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35853404>

Cinciripini PM, Kypriotakis G, Green C, Lawrence D, Anthenelli RM, et al. The effects of varenicline, bupropion, nicotine patch, and placebo on smoking cessation among smokers with major depression: A randomized clinical trial. *Depress Anxiety*, 2022; 39(5):429-40. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35535436>

Bondo P, Hoye A, Lochen ML, and Bramness JG. The relationship between smokeless tobacco (snus) and anxiety and depression amongst adults and elderly. A comparison to smoking in the tromsø study. *Addiction*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35603916>

Abdollahpour I, Golestannejad Z, Salimi Y, Nedjat S, Aguilar-Palacioc I, et al. Joint waterpipe and cigarette smoking as key correlate of history of depression in iranian population: A cross-sectional study. *Tob Use Insights*, 2022; 15:1179173X221100402. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35600561>

Tucker CJ, Bello MS, Weinberger AH, D'Orazio LM, Kirkpatrick MG, et al. Association of depression symptom level with smoking urges, cigarette withdrawal, and smoking reinstatement: A preliminary laboratory study. *Drug and Alcohol Dependence*, 2022; 232:109267. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35042097>

Pang RD, Chai SH, Tucker CJ, Weinberger AH, D'Orazio LM, et al. Effects of cigarette abstinence on negative and positive affect by depression symptom levels: A lab study. *J Affect Disord*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35341814>

Martinez-Vispo C, Lopez-Duran A, Senra C, and Becona E. Brooding rumination and anxiety sensitivity: Associations with depressive and anxiety symptoms in treatment-seeking smokers. *Psicothema*, 2022; 34(1):49-55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35048895>

Kiviruusu O, Berg N, Piirtola M, Viertio S, Suvisaari J, et al. Life course associations between smoking and depressive symptoms. A 30-year finnish follow-up study. *Nord J Psychiatry*, 2022;1. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35316159>

Fatani BZ, Al-Yahyawi H, Raggam A, Al-Ahdal M, Alzyoud S, et al. Perceived stress and willingness to quit smoking among patients with depressive and anxiety disorders seeking treatment. *Health Sci Rep*, 2022; 5(2):e503. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35229052>

Yonek JC, Meacham MC, Shumway M, Tolou-Shams M, and Satre DD. Smoking reduction is associated with lower alcohol consumption and depressive symptoms among young adults over one year. *Drug and Alcohol Dependence*, 2021; 227:108922. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34364192>

Tam J, Warner KE, Zivin K, Taylor GMJ, and Meza R. The potential impact of widespread cessation treatment for smokers with depression. *American Journal of Preventive Medicine*, 2021; 61(5):674–82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34244005>

Schnoll R, Paul Wileyto E, Bauer AM, Fox E, Leone F, et al. Comparing the rate of nicotine metabolism among smokers with current or past major depressive disorder. *American Journal on Addictions*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33772971>

Liu NH, Wu C, Perez-Stable EJ, and Munoz RF. Longitudinal association between smoking abstinence and depression severity in those with baseline current, past, and no history of major depressive episode in an international online tobacco cessation study. *Nicotine & Tobacco Research*, 2021; 23(2):267–75. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32149344>

Kang SH, Jeong W, Jang SI, and Park EC. The effect of depression status change on daily cigarette smoking amount according to sex: An eleven-year follow up study of the korea welfare panel study. *BMC Public Health*, 2021; 21(1):1304. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34217238>

Gonzalez-Roz A, Secades-Villa R, Garcia-Fernandez G, Martinez-Loredo V, and Alonso-Perez F. Depression symptom profiles and long-term response to cognitive behavioral therapy plus contingency management for smoking cessation. *Drug and Alcohol Dependence*, 2021; 225:108808. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34198211>

Garcia-Perez A, Weidberg S, Gonzalez-Roz A, Alonso-Perez F, and Secades-Villa R. Corrigendum to "relationship between delay discounting and depression in cigarette smokers and non-smokers" [addict. Behav. 103 (2020) 106251]. *Addictive Behaviors*, 2021:106809. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33423793>

de Boer N, Vermeulen J, Lin B, van Os J, Ten Have M, et al. Longitudinal associations between alcohol use, smoking, genetic risk scoring and symptoms of depression in the general population: A prospective 6-year cohort study. *Psychological Medicine*, 2021:1-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35023464>

Yao Y, Xu Y, Cai Z, Liu Q, Ma Y, et al. Determination of shared genetic etiology and possible causal relations between tobacco smoking and depression. *Psychological Medicine*, 2020:1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32249730>

Weinberger AH, Chaiton MO, Zhu J, Wall MM, Hasin DS, et al. Trends in the prevalence of current, daily, and nondaily cigarette smoking and quit ratios by depression status in the u.s.: 2005-2017. *American Journal of Preventive Medicine*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32156490>

Weidberg S, Gonzalez-Roz A, Garcia-Fernandez G, and Secades-Villa R. Activation level as a mediator between behavioral activation, sex, and depression among treatment-seeking smokers. *Addictive Behaviors*, 2020:106715. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33131968>

Watson NL, Heffner JL, Mull KE, McClure JB, and Bricker JB. Which method of assessing depression and anxiety best predicts smoking cessation: Screening instruments or self-reported conditions? *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32484870>

Waters AF, Peltier MR, Roys MR, Stewart SA, and Copeland AL. Smoking and suicidal ideation among college students: Smoking expectancies as potential moderators. *J Am Coll Health*, 2020:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32027235>

Sathian B, Menezes RG, Asim M, Mekkodathil A, Sreedharan J, et al. Cigarette smoking dose-response and suicidal ideation among young people in Nepal: A cross-sectional study. *Nepal J Epidemiol*, 2020; 10(1):821-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32257512>

Riblet NB, Gottlieb DJ, Hoyt JE, Watts BV, and Shiner B. An analysis of the relationship between chronic obstructive pulmonary disease, smoking and depression in an integrated healthcare system. *General Hospital Psychiatry*, 2020; 64:72-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32279024>

Ranjit A, Latvala A, Kinnunen TH, Kaprio J, and Korhonen T. Depressive symptoms predict smoking cessation in a 20-year longitudinal study of adult twins. *Addictive Behaviors*, 2020; 108:106427. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32361366>

Martinez-Vispo C, Lopez-Duran A, Senra C, Rodriguez-Cano R, Fernandez Del Rio E, et al. Environmental reward and depressive symptoms in the relationship between avoidance and cigarette dependence in treatment-seeking smokers. *Psicothema*, 2020; 32(2):176-81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32249742>

Martinez-Vispo C, Lopez-Duran A, Senra C, Rodriguez-Cano R, Fernandez Del Rio E, et al. Behavioral activation and smoking cessation outcomes: The role of depressive symptoms. *Addictive Behaviors*, 2020; 102:106183. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31809878>

Kurtulus S, Can R, and Sak ZHA. Assessment of the relationship between smoking and depression in pregnant women. *Journal of Immigrant and Minority Health*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32632720>

Harrison R, Munafo MR, Davey Smith G, and Wootton RE. Examining the effect of smoking on suicidal ideation and attempts: Triangulation of epidemiological approaches. *The British Journal of Psychiatry*, 2020;1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32290872>

Han JW and Lee H. The effects of hardcore smokers' depression and self-esteem on daily smoking amount. *Archives of Psychiatric Nursing*, 2020; 34(3):149-58. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32513465>

Giovenco DP, Spillane TE, Baig SA, Dumas SE, Dongchung TY, et al. Demographic and psychological moderators of the relationship between neighborhood cigarette advertising and current smoking in new york city. *Health Place*, 2020; 66:102441. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32947186>

Garcia-Perez A, Vallejo-Seco G, Weidberg S, Gonzalez-Roz A, and Secades-Villa R. Long-term changes in delay discounting following a smoking cessation treatment for patients with depression. *Drug and Alcohol Dependence*, 2020:108007. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32370930>

Cho MS. Use of alcohol, tobacco, and caffeine and suicide attempts: Findings from a nationally representative cross-sectional study. *J Prim Care Community Health*, 2020; 11:2150132720913720. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32193974>

Bazargan M, Cobb S, Castro Sandoval J, and Assari S. Smoking status and well-being of underserved african american older adults. *Behav Sci (Basel)*, 2020; 10(4). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32326442>

Aubin HJ, Legleye S, Thomas D, and Berlin I. Tobacco smoking: The likely confounder of the association between heart diseases and suicide. *J Intern Med*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32237242>

Alibekova R, Huang JP, Lee TS, Au HK, and Chen YH. Corrigendum to "effects of smoking on perinatal depression and anxiety in mothers and fathers: A prospective cohort study" *journal of affective disorders* 193(2016) 18-26]. *J Affect Disord*, 2020; 274:1218. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32663956>

Secades-Villa R, Gonzalez-Roz A, Vallejo-Seco G, Weidberg S, Garcia-Perez A, et al. Additive effectiveness of contingency management on cognitive behavioural treatment for smokers with depression: Six-month abstinence and depression outcomes. *Drug and Alcohol Dependence*, 2019; 204:107495. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31491583>

Rubin LF, Haaga DAF, Pearson JL, and Gunthert KC. Depression as a moderator of the prospective relationship between mood and smoking. *Health Psychology*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31682148>

Rogers ES. Financial distress and smoking-induced deprivation in smokers with depression. *American Journal of Health Behavior*, 2019; 43(1):219–27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30522579>

Ranjit A, Korhonen T, Buchwald J, Heikkila K, Tuulio-Henriksson A, et al. Testing the reciprocal association between smoking and depressive symptoms from adolescence to adulthood: A longitudinal twin study. *Drug and Alcohol Dependence*, 2019; 200:64-70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31100637>

Ranjit A, Buchwald J, Latvala A, Heikkila K, Tuulio-Henriksson A, et al. Predictive association of smoking with depressive symptoms: A longitudinal study of adolescent twins. *Prev Sci*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31069603>

Petroulia I, Kyriakos CN, Papadakis S, Tzavara C, Filippidis FT, et al. Corrigendum: Patterns of tobacco use, quit attempts, readiness to quit and self-efficacy among smokers with anxiety or depression: Findings among six countries of the eurest-plus itc europe surveys. *Tobacco Induced Diseases*, 2019; 17:83. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31889945>

Marsden DG, Loukas A, Chen B, Perry CL, and Wilkinson AV. Associations between frequency of cigarette and alternative tobacco product use and depressive symptoms: A longitudinal study of young adults. *Addictive Behaviors*, 2019; 99:106078. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31430619>

Lee JY, Brook JS, Finch SJ, Kim W, and Brook DW. Single and dual diagnoses of major depressive disorder and posttraumatic stress disorder predicted by triple comorbid trajectories of tobacco, alcohol, and marijuana use among urban adults. *Substance Abuse*, 2019:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30888260>

Lee EJ. Long-term effects of smoking cessation on depressive symptoms, resilience, coping skills, and serotonin. *Psychiatr Q*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31845167>

Kim HH, Kim HJ, and Kim EM. Exploring the association between suicidality and smoking behavior among school-based children in lao people's democratic republic (pdr). *J Psychoactive Drugs*, 2019:1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31888423>

Iyer AS, Holm KE, Bhatt SP, Kim V, Kinney GL, et al. Symptoms of anxiety and depression and use of anxiolytic-hypnotics and antidepressants in current and former smokers with and without copd - a cross sectional analysis of the copdgene cohort. *Journal of Psychosomatic Research*, 2019; 118:18-26. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30782350>

Ho CSH, Tan ELY, Ho RCM, and Chiu MYL. Relationship of anxiety and depression with respiratory symptoms: Comparison between depressed and non-depressed smokers in Singapore. *International Journal of Environmental Research and Public Health*, 2019; 16(1). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30626156>

Heffner JL, Watson NL, Serfozo E, Mull KE, MacPherson L, et al. A behavioral activation mobile health app for smokers with depression: Development and pilot evaluation in a single-arm trial. *JMIR Form Res*, 2019; 3(4):e13728. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31774405>

Gonzalez-Roz A, Secades-Villa R, Weidberg S, Muniz J, and MacKillop J. Characterizing the reinforcing value of tobacco using a cigarette purchase task: An item response theory approach. *Experimental*

and Clinical Psychopharmacology, 2019. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31556676>

Garcia-Perez A, Weidberg S, Gonzalez-Roz A, Alonso-Perez F, and Secades-Villa R. Relationship between delay discounting and depression in cigarette smokers and non-smokers. Addictive Behaviors, 2019; 103:106251. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31874376>

Fujita T, Babazono A, Harano Y, and Jiang P. Risk of depressive disorders after tobacco smoking cessation: A retrospective cohort study in fukuoka, Japan. BMJ Open, 2019; 9(3):e025124. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30904860>

Cheslack-Postava K, Wall MM, Weinberger AH, and Goodwin RD. Increasing depression and substance use among former smokers in the United States, 2002-2016. American Journal of Preventive Medicine, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31443956>

Carroll AJ, Huffman MD, Zhao L, Jacobs DR, Stewart JC, et al. Associations between depressive symptoms, cigarette smoking, and cardiovascular health: Longitudinal results from cardia. J Affect Disord, 2019; 260:583-91. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31539696>

Carroll AJ. Elucidating directionality between smoking and depression. Journal of Psychosomatic Research, 2019; 125:109790. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31421321>

Akambase JA, Miller NE, Garrison GM, Stadem P, Talley H, et al. Depression outcomes in smokers and nonsmokers: Comparison of collaborative care management versus usual care. J Prim Care Community Health, 2019; 10:2150132719861265. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31303098>

Weinberger AH, Bandiera FC, Leventhal AM, Dierker LC, Gbedemah M, et al. Socioeconomic disparities in smoking among u.S. Adults with depression, 2005-2014. American Journal of Preventive Medicine, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29674184>

Weidberg S, Vallejo-Seco G, Gonzalez-Roz A, Garcia-Perez A, and Secades-Villa R. In-treatment cigarette demand among treatment-seeking smokers with depressive symptoms. Addictive Behaviors, 2018; 82:35-43. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29482033>

Wang K, Liu Y, Ouedraogo Y, Wang N, Xie X, et al. Principal component analysis of early alcohol, drug and tobacco use with major depressive disorder in US adults. Journal of Psychiatric Research, 2018; 100:113-20. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29518578>

Tomita A and Manuel JJ. Evidence on the association between cigarette smoking and incident depression from the south african national income dynamics study 2008-2015: Mental health implications for a resource-limited setting. Nicotine & Tobacco Research, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30107603>

Taylor GMJ and Munafo MR. What about treatment of smoking to improve survival and reduce depression? Lancet Psychiatry, 2018; 5(6):464. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29857836>

Stubbs B, Vancampfort D, Firth J, Solmi M, Siddiqi N, et al. Association between depression and smoking: A global perspective from 48 low- and middle-income countries. *Journal of Psychiatric Research*, 2018; 103:142-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29852421>

Sharpe M. What about treatment of smoking to improve survival and reduce depression? - authors' reply. *Lancet Psychiatry*, 2018; 5(6):464-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29857837>

Rodriguez-Cano R, Paulus DJ, Zvolensky MJ, Lopez-Duran A, Martinez-Vispo C, et al. Depressive symptoms in the trajectory of craving during smoking cessation treatment: A latent growth curve model. *American Journal of Drug and Alcohol Abuse*, 2018:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29420081>

Rahman MS, Forsell Y, Hallgren M, and Galanti MR. Tobacco use does not influence the response to non-pharmacologic depression treatment: A secondary analysis of the regassa randomized controlled trial. *Psychiatry Res*, 2018; 261:442-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29395870>

Raffetti E, Donato F, Forsell Y, and Galanti MR. Longitudinal association between tobacco use and the onset of depressive symptoms among swedish adolescents: The kupol cohort study. *Eur Child Adolesc Psychiatry*, 2018; 28(5):695-704. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30315361>

Porcu M, Machado R, Urbano M, Verri WA, Jr., Rossaneis AC, et al. Depressed female smokers have higher levels of soluble tumor necrosis factor receptor 1. *Addictive Behaviors Reports*, 2018; 7:90-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29892702>

Lopez Ibanez de Aldecoa A, Cedillo Gomez S, Bruguera E, Garcia-Portilla MP, and Bobes J. Smoking status and cost of illness in patients with depressive disorder based on the national health survey in Spain. *Substance Use and Misuse*, 2018:1-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30585110>

Lemieux AM, Nakajima M, Saif-Ali R, Al-Habori M, Dokam A, et al. Anger, anxiety, and depressive affect as predictors of stress-induced cortisol production in khat and tobacco users. *Addictive Behaviors*, 2018; 82:195-201. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29551550>

Lee J, Kim TH, Min S, Kim MH, Park KC, et al. Depressive symptoms and suicidal behaviours in adolescent non-daily smokers compared to daily smokers and never-smokers in Korea: National cross-sectional study. *PLoS One*, 2018; 13(11):e0207182. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6241131/pdf/pone.0207182.pdf>

Lechner WV, Sidhu NK, Cioe PA, and Kahler CW. Effects of time-varying changes in tobacco and alcohol use on depressive symptoms following pharmaco-behavioral treatment for smoking and heavy drinking. *Drug and Alcohol Dependence*, 2018; 194:173-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30445275>

Komiyama M, Yamakage H, Satoh-Asahara N, Ozaki Y, Morimoto T, et al. Sex differences in nicotine dependency and depressive tendency among smokers. *Psychiatry Res*, 2018; 267:154-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29908483>

Jung M. The relationship between alcohol abuse and suicide risk according to smoking status: A cross-sectional study. *J Affect Disord*, 2018; 244:164-70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30342376>

Jiang H, Li S, and Yang J. Work stress and depressive symptoms in fishermen with a smoking habit: A mediator role of nicotine dependence and possible moderator role of expressive suppression and cognitive reappraisal. *Front Psychol*, 2018; 9:386. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29632504>

Jia H, Zack MM, Gottesman, II, and Thompson WW. Associations of smoking, physical inactivity, heavy drinking, and obesity with quality-adjusted life expectancy among US adults with depression. *Value Health*, 2018; 21(3):364-71. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29566844>

Janes AC, Zegel M, Ohashi K, Betts J, Molokotos E, et al. Nicotine normalizes cortico-striatal connectivity in non-smoking individuals with major depressive disorder. *Neuropsychopharmacology*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29795403>

Huffman AL, Bromberg JE, and Augustson EM. Lifetime depression, other mental illness, and smoking cessation. *American Journal of Health Behavior*, 2018; 42(4):90-101. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29973314>

Heinzman JT, Hoth KF, Cho MH, Sakornsakolpat P, Regan EA, et al. Gwas and systems biology analysis of depressive symptoms among smokers from the copdgene cohort. *J Affect Disord*, 2018; 243:16-22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30219690>

Gonzalez-Roz A, Secades-Villa R, and Muniz J. Validity evidence of the behavioral activation for depression scale-short form among depressed smokers. *Int J Clin Health Psychol*, 2018; 18(2):162-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30487921>

Endrighi R, McQuaid EL, Bartlett YK, Clawson AH, and Borrelli B. Parental depression is prospectively associated with lower smoking cessation rates and poor child asthma outcomes. *Annals of Behavioral Medicine*, 2018; 52(3):195-203. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29538661>

Doran N, Dubrava S, and Anthenelli RM. Effects of varenicline, depressive symptoms, and region of enrollment on smoking cessation in depressed smokers. *Nicotine & Tobacco Research*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29471329>

Cohn AM. Never, non-daily, and daily smoking status and progression to daily cigarette smoking as correlates of major depressive episode in a national sample of youth: Results from the national survey of drug use and health 2013 to 2015. *Addictive Behaviors*, 2018; 84:118-25. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29684762>

Chang HB, Munroe S, Gray K, Porta G, Douaihy A, et al. The role of substance use, smoking, and inflammation in risk for suicidal behavior. *J Affect Disord*, 2018; 243:33-41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30223137>

Barnes SA, Sheffler DJ, Semenova S, Cosford NDP, and Beshpalov A. Metabotropic glutamate receptor 5 as a target for the treatment of depression and smoking: Robust preclinical data but inconclusive clinical efficacy. *Biological Psychiatry*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29628194>

Badr HE and Francis K. Psychosocial perspective and suicidal behaviors correlated with adolescent male smoking and illicit drug use. *Asian J Psychiatr*, 2018; 37:51-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30118993>

Alghzawi H. Probability and correlates of nicotine dependence among smokers with and without major depressive disorder: Results from the national epidemiology survey on alcohol and related conditions. *Perspect Psychiatr Care*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29350765>

Aldi GA, Bertoli G, Ferraro F, Pezzuto A, and Cosci F. Effectiveness of pharmacological or psychological interventions for smoking cessation in smokers with major depression or depressive symptoms: A systematic review of the literature. *Substance Abuse*, 2018; 39(3):289–306. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29436984>

Witt DR, Garrison GM, Gonzalez CA, Witt TJ, and Angstman KB. Six-month outcomes for collaborative care management of depression among smoking and nonsmoking patients. *Health Serv Res Manag Epidemiol*, 2017; 4:2333392817721648. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28890910>

Weinberger AH, Gbedemah M, Wall MM, Hasin DS, Zvolensky MJ, et al. Depression among non-daily smokers compared to daily smokers and never-smokers in the United States: An emerging problem. *Nicotine & Tobacco Research*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28339571>

Taylor G, Aveyard P, Meer RV, Toze D, Stuijzand B, et al. Impact of variation in functions and delivery on the effectiveness of behavioural and mood management interventions for smoking cessation in people with depression: Protocol for a systematic review and meta-analysis. *BMJ Open*, 2017; 7(11):e018617. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29150479>

Secades-Villa R, Weidberg S, Gonzalez-Roz A, Reed DD, and Fernandez-Hermida JR. Cigarette demand among smokers with elevated depressive symptoms: An experimental comparison with low depressive symptoms. *Psychopharmacology (Berl)*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29143193>

Secades-Villa R, Gonzalez-Roz A, Garcia-Perez A, and Becona E. Psychological, pharmacological, and combined smoking cessation interventions for smokers with current depression: A systematic review and meta-analysis. *PLoS One*, 2017; 12(12):e0188849. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29206852>

Rodriguez-Cano R, Paulus DJ, Lopez-Duran A, Martinez-Vispo C, Fernandez Del Rio E, et al. The interplay of history of depression and craving in terms of smoking relapse among treatment seeking smokers. *Journal of Addictive Diseases*, 2017; 36(3):175-82. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28388278>

Rodriguez-Cano R, Lopez-Duran A, Paulus DJ, Martinez-Vispo C, Fernandez Del Rio E, et al. The interaction of depressive symptoms and hazardous drinking in relation to tobacco craving among treatment seeking depressed smokers: Sex differences. *Journal of Addiction Medicine*, 2017.

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29219856>

Rejas-Gutierrez J, Bruguera E, and Cedillo S. Modelling a budgetary impact analysis for funding drug-based smoking cessation therapies for patients with major depressive disorder in Spain. *European Psychiatry*, 2017; 45:41-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28728094>

Oliveira P, Ribeiro J, Donato H, and Madeira N. Smoking and antidepressants pharmacokinetics: A systematic review. *Annals of General Psychiatry*, 2017; 16:17. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28286537>

Kostrubiak DE, Vacchi-Suzzi C, Smith DM, and Meliker JR. Blood cadmium and depressive symptoms: Confounded by cigarette smoking. *Psychiatry Res*, 2017; 256:444-7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28709058>

Kopetz C, MacPherson L, Mitchell AD, Houston-Ludlam AN, and Wiers RW. A novel training approach to activate alternative behaviors for smoking in depressed smokers. *Experimental and Clinical Psychopharmacology*, 2017; 25(1):50-60. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28150972>

Hogarth L, Mathew AR, and Hitsman B. Current major depression is associated with greater sensitivity to the motivational effect of both negative mood induction and abstinence on tobacco-seeking behavior. *Drug and Alcohol Dependence*, 2017; 176:1-6. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28460322>

Green M, Turner S, and Sareen J. Smoking and suicide: Biological and social evidence and causal mechanisms. *Journal of Epidemiology and Community Health*, 2017. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28679536>

Gonzalez MO, Goudriaan AE, Perico CA, Waisman Campos M, de Andrade AG, et al. Neural correlates of depressive symptoms in smokers - a systematic review of imaging studies. *Substance Use and Misuse*, 2017:1-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28742414>

Becona E, Martinez-Vispo C, Senra C, Lopez-Duran A, Rodriguez-Cano R, et al. Cognitive-behavioral treatment with behavioral activation for smokers with depressive symptomatology: Study protocol of a randomized controlled trial. *BMC Psychiatry*, 2017; 17(1):134. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28390417>

Abrantes AM, Farris SG, Garnaat SL, Minto A, Brown RA, et al. The role of physical activity enjoyment on the acute mood experience of exercise among smokers with elevated depressive symptoms.

Ment Health Phys Act, 2017; 12:37-43. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28989344>

Stepankova L, Kralikova E, Zvolenska K, Pankova A, Ovesna P, et al. Depression and smoking cessation: Evidence from a smoking cessation clinic with 1-year follow-up. *Annals of Behavioral Medicine*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28035641>

Cooper J, Borland R, Yong HH, and Fotuhi O. The impact of quitting smoking on depressive symptoms: Findings from the international Tobacco Control four-country survey. *Addiction*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26918680>

Yue Y, Hong L, Guo L, Gao X, Deng J, et al. Gender differences in the association between cigarette smoking, alcohol consumption and depressive symptoms: A cross-sectional study among chinese adolescents. *Sci Rep*, 2015; 5:17959. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26639938>

Williams JM and Steinberg ML. Report ignores risk factor of tobacco in assessing suicidality. *Journal of Clinical Psychiatry*, 2015; 76(11):1570–1. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26646034>

Rodriguez-Cano R, Lopez-Duran A, Del Rio EF, Martinez-Vispo C, Martinez U, et al. Smoking cessation and depressive symptoms at 1-, 3-, 6-, and 12-months follow-up. *J Affect Disord*, 2015; 191:94–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26655118>

Morozova M, Rabin RA, and George TP. Co-morbid tobacco use disorder and depression: A re-evaluation of smoking cessation therapy in depressed smokers. *American Journal on Addictions*, 2015; 24(8):687–94. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26354720>

Kawada T. Comment on "atypical epidemiologic finding in association between depression and alcohol use or smoking in Korean male: Korean longitudinal study of aging". *Psychiatry Investig*, 2015; 12(4):573–4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26508973>

Foster DW, Khalil GE, Farris SG, Barnighausen TW, and Prokhorov AV. Effects of readiness to change, quit history, and health perceptions on depressive symptoms among young adult smokers. *Int J Psychol Psychoanal*, 2015; 1(1). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26457334>

Schroeder S. Depression, smoking, and heart disease: How can psychiatrists be effective? *The American Journal of Psychiatry*, 2011; 168(9):876–8. Available from: <http://ajp.psychiatryonline.org/cgi/content/full/168/9/876>

Khaled SM, Bulloch AG, Williams JV, Lavorato DH, and Patten SB. Major depression is a risk factor for shorter time to first cigarette irrespective of the number of cigarettes smoked per day: Evidence from a national population health survey. *Nicotine & Tobacco Research*, 2011; 13(11):1059–67. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21832274>

9A.3.5.2 Anxiety

Redmond, BY, Bizier, A, Salwa, A, Brown, RA, Garey, L, & Zvolensky, MJ. (2024). Transdiagnostic Risk Factors for Reasons for Smoking: Evaluating the Concurrent Role of Distress Tolerance and

Anxiety Sensitivity. *Int J Behav Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39480619>

Smits, JAJ, Zvolensky, MJ, Cheney, MK, Rosenfield, D, Brown, RA, Manser, SS et al. (2024). Efficacy and implementation of exercise-based smoking cessation treatment for adults with high anxiety sensitivity (STEP): Study protocol for a randomized controlled trial. *Contemp Clin Trials*, 141, 107521. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38580103>

Blackwell, AKM, Daryan, S, Roy, D, Duffy, D, Hisler, G, Sawyer, K et al. (2024). Integrating smoking cessation treatment into usual online psychological care for people with common mental illness: Protocol for an online randomised feasibility and pilot study (ESCAPE digital). *Contemp Clin Trials*, 141, 107541. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38643854>

Steinberg, ML, Rosen, RL, Ganz, O, Wackowski, OA, Jeong, M, & Delnevo, CD. (2024). Communicating the benefits of quitting smoking on mental health increases motivation to quit in people with anxiety and/or depression. *Addict Behav*, 149, 107903. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37924583>

Wu, AD, Gao, M, Aveyard, P, & Taylor, G. (2023). Smoking Cessation and Changes in Anxiety and Depression in Adults With and Without Psychiatric Disorders. *JAMA Netw Open*, 6(5), e2316111. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37256615>

Bataineh, B S, Wilkinson, AV, Sumbe, A, Clendennen, SL, Chen, B, Messiah, SE, & Harrell, MB. (2023). The Association Between Tobacco and Cannabis Use and the Age of Onset of Depression and Anxiety Symptoms Among Adolescents and Young Adults. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37042355>

Correa-Fernandez, V, Blalock, JA, Piper, ME, Canino, G, & Wetter, DW. (2023). Acceptance and Commitment Therapy Wellness Program for Latine Adults Who Smoke and Have Psychological Distress: Protocol for a Feasibility Study. *JMIR Res Protoc*, 12, e44146. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37014678>

Suarez-Castro, D, Barroso-Hurtado, M, Martinez-Vispo, C, Becona, E, & Lopez-Duran, A. (2023). Boredom Susceptibility and Quit Smoking: The Role of Anxiety Symptoms. *J Psychol*, 1-10. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36944190>

Steinberg, ML, Rosen, RL, Ganz, O, Wackowski, OA, Jeong, M, & Delnevo, CD. (2023). Communicating the benefits of quitting smoking on mental health increases motivation to quit in people with anxiety and/or depression: A randomized trial of two health messages. *medRxiv*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36865337>

Fredman Stein, K, Sawyer, K, Daryan, S, Allen, J, & Taylor, G. (2022). Service-user experiences of an integrated psychological intervention for depression or anxiety and tobacco smoking in improving access to psychological therapies services: A qualitative investigation into mechanisms of change in quitting smoking. *Health Expect*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36482862>

Orr, MF, Heggeness, LF, Mehta, N, Moya, J, Manning, K, Garey, L et al. (2022). Anxiety sensitivity and cigarette use on cannabis use problems, perceived barriers for cannabis cessation, and self-efficacy for quitting among adults with cannabis use disorder. *Addict Behav*, 137, 107509. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36194977>

Zvolensky, MJ, Redmond, BY, Smit, T, Matoska, CT, Mayorga, NA, Gallagher, M. & Garey, L. (2022). Anxiety sensitivity and its association with perceived barriers for quitting, smoking inflexibility, and severity of quit problems among Black smokers. *J Ethn Subst Abuse*, 1-18. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36190317>

Denche-Zamorano, A, Mendoza-Munoz, DM, Pastor-Cisneros, R, Adsuar, JC, Carlos-Vivas, J, Franco-Garcia, JM et al. (2022). A Cross-Sectional Study on the Associations between Physical Activity Level, Depression, and Anxiety in Smokers and Ex-Smokers. *Healthcare (Basel)*, 10(8). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36011059>

Hwang, J, & Borah, P. (2022). Anxiety Disorder and Smoking Behavior: The Moderating Effects of Entertainment and Informational Television Viewing. *Int J Environ Res Public Health*, 19(15). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35954515>

Huang Z, Zhang L, Wang J, Xu L, Liu Z, et al. Social support and subjective well-being among postgraduate medical students: The mediating role of anxiety and the moderating role of alcohol and tobacco use. *Heliyon*, 2021; 7(12):e08621. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34988318>

Zvolensky MJ, Rogers AH, Garey L, Ditre JW, Shepherd JM, et al. The role of anxiety sensitivity in the relation between pain intensity with substance use and anxiety and depressive symptoms among smokers with chronic pain. *International Journal of Behavioral Medicine*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32588345>

Torres OV, Estep JC, Gwin M, and Villalta I. Gender differences in negative mood, emotional intelligence and tobacco use among young adults. *Substance Use and Misuse*, 2020:1-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32532170>

Smits JAJ, Zvolensky MJ, Otto MW, Piper ME, Baird SO, et al. Enhancing panic and smoking reduction treatment with d-cycloserine: A pilot randomized clinical trial. *Drug and Alcohol Dependence*, 2020; 208:107877. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32004998>

G SC, C RG, M JZ, M ML, N BS, et al. Obsessive-compulsive symptoms and cigarette smoking: An initial cross-sectional test of mechanisms of co-occurrence. *Cogn Behav Ther*, 2020; 49(5):385–97. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32324104>

Ayers CR, Heffner JL, Russ C, Lawrence D, McRae T, et al. Efficacy and safety of pharmacotherapies for smoking cessation in anxiety disorders: Subgroup analysis of the randomized, active- and placebo-controlled eagles trial. *Depress Anxiety*, 2020; 37(3):247–60. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31850603>

Zvolensky MJ, Bakhshaie J, Shepherd JM, Garey L, Peraza N, et al. Anxiety sensitivity and smoking outcome expectancies among spanish-speaking latinx adult smokers. *Experimental and Clinical Psychopharmacology*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30896240>

Watson NL, Heffner JL, Mull KE, McClure JB, and Bricker JB. Comparing treatment acceptability and 12-month cessation rates in response to web-based smoking interventions among smokers who do and do not screen positive for affective disorders: Secondary analysis. *Journal of Medical Internet Research*, 2019; 21(6):e13500. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31219052>

Paulus DJ, Gallagher MW, Raines AM, Schmidt NB, and Zvolensky MJ. Intraindividual change in anxiety sensitivity and alcohol use severity 12-months following smoking cessation treatment. *Behav Res Ther*, 2019; 116:10-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30731131>

Buckner JD, Zvolensky MJ, and Lewis EM. Smoking and social anxiety: The role of false safety behaviors. *Cogn Behav Ther*, 2019:1-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31847703>

Zvolensky MJ, Bakhshaie J, Shepherd JM, Peraza N, Garey L, et al. Anxiety sensitivity and smoking among spanish-speaking latinx smokers. *Addictive Behaviors*, 2018; 90:55-61. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30359848>

Watson NL, DeMarree KG, and Cohen LM. Cigarette craving and stressful social interactions: The roles of state and trait social anxiety and smoking to cope. *Drug and Alcohol Dependence*, 2018; 185:75-81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29428323>

Petroulia I, Kyriakos CN, Papadakis S, Tzavara C, Filippidis FT, et al. Patterns of tobacco use, quit attempts, readiness to quit and self-efficacy among smokers with anxiety or depression: Findings among six countries of the eurest-plus itc europe surveys. *Tobacco Induced Diseases*, 2018; 16:A9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31516463>

Langdon KJ, Bakhshaie J, Lopez A, Tavakoli N, Garey L, et al. Anxiety sensitivity physical and cognitive concerns in relation to smoking-oriented cognition: An examination among treatment-seeking adults who smoke. *Journal of Addiction Medicine*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29438156>

Johnson AL, O'Bryan EM, Kraemer KM, McLeish AC, Zvolensky MJ, et al. The role of anxiety sensitivity-physical concerns in terms of quit day withdrawal symptoms and cravings: A pilot test among smokers with asthma. *The Journal of Asthma*, 2018:1-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29482398>

Guillot CR, Blumenthal H, Zvolensky MJ, and Schmidt NB. Anxiety sensitivity components in relation to alcohol and cannabis use, motives, and problems in treatment-seeking cigarette smokers. *Addictive Behaviors*, 2018; 82:166-73. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29544169>

Zvolensky MJ, Rodriguez-Cano R, Paulus DJ, Kotov R, Bromet E, et al. Respiratory problems and anxiety sensitivity in smoking lapse among treatment seeking smokers. *Addictive Behaviors*, 2017; 75:25-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28666154>

Zvolensky MJ, Bakhshaie J, Norton PJ, Smits JAJ, Buckner JD, et al. Visceral sensitivity, anxiety, and smoking among treatment-seeking smokers. *Addictive Behaviors*, 2017; 75:1-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28654824>

Yang MJ, Zvolensky MJ, and Leyro TM. The indirect effect of panic disorder on smoking cognitions via difficulties in emotion regulation. *Addictive Behaviors*, 2017; 72:126-32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28395248>

Nam E and Fukui S. Tobacco, alcohol, and drug use of people with mood and anxiety disorders: Differential impact of family and friends. *Journal of Dual Diagnosis*, 2017;0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28107107>

Lee JY, Brook JS, Finch SJ, De La Rosa M, and Brook DW. Joint trajectories of cigarette smoking and depressive symptoms from the mid-20s to the mid-30s predicting generalized anxiety disorder. *Journal of Addictive Diseases*, 2017; 36(3):158-66. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28281938>

Farris SG, Uebelacker LA, Brown RA, Price LH, Desaulniers J, et al. Anxiety sensitivity predicts increased perceived exertion during a 1-mile walk test among treatment-seeking smokers. *Journal of Behavioral Medicine*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28451904>

Abrams K, Krimmel S, Johnson S, Cieslowski K, Strnad H, et al. Nicotine deprivation attenuates panic reactivity in smokers: Findings from a placebo-controlled nicotine patch study. *Depress Anxiety*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28489321>

Watson NL, Heffner JL, McClure JB, and Bricker JB. Relationships between social anxiety and smoking-specific experiential avoidance. *Journal of Dual Diagnosis*, 2016;0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27739936>

9A.3.5.3 Suicidal ideation

Park, HJ, Han, B, Kim, B, Han, K, Kim, S, Kim, H et al. (2024). Relationship between smoking experience and risk of suicide mortality in South Korean adults: A nationwide population-based retrospective cohort study. *J Affect Disord*, 367, 67-74. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39222855>

Wang, M, Qin, A, Wei, Z, & Sun, L. (2023). Differentiating the associations between alcohol use, cigarette smoking, and conditional suicidal behaviors among adolescents. *J Affect Disord*, 341, 112-118. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37634822>

Herbst, E, Hoggatt, KJ, & McCaslin, S. (2023). Cigarette Smoking Among Veterans at High Risk for Suicide: Challenges and Opportunities for Intervention. *Mil Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37525945>

Adviento, BA, Regan, EA, Make, BJ, Han, MK, Foreman, MG, Iyer, AS et al. (2022). Clinical Markers Associated with Risk of Suicide or Drug Overdose Among Heavy Smokers - A Longitudinal Follow-Up Study of the COPDGene Cohort. *Chest*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36167120>

9A.3.5.4 Attention-deficit/hyperactivity disorder (ADHD)

McNealy, KR, Weyrich, L, & Bevins, RA. (2023). The co-use of nicotine and prescription psychostimulants: A review of their behavioral and neuropharmacological interactions. *Drug Alcohol Depend*, 248, 109906. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37216808>

- Smethells, JR, Burroughs, D, Saykao, A, & LeSage, MG. (2023). The relative reinforcing efficacy of nicotine in an adolescent rat model of attention-deficit hyperactivity disorder. *Front Psychiatry*, 14, 1154773. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37255676>
- Taylor MR, Carrasco K, Carrasco A, and Basu A. Tobacco and adhd: A role of mao-inhibition in nicotine dependence and alleviation of adhd symptoms. *Front Neurosci*, 2022; 16:845646. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35495050>
- Kollins SH, Sweitzer MM, McClernon FJ, and Perkins KA. Increased subjective and reinforcing effects of initial nicotine exposure in young adults with attention deficit hyperactivity disorder (adhd) compared to matched peers: Results from an experimental model of first-time tobacco use. *Neuropsychopharmacology*, 2020; 45(5):851–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31785588>
- Mitchell JT, McClernon FJ, Beckham JC, Brown RA, Lejuez CW, et al. Smoking abstinence effects on emotion dysregulation in adult cigarette smokers with and without attention-deficit/hyperactivity disorder. *Drug and Alcohol Dependence*, 2019; 205:107594. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31639512>
- Mitchell JT, Howard AL, Belendiuk KA, Kennedy TM, Stehli A, et al. Cigarette smoking progression among young adults diagnosed with adhd in childhood: A 16-year longitudinal study of children with and without adhd. *Nicotine & Tobacco Research*, 2019; 21(5):638–47. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29538764>
- Luo SX, Covey LS, Hu MC, Winhusen TM, and Nunes EV. Differential posttreatment outcomes of methylphenidate for smoking cessation for individuals with adhd. *American Journal on Addictions*, 2019; 28(6):497–502. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31538372>
- Gehricke JG, Gevorkian J, Stehli A, Alejo SD, Dawson M, et al. Discrepancies in the validity of self-reported cigarette smoking in adults with and without adhd. *Journal of Dual Diagnosis*, 2019:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31156069>
- van Amsterdam J, van der Velde B, Schulte M, and van den Brink W. Causal factors of increased smoking in adhd: A systematic review. *Substance Use and Misuse*, 2018; 53(3):432–45. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29039714>
- Luo SX, Wall M, Covey L, Hu MC, Scodes JM, et al. Exploring longitudinal course and treatment-baseline severity interactions in secondary outcomes of smoking cessation treatment in individuals with attention-deficit hyperactivity disorder. *American Journal of Drug and Alcohol Abuse*, 2018; 44(6):653–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29370538>
- Lee HJ, Tran DD, and Morrell HER. Smoking, adhd, and problematic video game use: A structural modeling approach. *Cyberpsychol Behav Soc Netw*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29652524>
- Sweitzer MM, Kollins SH, Kozink RV, Hallyburton M, English J, et al. Adhd, smoking withdrawal and inhibitory control: Results of a neuroimaging study with methylphenidate challenge.

Neuropsychopharmacology, 2017. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/29052617>

Lee CT, McClernon FJ, Kollins SH, and Fuemmeler BF. Childhood adhd symptoms and future illicit drug use: The role of adolescent cigarette use. *J Psychiatr Psychol*, 2017. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/29049706>

Janes AC, Gilman JM, Frederick BB, Radoman M, Pachas G, et al. Salience network coupling is linked to both tobacco smoking and symptoms of attention deficit hyperactivity disorder (adhd). *Drug and Alcohol Dependence*, 2017; 182:93-7. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/29175464>

Galera C, Salla J, Montagni I, Hanne-Poujade S, Salamon R, et al. Stress, attention deficit hyperactivity disorder (adhd) symptoms and tobacco smoking: The i-share study. *European Psychiatry*, 2017; 45:221-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28957791>

Bidwell LC, Karoly HC, Hutchison KE, and Bryan AD. Adhd symptoms impact smoking outcomes and withdrawal in response to varenicline treatment for smoking cessation. *Drug and Alcohol Dependence*, 2017; 179:18–24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28738266>

Bidwell LC, Balestrieri SG, Colby SM, Knopik VS, and Tidey JW. Abstinence-induced withdrawal severity among adolescent smokers with and without adhd: Disentangling effects of nicotine and smoking reinstatement. *Psychopharmacology (Berl)*, 2017. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/29018894>

Rhodes JD, Pelham WE, Gnagy EM, Shiffman S, Derefinko KJ, et al. Cigarette smoking and adhd: An examination of prognostically relevant smoking behaviors among adolescents and young adults. *Psychol Addict Behav*, 2016; 30(5):588-600. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/27824233>

Pagano ME, Delos-Reyes CM, Wasilow S, Svala KM, and Kurtz SP. Smoking cessation and adolescent treatment response with comorbid adhd. *Journal of Substance Abuse Treatment*, 2016; 70:21-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27692184>

9A.3.5.5 Substance use disorders

Nguyen, N, Bold, KW, & McClure, E A. (2024). Urgent need for treatment addressing co-use of tobacco and cannabis: An updated review and considerations for future interventions. *Addict Behav*, 158, 108118. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39089194>

Sedani, AE, Frank-Pearce, SG, Beebe, LA, Campbell, JE, Peck, JD, Chou, AF, & Cohn, AM. (2024). Cannabis Use and Subsequent CIGARETTE Discontinuation among U.S. Adults in the Population Assessment of Tobacco and Health (PATH) Study, Waves 1-5. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39178320>

Erinoso, O, Watts, T, Koning, S, Lu, M, Wagner, K, & Pearson, J. (2024). Choice of smoking cessation products among people with substance use problems in the US: Findings from the Population Assessment of Tobacco and Health (PATH) study Wave 6. *Addict Behav*, 158, 108104. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39042998>

- Lee, SW, Yu, PC, Yen, TT, Hsu, CY, Lai, LJ, Chen, IC, & Chang, T G. (2024). Impact of the Ottawa model on opiate screening and smoking cessation in methadone-treated patients with opioid use disorder: A retrospective cohort analysis. *Tob Induc Dis*, 22. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39076842>
- Bertucci, S, Schultz, E, Stevenson, B, & Fu, SS. (2024). An opt-out tobacco treatment group intervention within an intensive substance use disorders treatment program: Initial outcomes in the Veterans Health Administration. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38818765>
- Fuchshuber, J, Schober, H, Wohlmuth, M, Senra, H, Rominger, C, Schwerdtfeger, A, & Unterrainer, H F. (2024). The effectiveness of a standardized tobacco cessation program on psychophysiological parameters in patients with addiction undergoing long-term rehabilitation: a quasi-experimental pilot study. *BMC Med*, 22(1), 184. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38693570>
- Holt, AG, Hussong, A, Castro, MG, Bossenbroek Fedoriw, K, Schmidt, AM, Prentice, A, & Ware, OD. (2024). Smoking Policies of Outpatient and Residential Substance Use Disorder Treatment Facilities in the United States. *Tob Use Insights*, 17, 1179173X241254803. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38752184>
- McGowan, LJ, John, DA, Kenny, RPW, Joyes, EC, Adams, EA, Shabaninejad, H et al. (2024). Improving oral health and related health behaviours (substance use, smoking, diet) in people with severe and multiple disadvantage: A systematic review of effectiveness and cost-effectiveness of interventions. *PLoS One*, 19(4), e0298885. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38635524>
- Campbell, BK, Le, T, McCuistian, C, Bonniot, C, Delucchi, K, & Guydish, J. (2024). Policy and Practice Changes Associated with a Tobacco-Free Initiative in Residential Substance Use Disorder Treatment. *J Psychoactive Drugs*, 1-10. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38363096>
- Cruvinel, E, Mussulman, L, Scheuermann, T, Shergina, E, He, J, Sherman, S et al. (2024). Hospital-Initiated Smoking Cessation Among Patients Admitted with Behavioral Health Conditions. *J Gen Intern Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38326585>
- Rogova, A, Leal, IM, Britton, M, Chen, TA, Lowenstein, LM, Kyburz, B et al. (2024). Implementing a tobacco-free workplace program at a substance use treatment center: a case study. *BMC Health Serv Res*, 24(1), 201. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38355470>
- West, AL, Correll, L, & Kirschner, JH. (2024). Reducing tobacco and nicotine use among women in treatment for substance use disorder: evaluation of the knit to quit program. *Arts Health*, 1-16. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38420998>
- Neddermann-Carrillo, S, Abidi, L, Gea, A, Bes-Rastrollo, M, de la Fuente-Arrillaga, C, Lahortiga-Ramos, F et al. (2024). Tobacco and alcohol co-use: Lifestyle and sociodemographic factors, and personality aspects as potential predictors in the "Seguimiento Universidad de Navarra" cohort. *Res Nurs Health*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38217468>
- Voci, S, Veldhuizen, S, Ivanova, A, Melamed, OC, Selby, P, & Zawertailo, L. (2024). Cannabis Use Among Adults in Cigarette Smoking Cessation Treatment in Ontario, Canada: Prevalence and

Association With Tobacco Cessation Outcome, 2015-2021. *Am J Public Health*, 114(1), 98-107. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38091559>

Longitudinal Associations Between Use of Tobacco and Cannabis Among People Who Smoke Cigarettes in Real-world Smoking Cessation Treatment: Erratum. (2023). *J Addict Med*, 17(6), 748. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37934555>

Han, K, Wang, T, Shen, F, Li, T, & Zhou, L. (2023). Does smoking associated with increased prescription opioid use and misuse? Evidence from U.S. adults. *Int J Surg*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38000048>

Rakesh, G, Alcorn, JL, 3rd, Khanal, R, Himelhoch, SS, & Rush, CR. (2023). Comparing cigarette-cue attentional bias between people with HIV/AIDS and people with opioid use disorder who smoke. *Health Psychol Behav Med*, 11(1), 2255028. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37693107>

Druckrey-Fiskaaen, KT, Furulund, E, Madebo, T, Carlsen, SL, Fadnes, LT, Lid, TG, & for, A L A R S G. (2023). A qualitative study on people with opioid use disorders' perspectives on smoking and smoking cessation interventions. *Front Psychiatry*, 14, 1185338. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37636821>

Pagano, A, Le, T, Fong, T, Reid, R, Delucchi, K, & Guydish, J. (2023). Gambling, tobacco use, and health among individuals in substance use disorder treatment. *Am J Addict*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37543853>

Guydish, J, McCuistian, C, Hosakote, S, Le, T, Masson, CL, Campbell, BK, & Delucchi, K. (2023). A cluster-randomized trial of a brief multi-component intervention to improve tobacco outcomes in substance use treatment. *Subst Abuse Treat Prev Policy*, 18(1), 34. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37328775>

Jhanjee, S, Charan, D, Mishra, AK, Kaloiya, GS, Jain, R, & Dayal, P. (2023). Effectiveness of smoking cessation intervention in opioid-dependent male subjects on buprenorphine maintenance treatment: An open-label trial. *Ind Psychiatry J*, 32(1), 164-171. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37274578>

Gravely, S, Driezen, P, McClure, EA, Smith, DM, & Fong, GT. (2023). Prevalence of depressive symptoms and cannabis use among adult cigarette smokers in Canada: cross-sectional findings from the 2020 International Tobacco Control Policy Evaluation Project Canada Smoking and Vaping Survey. *CMAJ Open*, 11(3), E516-E526. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37311596>

Kelty, E, Havard, A, & Preen, DB. (2023). Trends in smoking during pregnancy stratified by the use of opioid agonist therapy and the contribution of smoking to poor outcome in neonates prenatally exposed to opioid agonist treatment. *Arch Womens Ment Health*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37368055>

- Harton, MR, Adams, ZW, & Parker, MA. (2023). Associations of alcohol use disorder, cannabis use disorder, and nicotine dependence with concurrent opioid use disorder in U.S. adults. *Exp Clin Psychopharmacol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37166911>
- Peeters, LD, Wills, LJ, Cuzzo, AM, Ivanich, KL, & Brown, RW. (2023). Reinstatement of nicotine conditioned place preference in a transgenerational model of drug abuse vulnerability in psychosis: Impact of BDNF on the saliency of drug associations. *Psychopharmacology (Berl)*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37160431>
- Britton, M, Martinez Leal, I, Jafry, MZ, Chen, TA, Rogova, A, Kyburz, B et al. (2023). Influence of Provider and Leader Perspectives about Concurrent Tobacco-Use Care during Substance-Use Treatment on Their Tobacco Intervention Provision with Clients: A Mixed-Methods Study. *Int J Environ Res Public Health*, 20(7). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37047876>
- Freichel, R, Kroon, E, Kuhns, L, Filbey, F, Veer, IM, Wiers, R, & Cousijn, J. (2023). Cannabis Use Disorder Symptoms in Weekly Cannabis Users: A Network Comparison Between Daily Cigarette Users and Nondaily Cigarette Users. *Cannabis Cannabinoid Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37074121>
- Lobbe, C, Bahnisch, J, Lin, C, Demirkol, A, & Murnion, B. (2023). How do patients and staff in an opioid agonist treatment service view smoking cessation medications and e-cigarettes? *Drug Alcohol Rev*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36877583>
- Martin, EL, Neelon, B, Brady, KT, Guille, C, Baker, NL, Ramakrishnan, V et al. (2023). Differential prevalence of Adverse Childhood Experiences (ACEs) by gender and substance used in individuals with cannabis, cocaine, opioid, and tobacco use disorders. *Am J Drug Alcohol Abuse*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36881810>
- Martinez, C, Lisha, N, McCuistian, C, Strauss, E, Deluchi, K, & Guydish, J. (2023). Comparing client and staff reports on tobacco-related knowledge, attitudes, beliefs and services provided in substance use treatment. *Tob Induc Dis*, 21, 45. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36969981>
- Renton, N, Mwafy, A, Morgan, T, Nicholson, A, & Sherman, S. (2023). Smoking Cessation in Smokers with Alcohol Use Disorder: Does Age Matter? *J Gen Intern Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36941427>
- Shephard, A, Dolek, S, & Barrett, SP. (2023). Investigating predictors of problematic alcohol, cannabis, and nicotine use among legal users of all three substances. *Front Psychiatry*, 14, 1110415. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36911131>
- Walters, KJ, Baker, NL, Tomko, RL, Gray, KM, Carpenter, MJ, & McClure, EA. (2023). Determining the impact of cannabis use and severity on tobacco cessation outcomes: study protocol for a prospective tobacco treatment trial. *BMC Psychol*, 11(1), 25. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36698194>
- Feliu, A, Fernandez, E, Castellano, Y, Enriquez, M, Saura, J, Cabezas, C et al. (2022). Tobacco cessation among smokers under substance use treatment for alcohol and/or cannabis: study protocol and pilot study. *Addict Sci Clin Pract*, 17(1), 66. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36451226>

Chassagne, J, Raynal, P, Bronchain, J, & Chabrol, H. (2022). Smoking Mostly Alone as a Risk Factor for Cannabis Use Disorders and Depressive Symptoms. *Int J Ment Health Addict*, 1-10. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36415673>

Creedon, TB, Wayne, GF, Progovac, AM, Levy, DE, & Cook, BL. (2022). Trends in cigarette use and health insurance coverage among US adults with mental health and substance use disorders. *Addiction*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36385708>

Salameh, TN, Hall, LA, & Hall, MT. (2022). Predictors of cigarette smoking in pregnant women with substance use disorders. *J Addict Dis*, 1-8. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36325923>

Bjornestad, ED, Vederhus, JK, & Clausen, T. (2022). High smoking and low cessation rates among patients in treatment for opioid and other substance use disorders. *BMC Psychiatry*, 22(1), 649. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36261791>

Dobbie, F, Miller, M, Kam, MHM, McKenna, A, Glen, C, & McCallum, A. (2022). DASHES Protocol: Development and Feasibility Testing of a Tailored Community Programme to Support People in Recovery from Problematic Alcohol and Drug Use to Cut Down or Stop Smoking Using Co-Creation. *Int J Environ Res Public Health*, 19(20). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36294287>

Green, R, Lin, J, Montoya, AK, Bello, M S, Grodin, EN, Ryu, H et al. (2022). Characteristics associated with treatment seeking for smoking cessation among heavy-drinking research participants. *Front Psychiatry*, 13, 951364. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36245856>

Salameh, TN, Hall, LA, & Hall, MT. (2022). Cigarette Smoking Cessation Counselling in Pregnant Smokers with Mental Illness/Substance Use Disorders. *West J Nurs Res*, 1939459221127803. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36196024>

Kathuria, H, Shankar, D, Cobb, V, Newman, J, Bulekova, K, Werntz, S, & Borrelli, B. (2022). Integrating Social Determinants of Health With Tobacco Treatment for Individuals With Opioid Use Disorder: Feasibility and Acceptability Study of Delivery Through Text Messaging. *JMIR Form Res*, 6(9), e36919. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36048509>

Kolodner, G, DiClemente, CC, & Miller, MM. (2022). Nicotine Addiction: A Burning Issue in Addiction Psychiatry. *Psychiatr Clin North Am*, 45(3), 451-465. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36055732>

Druckrey-Fiskaaen, KT, Furulund, E, Daltveit, JT, Vold, JH, Lid, TG, Madebo, T et al. (2022). Integration of smoking cessation into standard treatment for patients receiving opioid agonist therapy who are smoking tobacco: protocol for a randomised controlled trial (ATLAS4LAR). *Trials*, 23(1), 663. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35978355>

Matthews, H, Diamond, JB, Morrison, D, Teitelbaum, SA, & Merlo, LJ. (2022). Patient experiences with tobacco use during substance use disorder treatment and early recovery: a mixed method analysis of phone interview responses. *J Addict Dis*, 1-7. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35930400>

Garnett, C, Oldham, M, Shahab, L, Tattan-Birch, H, & Cox, S. (2022). Characterising smoking and smoking cessation attempts by risk of alcohol dependence: A representative, cross-sectional study of

adults in England between 2014-2021. *Lancet Reg Health Eur*, 18, 100418. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35814338>

Lima, DR, Guimaraes-Pereira, BBS, Mannes, ZL, Carvalho, CFC, Loreto, AR, Davanso, LC et al. (2022). The effect of a real-world intervention for smoking cessation in Adults with and without comorbid psychiatric and substance use disorders: A one-year follow-up study. *Psychiatry Res*, 315, 114722. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35841703>

Min, JY, Levin, J, & Weinberger, AH. (2022). Associations of tobacco cigarette use and dependence with substance use disorder treatment completion by sex/gender and race/ethnicity. *J Subst Abuse Treat*, 140, 108834. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35803029>

Pagano, A, McCuistian, C, Le, T, Campbell, BK, Delucchi, K, Woodward-Lopez, G, & Guydish, J. (2022). Smoking Behavior and Wellness among Individuals in Substance Use Disorder Treatment. *J Psychoactive Drugs*, 1-12. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35815722>

Secades-Villa, R, Aonso-Diego, G, & Gonzalez-Roz, A. (2022). A randomized controlled trial of contingency management for smoking cessation in substance use treatment patients. *Int J Clin Health Psychol*, 22(3), 100314. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35662791>

Swithenbank, Z, Harrison, R, & Porcellato, L. (2022). Service user perceptions of smoking cessation in residential substance use treatment. *PLoS One*, 17(6), e0270045. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35731791>

van Amsterdam J and van den Brink W. Smoking as an outcome moderator in the treatment of alcohol use disorders. *Alcohol Alcohol*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35589093>

Patel SY, Wayne GF, Progovac AM, Flores M, Moyer M, et al. Effects of medicaid coverage on receipt of tobacco dependence treatment among medicaid beneficiaries with substance use disorder. *Health Serv Res*, 2022. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35584242>

Glasser AM, Nemeth JM, Quisenberry AJ, Shoben AB, Trapl ES, et al. Cigarillo flavor and motivation to quit among co-users of cigarillos and cannabis: A structural equation modeling approach. *International Journal of Environmental Research and Public Health*, 2022; 19(9). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35565122>

Fokuo JK, McCuistian CL, Masson CL, Gruber VA, Straus E, et al. Pre-implementation assessment of tobacco cessation interventions in substance use disorder residential programs in california. *Substance Use and Misuse*, 2022:1-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35621319>

Shankar D, Borrelli B, Cobb V, Quintiliani LM, Palfai T, et al. Text-messaging to promote smoking cessation among individuals with opioid use disorder: Quantitative and qualitative evaluation. *BMC Public Health*, 2022; 22(1):668. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35387648>

Garcia-Perez A, Aonso-Diego G, Weidberg S, and Secades-Villa R. Effects of episodic future thinking on reinforcement pathology during smoking cessation treatment among individuals with substance

use disorders. *Psychopharmacology (Berl)*, 2022; 239(2):631-42. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35020047>

Aonso-Diego G, Gonzalez-Roz A, Weidberg S, Garcia-Fernandez G, and Secades-Villa R. Smoking cessation treatment attendance among smokers with substance use disorders. *Adicciones*, 2022; 0(0):1718. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35472160>

Zvolensky MJ, Rogers AH, Garey L, Shepherd JM, and Ditre JW. Opioid misuse among smokers with chronic pain: Relations with substance use and mental health. *Behav Med*, 2021; 47(4):335-43. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34719341>

Weinberger AH, Platt J, Zhu J, Levin J, Ganz O, et al. Cigarette use and cannabis use disorder onset, persistence, and relapse: Longitudinal data from a representative sample of US adults. *Journal of Clinical Psychiatry*, 2021; 82(4). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34232581>

Taing M, Nitturi V, Chen TA, Kyburz B, Martinez Leal I, et al. Implementation and outcomes of a comprehensive tobacco free workplace program in opioid treatment centers. *International Journal of Environmental Research and Public Health*, 2021; 19(1). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35010499>

Streck JM, Kalkhoran S, Bearnot B, Gupta PS, Kalagher KM, et al. Perceived risk, attitudes, and behavior of cigarette smokers and nicotine vapers receiving buprenorphine treatment for opioid use disorder during the COVID-19 pandemic. *Drug and Alcohol Dependence*, 2021; 218:108438. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33271434>

Siegel SD, Brooks M, Ragozine-Bush HE, Schnoll RA, and Curriero FC. The co-occurrence of smoking and alcohol use disorder in a hospital-based population: Applying a multimorbidity framework using geographic information system methods. *Addictive Behaviors*, 2021; 118:106883. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33714034>

Selva Kumar D, Peterson M, Zhang C, Fagan P, and Nahvi S. The impact of menthol cigarette use on quit attempts and abstinence among smokers with opioid use disorder. *Addictive Behaviors*, 2021; 118:106880. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33706070>

Rodriguez-Cano R, Paulus DJ, Derrick JL, Blalock JA, and Zvolensky MJ. Emotion dysregulation and hazardous drinking in relation to suicidal ideation among spanish-speaking latinx daily-smokers. *Journal of Substance Abuse Treatment*, 2021; 132:108508. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34098209>

Reed ZE, Wootton RE, and Munafo MR. Using mendelian randomization to explore the gateway hypothesis: Possible causal effects of smoking initiation and alcohol consumption on substance use outcomes. *Addiction*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34590374>

Parker MA, Cordoba-Grueso WS, Streck JM, Goodwin RD, and Weinberger AH. Intersectionality of serious psychological distress, cigarette smoking, and substance use disorders in the United States: 2008-2018. *Drug and Alcohol Dependence*, 2021; 228:109095. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34601273>

Nieva G, Ballbe M, Cano M, Carcole B, Fernandez T, et al. Smoking cessation interventions in substance use disorders treatment centers of catalonia: The abandoned addiction. *Adicciones*, 2021; 0(0):1492. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33768256>

Nieto SJ, Venegas A, Burnette EM, MacKillop J, and Ray LA. Additive roles of tobacco and cannabis co-use in relation to delay discounting in a sample of heavy drinkers. *Psychopharmacology (Berl)*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34652499>

Nguyen N, Neilands TB, Lisha NE, Lyu JC, Olson SS, et al. Longitudinal associations between use of tobacco and cannabis among people who smoke cigarettes in real-world smoking cessation treatment. *Journal of Addiction Medicine*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34619713>

Naude GP, Reed DD, Jarmolowicz DP, Martin LE, Fox AT, et al. Single- and cross-commodity discounting among adults who use alcohol and cannabis: Associations with tobacco use and clinical indicators. *Drug and Alcohol Dependence*, 2021; 229(Pt B):109082. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34634563>

Motschman CA and Tiffany ST. Combined smoking and alcohol cues: Effects on craving, drug-seeking, and consumption. *Alcohol Clin Exp Res*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34469584>

Montgomery L, Winhusen T, Scodes J, Pavlicova M, Twitty D, et al. Reductions in tobacco use in naltrexone, relative to buprenorphine-maintained individuals with opioid use disorder: Secondary analysis from the national drug abuse treatment clinical trials network. *Journal of Substance Abuse Treatment*, 2021; 130:108489. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34118716>

Minian N, Noormohamed A, Lingam M, Zawertailo L, Le Foll B, et al. Integrating a brief alcohol intervention with tobacco addiction treatment in primary care: Qualitative study of health care practitioner perceptions. *Addiction Science & Clinical Practice*, 2021; 16(1):17. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33726843>

McClure JB and Lapham G. Tobacco quitline engagement and outcomes among primary care patients reporting use of tobacco or dual tobacco and cannabis: An observational study. *Substance Abuse*, 2021; 42(4):417–22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33270541>

Lien L, Bolstad I, and Bramness JG. Smoking among inpatients in treatment for substance use disorders: Prevalence and effect on mental health and quality of life. *BMC Psychiatry*, 2021; 21(1):244. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33975577>

Lee J and Thrul J. Trends in opioid misuse by cigarette smoking status among US adolescents: Results from national survey on drug use and health 2015-2018. *Preventive Medicine*, 2021; 153:106829. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34624387>

Kwon I, Montebello M, and Bittoun R. Tobacco dependence management in a smoke-free inpatient drug and alcohol unit. *Australas Psychiatry*, 2021; 29(1):14–21. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33301381>

Kunas SL, Stuke H, Heinz A, Strohle A, and BERPPOHL F. Evidence for a hijacked brain reward system but no desensitized threat system in quitting motivated smokers: An fmri study. *Addiction*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34312937>

King A and Fucito L. Cigarette smoking and heavy alcohol drinking: The challenges and opportunities for combination treatments. *American Journal of Psychiatry*, 2021; 178(9):783-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34516230>

Ho RKS, Fok PWY, and Chan HCH. Pattern and determinants of alcohol and tobacco co-use and its relationship with smoking cessation in hong kong. *Tob Prev Cessat*, 2021; 7:21. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33778215>

Haeny AM, Gueorguieva R, Montgomery L, Bold KW, Fucito LM, et al. The effect of varenicline on smoking and drinking outcomes among black and white adults with alcohol use disorder and co-occurring cigarette smoking: A secondary analysis of two clinical trials. *Addictive Behaviors*, 2021; 122:106970. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34216871>

Guillaumier A, Skelton E, Tzelepis F, D'Este C, Paul C, et al. Patterns and predictors of nicotine replacement therapy use among alcohol and other drug clients enrolled in a smoking cessation randomised controlled trial. *Addictive Behaviors*, 2021; 119:106935. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33848758>

Goodwin RD, Wu M, and Davidson L. An empirical perspective on cigarette use in substance use recovery. *Psychological Medicine*, 2021:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34261564>

Gonzalez M and Sanders-Jackson A. Traditional cigarette and poly-tobacco use are associated with medical opioid use in rural areas of the US. *International Journal of Environmental Research and Public Health*, 2021; 18(22). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34831619>

Felicione NJ, Ozga JE, Dino G, Berry JH, Sullivan CR, et al. Timing of smoking cessation treatment integrated into outpatient treatment with medications for opioid use disorder: Feasibility trial. *Journal of Substance Abuse Treatment*, 2021:108579. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34452780>

Eastwood B, Clare T, Dockrell MJ, Locker J, Chowdary Q, et al. Reciprocal influences of tobacco use on illicit opioid and alcohol use during the first six-months of specialist addiction treatment. *Drug and Alcohol Dependence*, 2021; 218:108418. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33262003>

Cui Y, Linares Abrego P, Yoon JH, Karam-Hage M, Cinciripini PM, et al. Behavioral economic assessment of alcohol and cigarette demand in smokers with alcohol use disorder. *Front Psychiatry*, 2021; 12:674607. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34248711>

Courtney JB, Russell MA, and Conroy DE. Tobacco and cannabis use as moderators of the association between physical activity and alcohol use across the adult lifespan in the United States: Nhanes, 2005-2016. *Preventive Medicine*, 2021; 155:106931. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34954238>

Australian Institute of Health and Welfare. Alcohol and other drug treatment services in Australia: Key findings. Cat. no. HSE 242. Canberra: AIHW, 2021. Available from: <https://www.aihw.gov.au/getmedia/373321cd-bcef-473c-95b0-56b3ee55c3bb/Alcohol-and-other-drug-treatment-services-in-Australia.pdf.aspx?inline=true>.

Aonso-Diego G, Gonzalez-Roz A, Martinez-Loredo V, Krotter A, and Secades-Villa R. Episodic future thinking for smoking cessation in individuals with substance use disorder: Treatment feasibility and acceptability. *Journal of Substance Abuse Treatment*, 2021; 123:108259. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33612193>

Alkan O, Abar H, and Gencer O. Analysis of factors affecting alcohol and tobacco concurrent use by bivariate probit model in turkey. *Environ Sci Pollut Res Int*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33586099>

Alghamdi AA, Gartner C, Najman J, and Smirnov A. Co-administering cannabis with tobacco and persistent tobacco smoking after 4(1/2) years in young adults who use stimulants: A prospective population-based study. *J Psychoactive Drugs*, 2021:1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34254884>

Ahmed Alghamdi A, Gartner C, Najman JM, Chan G, and Smirnov A. The relationship between cannabis and tobacco co-administration and long-term patterns of cannabis use in young adults who use stimulants: A prospective population-based study. *Substance Use and Misuse*, 2021:1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34796786>

Zawertailo L, Ivanova A, Ng G, Le Foll B, and Selby P. Safety and efficacy of varenicline for smoking cessation in alcohol-dependent smokers in concurrent treatment for alcohol use disorder: A pilot, randomized placebo-controlled trial. *J Clin Psychopharmacol*, 2020; 40(2):130–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32068562>

Yoon JH, Suchting R, Cassidy RN, Bolin PK, Omar Y, et al. Assessment of demand for methamphetamine and cigarettes among individuals with methamphetamine use disorder. *Experimental and Clinical Psychopharmacology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32202813>

Walsh H, McNeill A, Purssell E, and Duaso M. A systematic review and bayesian meta-analysis of interventions which target or assess co-use of tobacco and cannabis in single- or multi-substance interventions. *Addiction*, 2020; 115(10):1800–14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32003088>

Vlad C, Arnsten JH, and Nahvi S. Achieving smoking cessation among persons with opioid use disorder. *CNS Drugs*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32107731>

Thrul J, Gubner NR, Nguyen N, Nguyen C, Goodell EA, et al. Perceived reward from using cigarettes with alcohol or cannabis and concurrent use: A smartphone-based daily diary study. *Addictive Behaviors*, 2020; 114:106747. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33307406>

Streck JM, Sigmon SC, Priest J, Bergeria CL, Davis DR, et al. Investigating tobacco withdrawal in response to reduced nicotine cigarettes among smokers with opioid use disorder and other

vulnerabilities. *Experimental and Clinical Psychopharmacology*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32027158>

Streck JM, Parker MA, Weinberger AH, Rigotti NA, and Park ER. Association of cigarette use and substance use disorders among US adults with and without a recent diagnosis of cancer. *Curr Oncol*, 2020; 28(1):86-93. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33704178>

Rogers AH, Shepherd JM, Buckner JD, Garey L, Manning K, et al. Current cannabis use and smoking cessation among treatment seeking combustible smokers. *Drug and Alcohol Dependence*, 2020; 209:107928. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32092636>

Rhodes BE and Gottfredson NC. Effects of tobacco on affect and craving during opioid addiction recovery: An ecological momentary assessment study. *Addictive Behaviors*, 2020; 106:106358. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32151893>

Rabin RA. Commentary on walsh et al. (2020): Tobacco and cannabis co-use- considerations for treatment. *Addiction*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32293768>

Pericot-Valverde I, Heo M, Akiyama MJ, Norton BL, Agyemang L, et al. Factors and hcv treatment outcomes associated with smoking among people who inject drugs on opioid agonist treatment: Secondary analysis of the prevail randomized clinical trial. *BMC Infect Dis*, 2020; 20(1):928. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33276738>

Parker MA, Weinberger AH, and Villanti AC. Quit ratios for cigarette smoking among individuals with opioid misuse and opioid use disorder in the United States. *Drug and Alcohol Dependence*, 2020; 214:108164. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32652375>

Nguyen N, Nguyen C, and Thrul J. Digital health for assessment and intervention targeting tobacco and cannabis co-use. *Curr Addict Rep*, 2020; 7(3):268–79. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33643768>

Moey CH, Yee A, and Muhamud Kayat SB. Tobacco use disorder: Prevalence, associated factors and its influence on quality of life among patients on methadone assisted treatment. *Journal of Addictive Diseases*, 2020:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32329412>

Minian N, Ivanova A, Voci S, Veldhuizen S, Zawertailo L, et al. Computerized clinical decision support system for prompting brief alcohol interventions with treatment seeking smokers: A sex-based secondary analysis of a cluster randomized trial. *International Journal of Environmental Research and Public Health*, 2020; 17(3). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32041190>

Meadows AL, Guenthner G, Broster LS, Luo AY, and Martin CA. Assessing adolescent tobacco use in a substance use treatment program with self-report and biological measures. *Addictive Behaviors*, 2020; 107:106424. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32251874>

McHugh RK, Janes AC, Griffin ML, Taghian N, Greenfield SF, et al. Clinical correlates of smoking status in men and women with opioid use disorder. *Substance Use and Misuse*, 2020:1-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32251874>

Im.nih.gov/pubmed/32037945 McClure EA, Rabin RA, Lee DC, and Hindocha C. Treatment implications associated with cannabis and tobacco co-use. *Curr Addict Rep*, 2020; 7(4):533-44. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33777645>

Lima DR, Carvalho CFC, Guimaraes-Pereira BBS, Loreto AR, Frallonardo FP, et al. Abstinence and retention outcomes in a smoking cessation program among individuals with co-morbid substance use and mental disorders. *Journal of Psychiatric Research*, 2020; 125:121-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32272242>

Kranzler HR, Washio Y, Zindel LR, Lynch KG, Hand D, et al. Pregnant smokers receiving opioid agonist therapy have an elevated nicotine metabolite ratio: A replication study. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32303761>

Kaye JT, Johnson AL, Baker TB, Piper ME, and Cook JW. Searching for personalized medicine for binge drinking smokers: Smoking cessation using varenicline, nicotine patch, or combination nicotine replacement therapy. *J Stud Alcohol Drugs*, 2020; 81(4):426-35. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32800078>

Kahler CW, Cohn AM, Costantino C, Toll BA, Spillane NS, et al. Digital smoking cessation for heavy drinkers: Pilot randomized controlled trial. *JMIR Form Res*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32348286>

Kadri AN, Khodor S, Ali A, Nusairat L, Mahmood A, et al. National trends of tobacco, alcohol, and drug use in patients admitted with acute myocardial infarction. *Cardiovasc Revasc Med*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33229134>

Hobden B, Bryant J, Forshaw K, Oldmeadow C, Evans TJ, et al. Prevalence and characteristics associated with concurrent smoking and alcohol misuse within Australian general practice patients. *Aust Health Rev*, 2020; 44(1):125–31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30543764>

Hindocha C, Brose LS, Walsh H, and Cheeseman H. Cannabis use and co-use in tobacco smokers and non-smokers: Prevalence and associations with mental health in a cross-sectional, nationally representative sample of adults in Great Britain, 2020. *Addiction*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33345423>

Hallford G, Bakhshaie J, Rodriguez-Cano R, Shepherd JM, Schmidt NB, et al. A multiple mediation model of trait worry and hazardous drinking among treatment-seeking smokers: The roles of coping and conformity drinking motives. *Cogn Behav Ther*, 2020:1-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32352348>

Guillaumier A, Skelton E, Shakeshaft A, Farrell M, Tzelepis F, et al. Effect of increasing the delivery of smoking cessation care in alcohol and other drug treatment centres: A cluster-randomized controlled trial. *Addiction*, 2020; 115(7):1345–55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31762105>

Green R, MacKillop J, Hartwell EE, Lim AC, Baskerville WA, et al. Behavioral economic demand for alcohol and cigarettes in heavy drinking smokers: Evidence of asymmetric cross-commodity

reinforcing value. Nicotine & Tobacco Research, 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33247757>

Fucito LM, Wu R, O'Malley SS, Hanrahan TH, Ikomi JT, et al. An integrated behavioural intervention combined with varenicline for heavy-drinking smokers: A randomized pilot study. J Smok Cessat, 2020; 15(3):119-27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33312238>

Durazzo TC and Meyerhoff DJ. Cigarette smoking history is associated with poorer recovery in multiple neurocognitive domains following treatment for an alcohol use disorder. Alcohol, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31923562>

Cheatle MD, Falcone M, Dhingra L, and Lerman C. Independent association of tobacco use with opioid use disorder in patients of European ancestry with chronic non-cancer pain. Drug and Alcohol Dependence, 2020; 209:107901. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32126454>

Campbell BK, Le T, Kapiteni K, Gubner NR, and Guydish J. Correlates of lifetime blunt/spliff use among cigarette smokers in substance use disorders treatment. Journal of Substance Abuse Treatment, 2020; 116:108064. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32741500>

Antonelli M, Fattore L, Sestito L, Di Giuda D, Diana M, et al. Transcranial magnetic stimulation: A review about its efficacy in the treatment of alcohol, tobacco and cocaine addiction. Addictive Behaviors, 2020; 114:106760. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33316590>

Akhtar WZ, Mundt MP, Koepke R, Krechel S, Fiore MC, et al. Prevalence of tobacco use among rural-dwelling individuals who inject drugs. JAMA Netw Open, 2020; 3(3):e200493. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32154885>

Zhong BL, Xu YM, Xie WX, and Lu J. Cigarette smoking is significantly linked to sexual dissatisfaction in chinese heroin-dependent patients receiving methadone maintenance treatment. Front Psychiatry, 2019; 10:306. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31178762>

Yip D, Gubner N, Le T, Williams D, Delucchi K, et al. Association of medicaid expansion and health insurance with receipt of smoking cessation services and smoking behaviors in substance use disorder treatment. J Behav Health Serv Res, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31359228>

Wong HY, Subramaniyan M, Bullen C, Amer Siddiq AN, Danaee M, et al. The mobile-phone-based ico(tm) smokerlyzer((r)): Comparison with the pico(+) smokerlyzer((r)) among smokers undergoing methadone-maintained therapy. Tobacco Induced Diseases, 2019; 17:65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31582954>

Winhusen TM, Theobald J, and Lewis DF. Substance use outcomes in cocaine-dependent tobacco smokers: A mediation analysis exploring the role of sleep disturbance, craving, anxiety, and depression. Journal of Substance Abuse Treatment, 2019; 96:53-7. Available from: [https://www.journalofsubstanceabusetreatment.com/article/S0740-5472\(18\)30413-6/fulltext](https://www.journalofsubstanceabusetreatment.com/article/S0740-5472(18)30413-6/fulltext)

Webb Hooper M. Racial/ethnic differences in physiological stress and relapse among treatment seeking tobacco smokers. International Journal of Environmental Research and Public Health, 2019; 16(17). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31450707>

Wang R, Li B, Jiang Y, Guan Y, Wang G, et al. Smoking cessation mutually facilitates alcohol drinking cessation among tobacco and alcohol co-users: A cross-sectional study in a rural area of shanghai, China. *Tobacco Induced Diseases*, 2019; 17:85. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31889947>

Veliz P, Eisman A, McCabe SE, Evans-Polce R, McCabe VV, et al. E-cigarette use, polytobacco use, and longitudinal changes in tobacco and substance use disorder symptoms among u.S. Adolescents. *The Journal of Adolescent Health*, 2019. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31395513>

Romberg AR, Miller Lo EJ, Barton A, Xiao H, Vallone D, et al. Cigarette smoking, prescription opioid use and misuse among young adults: An exploratory analysis. *Preventive Medicine*, 2019:105845.

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31518628>

Rajabi A, Dehghani M, Shojaei A, Farjam M, and Motevalian SA. Association between tobacco smoking and opioid use: A meta-analysis. *Addictive Behaviors*, 2019; 92:225–35. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30685521>

Quisenberry AJ, Pittman J, Goodwin RD, Bickel WK, D'Urso G, et al. Smoking relapse risk is increased among individuals in recovery. *Drug and Alcohol Dependence*, 2019; 202:93-103. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31325822>

Peltzer K, Chao LW, Ramlagan S, and Szrek H. Daily tobacco use and problem drinking among urban adults in south africa: A longitudinal study. *Pan Afr Med J*, 2019; 32:51. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31143356>

Park SH. Smoking-related differential influence of alcohol consumption on the metabolic syndrome. *Substance Use and Misuse*, 2019:1-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31373528>

Pacek LR, Reboussin BA, Green KM, LaFlair LN, Storr CL, et al. Current tobacco use, nicotine dependence, and transitions across stages of alcohol involvement: A latent transition analysis approach. *Int J Methods Psychiatr Res*, 2019:e1789. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31141253>

Minian N, Baliunas D, Noormohamed A, Zawertailo L, Giesbrecht N, et al. The effect of a clinical decision support system on prompting an intervention for risky alcohol use in a primary care smoking cessation program: A cluster randomized trial. *Implement Sci*, 2019; 14(1):85. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31443663>

McCrabb S, Baker AL, Attia J, Balogh ZJ, Lott N, et al. Comorbid tobacco and other substance use and symptoms of anxiety and depression among hospitalised orthopaedic trauma patients. *BMC Psychiatry*, 2019; 19(1):28. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30654783>

McClure EA, Baker NL, Hood CO, Tomko RL, Squeglia LM, et al. Cannabis and alcohol co-use in a smoking cessation pharmacotherapy trial for adolescents and emerging adults. *Nicotine & Tobacco Research*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31612956>

Martin RA, Rohsenow DJ, and Tidey JW. Smokers with opioid use disorder may have worse drug use outcomes after varenicline than nicotine replacement. *Journal of Substance Abuse Treatment*, 2019; 104:22-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31370981>

Lee DC, Walker DD, Hughes JR, Brunette MF, Scherer E, et al. Sequential and simultaneous treatment approaches to cannabis use disorder and tobacco use. *Journal of Substance Abuse Treatment*, 2019; 98:39-46. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30665602>

Kladney M, Joudrey P, Cunningham CO, and Bachhuber MA. Pharmacotherapy prescribing to patients with concurrent tobacco and alcohol use disorder in a large, urban, integrated health system. *Journal of General Internal Medicine*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30604121>

Kelly PJ, Baker AL, Townsend CJ, Deane FP, Callister R, et al. Healthy recovery: A pilot study of a smoking and other health behavior change intervention for people attending residential alcohol and other substance dependence treatment. *Journal of Dual Diagnosis*, 2019; 15(3):207–16. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31122158>

Kelly JG, C; Bergman, B; Hoepfner, B. Smoking cessation in the context of recovery from drug and alcohol problems: Prevalence, predictors, and cohort effects in a national u.S. Sample Drug and Alcohol Dependence, 2019; 195:6-12. Available from: <https://www-clinicalkey-com-au.ezp.lib.unimelb.edu.au/#!/content/journal/1-s2.0-S0376871618308275?scrollTo=%23top>

John WS, Zhu H, Mannelli P, Subramaniam GA, Schwartz RP, et al. Prevalence and patterns of opioid misuse and opioid use disorder among primary care patients who use tobacco. *Drug and Alcohol Dependence*, 2019; 194:468-75. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30513477>

Hovhannisyan K, Gunther M, Raffing R, Wikstrom M, Adami J, et al. Compliance with the very integrated program (vip) for smoking cessation, nutrition, physical activity and comorbidity education among patients in treatment for alcohol and drug addiction. *International Journal of Environmental Research and Public Health*, 2019; 16(13). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31261620>

Guydish J, Yip D, Le T, Gubner NR, Williams DD, et al. Tobacco cessation services in addiction treatment: What do clients say? *Psychiatric Services*, 2019; appips201700565. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30630403>

Green R, Bujarski S, Lim AC, Venegas A, and Ray LA. Naltrexone and alcohol effects on craving for cigarettes in heavy drinking smokers. *Experimental and Clinical Psychopharmacology*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30628813>

Gonzalez-Roz A, Ruano L, Aonso-Diego G, Garcia-Perez A, Weidberg S, et al. Smoking cessation interventions in substance use treatment facilities: Clinical implications and recommendations for implementation. *Adicciones*, 2019; 0(0):1270. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31342079>

Gonzalez-Roz A, Postigo A, Aonso-Diego G, Garcia-Perez A, and Secades-Villa R. Reinforcer pathology among cigarette smokers with and without history of alcohol use disorder. *Psicothema*, 2019; 31(4):393-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31634083>

Fine DR, Bearnot BI, Rigotti NA, and Baggett TP. Smoking status and quit behaviors among health center patients with substance use disorders: A national study. *Drug and Alcohol Dependence*, 2019; 202:6-12. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31279257>

Fallin-Bennett A, Rademacher K, Dye H, Elswick A, Ashford K, et al. Perinatal navigator approach to smoking cessation for women with prevalent opioid dependence. *West J Nurs Res*, 2019;193945918825381. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30724661>

Cooperman NA, Rizvi SL, Hughes CD, and Williams JM. Field test of a dialectical behavior therapy skills training-based intervention for smoking cessation and opioid relapse prevention in methadone treatment. *Journal of Dual Diagnosis*, 2019:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30646819>

Bold KW, Rosen RL, Steinberg ML, Epstein EE, McCrady BS, et al. Smoking characteristics and alcohol use among women in treatment for alcohol use disorder. *Addictive Behaviors*, 2019; 101:106137. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31648138>

Boatner J, Patterson D, Herron F, Nugent W, and Rice T. Assessing the availability of pharmacotherapy options for tobacco cessation in tennessee's substance use facilities. *South Med J*, 2019; 112(7):406-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31282972>

Baldassarri SR, Fiellin DA, Savage ME, Madden LM, Beitel M, et al. Electronic cigarette and tobacco use in individuals entering methadone or buprenorphine treatment. *Drug and Alcohol Dependence*, 2019; 197:37-41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30769264>

Atzendorf J, Rauschert C, Seitz NN, Lochbuhler K, and Kraus L. The use of alcohol, tobacco, illegal drugs and medicines. *Dtsch Arztebl Int*, 2019; 116(35-36):577-84. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31587705>

Amorim TA, Lucchese R, Silva Neta EMD, Santos JSD, Vera I, et al. Determinants of mental health and abuse of psychoactive substances associated with tobacco use. A case-control study. *Cien Saude Colet*, 2019; 24(11):4141-52. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31664387>

Yee A, Hoong MC, Joyce YC, and Loh HS. Smoking cessation among methadone-maintained patients: A meta-analysis. *Substance Use and Misuse*, 2018; 53(2):276–85. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28857640>

Worley MJ, Isgro M, Heffner JL, Lee SY, Daniel BE, et al. Predictors of reduced smoking quantity among recovering alcohol dependent men in a smoking cessation trial. *Addictive Behaviors*, 2018; 84:263-70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29763835>

Weinberger AH, Pacek LR, Wall MM, Zvolensky MJ, Copeland J, et al. Trends in cannabis use disorder by cigarette smoking status in the United States, 2002-2016. *Drug and Alcohol Dependence*, 2018; 191:45–51. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30077055>

Weinberger AH, Pacek LR, Giovenco D, Galea S, Zvolensky MJ, et al. Cigarette use among individuals with alcohol use disorders in the United States, 2002-2016: Trends overall and by race/ethnicity. *Alcohol Clin Exp Res*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30408209>

Weinberger AH, Gbedemah M, Wall MM, Hasin DS, Zvolensky MJ, et al. Cigarette use is increasing among people with illicit substance use disorders in the United States, 2002-14: Emerging disparities in vulnerable populations. *Addiction*, 2018; 113(4):719–28. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29265574>

Vogel EA, Rubinstein ML, Prochaska JJ, and Ramo DE. Associations between marijuana use and tobacco cessation outcomes in young adults. *Journal of Substance Abuse Treatment*, 2018; 94:69-73. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30243420>

Streck JM, Heil SH, Higgins ST, Bunn JY, and Sigmon SC. Tobacco withdrawal among opioid-dependent smokers. *Experimental and Clinical Psychopharmacology*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29389213>

Skelton E, Tzelepis F, Shakeshaft A, Guillaumier A, Wood W, et al. Integrating smoking cessation care into routine service delivery in a medically supervised injecting facility: An acceptability study. *Addictive Behaviors*, 2018; 84:193–200. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29723802>

Serre F, Fatseas M, Denis C, Swendsen J, and Auriacombe M. Predictors of craving and substance use among patients with alcohol, tobacco, cannabis or opiate addictions: Commonalities and specificities across substances. *Addictive Behaviors*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29428330>

Roberts W, Shi JM, Tetraault JM, and McKee SA. Effects of varenicline alone and in combination with low-dose naltrexone on alcohol-primed smoking in heavy-drinking tobacco users: A preliminary laboratory study. *Journal of Addiction Medicine*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29438157>

Roberts W, Ralevski E, Verplaetse TL, McKee SA, and Petrakis IL. Tobacco use during a clinical trial of mecamylamine for alcohol dependence: Medication effects on smoking and associations with reductions in drinking. *Journal of Substance Abuse Treatment*, 2018; 94:91-6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30243424>

Raich A, Pinet C, Ballbe M, Mondon S, Tejedor R, et al. Multimodal treatment for smoking cessation with varenicline in alcoholic, methadone-maintained, and psychotic patients: A one-year follow-up. *Tobacco Induced Diseases*, 2018; 16:58. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31516455>

Parker MA, Streck JM, and Sigmon SC. Associations between opioid and nicotine dependence in nationally representative samples of United States adult daily smokers. *Drug and Alcohol Dependence*, 2018; 186:167-70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29579724>

Pagano A, Gubner NR, Le T, Yip D, Williams D, et al. Differences in tobacco use prevalence, behaviors, and cessation services by race/ethnicity: A survey of persons in addiction treatment.

Journal of Substance Abuse Treatment, 2018; 94:9-17. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30243423>

Pagano A, Gubner N, Le T, and Guydish J. Cigarette smoking and quit attempts among latinos in substance use disorder treatment. American Journal of Drug and Alcohol Abuse, 2018;1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29333890>

O'Malley SS, Zweben A, Fucito LM, Wu R, Piepmeier ME, et al. Effect of varenicline combined with medical management on alcohol use disorder with comorbid cigarette smoking: A randomized clinical trial. JAMA Psychiatry, 2018; 75(2):129–38. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29261824>

Nahvi S and Arnsten JH. Missed opportunities to test the neuropsychiatric safety--and efficacy--of varenicline among smokers with substance use disorders. Drug and Alcohol Dependence, 2018; 185:245-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30369710>

Minian N, Noormohamed A, Zawertailo L, Baliunas D, Giesbrecht N, et al. A method for co-creation of an evidence-based patient workbook to address alcohol use when quitting smoking in primary care: A case study. Res Involv Engagem, 2018; 4:4. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29445521>

Minami H, Bloom EL, Brinkman HR, Abrantes AM, Young CC, et al. Factors related to cigarette smoking and intent to quit among adolescent inpatients with psychiatric and substance use disorders. Drug and Alcohol Dependence, 2018; 186:215-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29609133>

McPherson S, Orr M, Lederhos C, McDonell M, Leickly E, et al. Decreases in smoking during treatment for methamphetamine-use disorders: Preliminary evidence. Behav Pharmacol, 2018; 29(4):370–4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29035917>

McCorkindale AN, Sizemova A, Sheedy D, Kril JJ, and Sutherland GT. Re-investigating the effects of chronic smoking on the pathology of alcohol-related human brain damage. Alcohol, 2018; 76:11-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30529017>

McClure EA, Tomko RL, Salazar CA, Akbar SA, Squeglia LM, et al. Tobacco and cannabis co-use: Drug substitution, quit interest, and cessation preferences. Experimental and Clinical Psychopharmacology, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30556733>

McClure EA, Baker NL, Sonne SC, Ghitza UE, Tomko RL, et al. Tobacco use during cannabis cessation: Use patterns and impact on abstinence in a national drug abuse treatment clinical trials network study. Drug and Alcohol Dependence, 2018; 192:59-66. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30218864>

Lim AC, Courtney KE, Moallem NR, Allen VC, Jr., Leventhal AM, et al. A brief smoking cessation intervention for heavy drinking smokers: Treatment feasibility and acceptability. Front Psychiatry, 2018; 9:362. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30147661>

Hurt RT, Ebbert JO, Croghan IT, Schroeder DR, Hurt RD, et al. Varenicline for tobacco-dependence treatment in alcohol-dependent smokers: A randomized controlled trial. *Drug and Alcohol Dependence*, 2018; 184:12–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29324248>

Huggett SB, Hatoum AS, Hewitt JK, and Stallings MC. The speed of progression to tobacco and alcohol dependence: A twin study. *Behavior Genetics*, 2018; 48(2):109-24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29427143>

Holahan CJ, Brennan PL, Schutte KK, Holahan CK, Hixon JG, et al. Drinking level versus drinking pattern and cigarette smoking among older adults. *Alcohol Clin Exp Res*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29417610>

Herrmann ES, Cooper ZD, Bedi G, Ramesh D, Reed SC, et al. Varenicline and nabilone in tobacco and cannabis co-users: Effects on tobacco abstinence, withdrawal and a laboratory model of cannabis relapse. *Addict Biol*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30378231>

Healey A, Roberts S, Sevdalis N, Goulding L, Wilson S, et al. A cost-effectiveness analysis of stop smoking interventions in substance use disorder populations undergoing treatment. *Nicotine & Tobacco Research*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29733376>

Hardy L, Parker S, Hartley L, and Hogarth L. A concurrent pictorial drug choice task marks multiple risk factors in treatment-engaged smokers and drinkers. *Behav Pharmacol*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30169375>

Gubner NR, Williams DD, Pagano A, Campbell BK, and Guydish J. Menthol cigarette smoking among individuals in treatment for substance use disorders. *Addictive Behaviors*, 2018; 80:135-41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29407684>

Gass JC, Morris DH, Winters J, VanderVeen JW, and Chermack S. Characteristics and clinical treatment of tobacco smokers enrolled in a substance use disorders clinic. *Journal of Substance Abuse Treatment*, 2018; 84:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29195588>

Ganavadiya R, Chandrashekar BR, Singh P, Gupta R, Rana PT, et al. Knowledge, attitude, and practice among tobacco and alcohol addicts before and after psychological intervention in a de-addiction center at madhya pradesh, India. *Ind Psychiatry J*, 2018; 27(1):27-40. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30416289>

Fallin-Bennett A, Barnett J, Ducas L, Wiggins AT, McCubbin A, et al. Pilot tobacco treatment intervention for women in residential treatment for substance use disorder. *J Obstet Gynecol Neonatal Nurs*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30273555>

Fairman BJ, Furr-Holden CD, and Johnson RM. When marijuana is used before cigarettes or alcohol: Demographic predictors and associations with heavy use, cannabis use disorder, and other drug-related outcomes. *Prev Sci*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29770947>

Dermody SS, Hendershot CS, Andrade AK, Novalen M, and Tyndale RF. Changes in nicotine metabolite ratio among daily smokers receiving treatment for alcohol use disorder. *Nicotine & Tobacco Research*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30561731>

Derefinko KJ, Salgado Garcia FI, and Sumrok DD. Smoking cessation for those pursuing recovery from substance use disorders. *Med Clin North Am*, 2018; 102(4):781-96. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29933829>

Cox S and Notley C. Response to campbell et al., (2018) - health risk perceptions and reasons for use of tobacco products among clients in addictions treatment. *Addictive Behaviors*, 2018; 90:378-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30529993>

Campbell BK, Le T, Gubner NR, and Guydish J. Health risk perceptions and reasons for use of tobacco products among clients in addictions treatment. *Addictive Behaviors*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30206006>

Campbell B, Yip D, Le T, Gubner N, and Guydish J. Relationship between tobacco use and health-related quality of life (hrqol) among clients in substance use disorders treatment. *J Psychoactive Drugs*, 2018:1-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30570409>

Braillon A. Re: 'Killing two birds with one stone? Association between tobacco and alcohol consumption.'. *Public Health*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29656790>

Anton RF, Latham PK, Voronin KE, Randall PK, Book SW, et al. Nicotine-use/smoking is associated with the efficacy of naltrexone in the treatment of alcohol dependence. *Alcohol Clin Exp Res*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29431852>

Weinberger AH, Gbedemah M, and Goodwin RD. Cigarette smoking quit rates among adults with and without alcohol use disorders and heavy alcohol use, 2002-2015: A representative sample of the United States population. *Drug and Alcohol Dependence*, 2017; 180:204–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28918239>

Swarbrick MA, Cook JA, Razzano LA, Jonikas JA, Gao N, et al. Correlates of current smoking among adults served by the public mental health system. *Journal of Dual Diagnosis*, 2017; 13(2):82-90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28426346>

Streck JM, Regan S, Chang Y, Kelley JHK, Singer DE, et al. Examining the effects of illicit drug use on tobacco cessation outcomes in the helping hand 2 randomized controlled trial. *Drug and Alcohol Dependence*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28739101>

Skelton E, Tzelepis F, Shakeshaft A, Guillaumier A, Dunlop A, et al. Addressing tobacco in Australian alcohol and other drug treatment settings: A cross-sectional survey of staff attitudes and perceived barriers. *Substance Abuse Treatment, Prevention, and Policy*, 2017; 12(1):20. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28464898>

Skelton E, Tzelepis F, Shakeshaft A, Guillaumier A, Dunlop A, et al. Smoking cessation care provision in Australian alcohol and other drug treatment services: A cross-sectional survey of staff self-reported practices. *Journal of Substance Abuse Treatment*, 2017; 77:101–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28476261>

Shah PA, Cunningham CO, Brisbane MT, DeLuca JP, and Nahvi S. Use of smoking cessation methods among patients receiving office-based buprenorphine maintenance treatment. *Journal of Addiction Medicine*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28885300>

Schacht JP, Randall PK, Latham PK, Voronin KE, Book SW, et al. Predictors of naltrexone response in a randomized trial: Reward-related brain activation, oprm1 genotype, and smoking status. *Neuropsychopharmacology*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28409564>

Sapkota M, Burnham EL, DeVasure JM, Sweeter JM, Hunter CD, et al. Malondialdehyde-acetaldehyde protein adducts are found exclusively in the lungs of smokers with alcohol use disorders and are associated with systemic anti-maa antibodies. *Alcohol Clin Exp Res*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28941289>

Salisbury-Afshar E. Interventions for tobacco cessation in patients being treated for or recovering from substance use disorders. *American Family Physician*, 2017; 96(9):572-3. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29094884>

Quinn PD, Hur K, Chang Z, Krebs EE, Bair MJ, et al. Incident and long-term opioid therapy among patients with psychiatric conditions and medications: A national study of commercial health care claims. *Pain*, 2017; 158(1):140-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27984526>

Przulj D, Hajek P, Snuggs S, and McRobbie H. Changes in alcohol consumption during a stop-smoking attempt and differences between smokers using nicotine replacement and smokers using varenicline. *Nicotine & Tobacco Research*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28521015>

Murphy CM, MacKillop J, Martin RA, Tidey JW, Colby SM, et al. Effects of varenicline versus transdermal nicotine replacement therapy on cigarette demand on quit day in individuals with substance use disorders. *Psychopharmacology (Berl)*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28500373>

Minian N, Baliunas D, Zawertailo L, Noormohamed A, Giesbrecht N, et al. Combining alcohol interventions with tobacco addictions treatment in primary care-the combat study: A pragmatic cluster randomized trial. *Implement Sci*, 2017; 12(1):65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28521782>

McKelvey K, Thrul J, and Ramo D. Impact of quitting smoking and smoking cessation treatment on substance use outcomes: An updated and narrative review. *Addictive Behaviors*, 2017; 65:161–70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27816663>

MacLean RR, Sofuoglu M, and Rosenheck R. Tobacco and alcohol use disorders: Evaluating multimorbidity. *Addictive Behaviors*, 2017; 78:59-66. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29127785>

Knudsen HK. Author's response: Smoking cessation treatment in substance use disorder settings. *American Journal of Drug and Alcohol Abuse*, 2017;1. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28402753>

Kahler CW, Cioe PA, Tzilos GK, Spillane NS, Leggio L, et al. A double-blind randomized placebo-controlled trial of oral naltrexone for heavy-drinking smokers seeking smoking cessation treatment. *Alcohol Clin Exp Res*, 2017; 41(6):1201-11. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28401564>

Isgro M, Doran N, Heffner JL, Wong E, Dinh E, et al. Type a/type b alcoholism predicts differential response to topiramate in a smoking cessation trial in dually diagnosed men. *J Stud Alcohol Drugs*, 2017; 78(2):232-40. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28317503>

Hufnagel A, Frick U, Ridinger M, and Wodarz N. Recovery from alcohol dependence: Do smoking indicators predict abstinence? *American Journal on Addictions*, 2017; 26(4):366-73. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28376287>

Hall SM, Humfleet G, Gasper J, Delucchi KL, Hersh D, et al. Cigarette smoking cessation intervention for buprenorphine treatment patients. *Nicotine & Tobacco Research*, 2017. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28549161>

Gubner NR, Guydish J, Humfleet GL, Benowitz NL, and Hall SM. Nicotine biomarkers and rate of nicotine metabolism among cigarette smokers taking buprenorphine for opioid dependency. *Drug and Alcohol Dependence*, 2017; 178:267-70. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28683421>

Gentry S, Craig J, Holland R, and Notley C. Smoking cessation for substance misusers: A systematic review of qualitative studies on participant and provider beliefs and perceptions. *Drug and Alcohol Dependence*, 2017; 180:178–92. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28910690>

Evins AE and Kelly JF. A call to action for treatment of comorbid tobacco and alcohol dependence. *JAMA Psychiatry*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29261821>

Das S, Hickman NJ, and Prochaska JJ. Treating smoking in adults with co-occurring acute psychiatric and addictive disorders. *Journal of Addiction Medicine*, 2017. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28441272>

Cooperman NA, Lu SE, Richter KP, Bernstein SL, and Williams JM. Pilot study of a tailored smoking cessation intervention for individuals in treatment for opioid dependence. *Nicotine & Tobacco Research*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29059389>

Berman BA, Jones L, Jones F, Jones A, Pacheco BA, et al. How can we help african american substance users stop smoking? Client and agency perspectives. *J Ethn Subst Abuse*, 2017:1-17. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29257942>

Beckham JC, Adkisson KA, Hertzberg J, Kimbrel NA, Budney AJ, et al. Mobile contingency management as an adjunctive treatment for co-morbid cannabis use disorder and cigarette smoking. *Addictive Behaviors*, 2017; 79:86-92. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29268146>

Beard E, West R, Michie S, and Brown J. Association between smoking and alcohol-related behaviours: A time-series analysis of population trends in england. *Addiction*, 2017. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28556467>

Ainscough TS, Brose LS, Strang J, and McNeill A. Contingency management for tobacco smoking during opioid addiction treatment: A randomised pilot study. *BMJ Open*, 2017; 7(9):e017467. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28864706>

Winhusen T, Theobald J, and Lewis D. Design considerations for a pilot trial using a novel approach for evaluating smoking-cessation medication in methadone-maintained smokers. *Contemporary Clinical Trials*, 2016; 47:334–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26944021>

Winhusen T and Lewis D. Cigarette smoking in pregnant substance users: Association with substance use and desire to quit. *Journal of Addictive Diseases*, 2016:1-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27802114>

Weinberger AH, Funk AP, and Goodwin RD. A review of epidemiologic research on smoking behavior among persons with alcohol and illicit substance use disorders. *Preventive Medicine*, 2016; 92:148–59. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27196143>

Thurgood SL, McNeill A, Clark-Carter D, and Brose LS. A systematic review of smoking cessation interventions for adults in substance abuse treatment or recovery. *Nicotine & Tobacco Research*, 2016; 18(5):993–1001. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26069036>

Secades-Villa R, Pericot-Valverde I, and Weidberg S. Relative reinforcing efficacy of cigarettes as a predictor of smoking abstinence among treatment-seeking smokers. *Psychopharmacology (Berl)*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27325392>

Ruther T, Ruderer A, Wirth C, Schuler V, Lang V, et al. Smoking cessation program for inpatients with substance use disorder: A quasi-randomized controlled trial of feasibility and efficacy. *Eur Addict Res*, 2016; 22(5):268–76. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27287751>

Roche DJ, Ray LA, Yardley MM, and King AC. Current insights into the mechanisms and development of treatments for heavy drinking cigarette smokers. *Curr Addict Rep*, 2016; 3(1):125–37. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27162709>

Richter KP, Hunt JJ, Cupertino AP, Gajewski BJ, Jiang Y, et al. Commitment and capacity for providing evidence-based tobacco treatment in u.S. Drug treatment facilities. *Substance Abuse*, 2016:0. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27897468>

Rabin RA, Ashare RL, Schnoll RA, Cinciripini PM, Hawk LW, Jr., et al. Does cannabis use moderate smoking cessation outcomes in treatment-seeking tobacco smokers? Analysis from a large multi-center trial. *American Journal on Addictions*, 2016; 25(4):291–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27187893>

Pollack HA. Commentary on guydish et al. (2016): The continuing deadly threat of tobacco use in addiction treatment. *Addiction*, 2016; 111(2):231–2. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26767337>

Pagano A, Tajima B, and Guydish J. Barriers and facilitators to tobacco cessation in a nationwide sample of addiction treatment programs. *Journal of Substance Abuse Treatment*, 2016; 67:22–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27296658>

Meyerhoff DJ. Time for cigarette smoke exposure chambers? *Alcohol Clin Exp Res*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26990486>

Martin RA, Cassidy RN, Murphy CM, and Rohsenow DJ. Barriers to quitting smoking among substance dependent patients predict smoking cessation treatment outcome. *Journal of Substance Abuse Treatment*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26979552>

Guydish J, Tajima B, Pramod S, Le T, Gubner NR, et al. Use of multiple tobacco products in a national sample of persons enrolled in addiction treatment. *Drug and Alcohol Dependence*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27449271>

Gubner NR, Kozar-Konieczna A, Szoltysek-Boldys I, Slodczyk-Mankowska E, Goniewicz J, et al. Cessation of alcohol consumption decreases rate of nicotine metabolism in male alcohol-dependent smokers. *Drug and Alcohol Dependence*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27107849>

Fallin A, Miller A, and Ashford K. Smoking among pregnant women in outpatient treatment for opioid dependence: A qualitative inquiry. *Nicotine & Tobacco Research*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26817489>

de Dios MA, Cano MA, Childress S, Vaughan E, Cerna Y, et al. Smoking status and substance use treatment outcomes among spanish speakers enrolled in substance abuse treatment. *J Clin Psychol*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27092710>

Cooperman NA, Lu SE, Richter KP, Bernstein SL, and Williams JM. Influence of psychiatric and personality disorders on smoking cessation among individuals in opiate dependence treatment. *Journal of Dual Diagnosis*, 2016:0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27064523>

Campbell BK, Le T, Andrews KB, Pramod S, and Guydish J. Smoking among patients in substance use disorders treatment: Associations with tobacco advertising, anti-tobacco messages, and perceived health risks. *American Journal of Drug and Alcohol Abuse*, 2016; 42(6):649-56. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27314450>

Bonevski B, Guillaumier A, Shakeshaft A, Farrell M, Tzelepis F, et al. An organisational change intervention for increasing the delivery of smoking cessation support in addiction treatment centres: Study protocol for a randomized controlled trial. *Trials*, 2016; 17(1):290. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27301489>

Bhuiyan J, Jonkman L, Connor S, and Giannetti V. Qualitative evaluation of perceptions of smoking cessation among clients at an alcohol and other drug treatment program. *Research in Social and Administrative Pharmacy*, 2016. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27856209>

Apollonio D, Philipps R, and Bero L. Interventions for tobacco use cessation in people in treatment for or recovery from substance use disorders. *Cochrane Database of Systematic Reviews*, 2016; 11:CD010274. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27878808>

Zawertailo LA, Baliunas D, Ivanova A, and Selby PL. Individualized treatment for tobacco dependence in addictions treatment settings: The role of current depressive symptoms on outcomes at 3 and 6

months. *Nicotine & Tobacco Research*, 2015; 17(8):937–45. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/26180218>

Yardley MM, Mirbaba MM, and Ray LA. Pharmacological options for smoking cessation in heavy-drinking smokers. *CNS Drugs*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26507831>

Wilson AJ, Bonevski B, Dunlop A, Shakeshaft A, Tzelepis F, et al. 'The lesser of two evils': A qualitative study of staff and client experiences and beliefs about addressing tobacco in addiction treatment settings. *Drug and Alcohol Review*, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/26332170>

Weinberger AH, Platt J, Jiang B, and Goodwin RD. Cigarette smoking and risk of alcohol use relapse among adults in recovery from alcohol use disorders. *Alcohol Clin Exp Res*, 2015; 39(10):1989–96.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26365044>

Stuyt EB. Enforced abstinence from tobacco during in-patient dual-diagnosis treatment improves substance abuse treatment outcomes in smokers. *American Journal on Addictions*, 2015; 24(3):252–

7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25907814>

Sigmon SC, Miller ME, Meyer AC, Saulsgiver K, Badger GJ, et al. Financial incentives to promote extended smoking abstinence in opioid-maintained patients: A randomized trial. *Addiction*, 2015.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26638126>

Shu C and Cook BL. Examining the association between substance use disorder treatment and smoking cessation. *Addiction*, 2015; 110(6):1015–24. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25683883>

Shi Y and Cummins SE. Smoking cessation services and smoke-free policies at substance abuse treatment facilities: National survey results. *Psychiatric Services*, 2015:appips201400328. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25686819>

Rohsenow DJ, Tidey JW, Martin RA, Colby SM, Sirota AD, et al. Contingent vouchers and motivational interviewing for cigarette smokers in residential substance abuse treatment. *Journal of Substance Abuse Treatment*, 2015; 55:29–38. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25805668>

Rohsenow DJ, Tidey JW, Kahler CW, Martin RA, Colby SM, et al. Intolerance for withdrawal discomfort and motivation predict voucher-based smoking treatment outcomes for smokers with substance use disorders. *Addictive Behaviors*, 2015; 43:18–24. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25531536>

Peters EN, Harrell PT, Hendricks PS, O'Grady KE, Pickworth WB, et al. Electronic cigarettes in adults in outpatient substance use treatment: Awareness, perceptions, use, and reasons for use. *American Journal on Addictions*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25809200>

Muilenburg JL, Laschober TC, and Eby LT. Relationship between low-income patient census and substance use disorder treatment programs' availability of tobacco cessation services. *J Drug Issues*, 2015; 45(1):69–79. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25530629>

Miller ME and Sigmon SC. Are pharmacotherapies ineffective in opioid-dependent smokers? Reflections on the scientific literature and future directions. *Nicotine & Tobacco Research*, 2015; 17(8):955–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26180219>

Mandal P, Jain R, Jhanjee S, and Sreenivas V. Psychological barriers to tobacco cessation in indian buprenorphine-naloxone maintained patients: A pilot study. *Indian J Psychol Med*, 2015; 37(3):299–304. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26664077>

Magee JC, Lewis DF, and Winhusen T. Evaluating nicotine craving, withdrawal, and substance use as mediators of smoking cessation in cocaine- and methamphetamine-dependent patients. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26048168>

Mackillop J, Murphy CM, Martin RA, Stojek M, Tidey JW, et al. Predictive validity of a cigarette purchase task in a randomized controlled trial of contingent vouchers for smoking in individuals with substance use disorders. *Nicotine & Tobacco Research*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26498173>

Laschober TC, Muilenburg JL, and Eby LT. Factors linked to substance use disorder counselors' (non)implementation likelihood of tobacco cessation 5 a's, counseling, and pharmacotherapy. *Journal of Addictive Behaviors, Therapy and Rehabilitation*, 2015; 4(1). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26005696>

Krauth D and Apollonio DE. Overview of state policies requiring smoking cessation therapy in psychiatric hospitals and drug abuse treatment centers. *Tobacco Induced Diseases*, 2015; 13:33. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26500471>

Krauth D and Apollonio D. Accuracy of popular media reporting on tobacco cessation therapy in substance abuse and mental health populations. *BMJ Open*, 2015; 5(3):e007169. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25814498>

Knudsen HK and Roman PM. Medicaid, private insurance, and the availability of smoking cessation interventions in substance use disorder treatment. *Psychiatric Services*, 2015:appips201400451. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26234332>

Hitschfeld MJ, Schneekloth TD, Ebbert JO, Hall-Flavin DK, Karpyak VM, et al. Female smokers have the highest alcohol craving in a residential alcoholism treatment cohort. *Drug and Alcohol Dependence*, 2015; 150:179–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25746235>

Guydish J, Gruber VA, Le T, Tajima B, Andrews KB, et al. A pilot study of a readiness group to increase initiation of smoking cessation services among women in residential addiction treatment. *Journal of Substance Abuse Treatment*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26825975>

Grigsby TJ, Forster M, and Sussman S. A perspective on cigarette smoking during alcohol and substance use treatment. *Substance Use and Misuse*, 2015:1–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25774483>

Cooperman NA, Richter KP, Bernstein SL, Steinberg ML, and Williams JM. Determining smoking cessation related information, motivation, and behavioral skills among opiate dependent smokers in

methadone treatment. Substance Use and Misuse, 2015. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/25559697>

Boehm G, Schroeder Y, and Schoberberger R. Inpatient smoking cessation therapy: Truth or dare?

Wien Klin Wochenschr, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26142170>

Winhusen TM, Kropp F, Theobald J, and Lewis DF. Achieving smoking abstinence is associated with decreased cocaine use in cocaine-dependent patients receiving smoking-cessation treatment. Drug and Alcohol Dependence, 2014; 134:391–5. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24128381>

Winhusen TM, Brigham GS, Kropp F, Lindblad R, Gardin JG, 2nd, et al. A randomized trial of concurrent smoking-cessation and substance use disorder treatment in stimulant-dependent smokers. Journal of Clinical Psychiatry, 2014; 75(4):336–43. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24345356>

Shealy SE and Winn JL. Integrating smoking cessation into substance use disorder treatment for military veterans: Measurement and treatment engagement efforts. Addictive Behaviors, 2014;

39(2):439–44. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24094921>

Secades-Villa R, Garcia-Rodriguez O, Lopez-Nunez C, Alonso-Perez F, and Fernandez-Hermida JR. Contingency management for smoking cessation among treatment-seeking patients in a community setting. Drug and Alcohol Dependence, 2014; 140:63–8. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24768410>

Rohsenow DJ, Martin RA, Monti PM, Colby SM, Day AM, et al. Motivational interviewing versus brief advice for cigarette smokers in residential alcohol treatment. Journal of Substance Abuse Treatment, 2014; 46(3):346–55. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24210533>

Patrick ME, Dunn KE, Badger GJ, Heil SH, Higgins ST, et al. Spontaneous reductions in smoking during double-blind buprenorphine detoxification. Addictive Behaviors, 2014; 39(9):1353–6. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24845165>

Nahvi S, Blackstock O, Sohler NL, Thompson D, and Cunningham CO. Smoking cessation treatment among office-based buprenorphine treatment patients. Journal of Substance Abuse Treatment, 2014; 47(2):175–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24912863>

Muilenburg JL, Laschober TC, and Eby LT. Organizational factors as predictors of tobacco cessation pharmacotherapy adoption in addiction treatment programs. Journal of Addiction Medicine, 2014; 8(1):59–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24365803>

Leggio L, Zywiak WH, Edwards SM, Tidey JW, Swift RM, et al. A preliminary double-blind, placebo-controlled randomized study of baclofen effects in alcoholic smokers. Psychopharmacology (Berl), 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24973894>

Lee DC, Budney AJ, Brunette MF, Hughes JR, Etter JF, et al. Treatment models for targeting tobacco use during treatment for cannabis use disorder: Case series. Addictive Behaviors, 2014; 39(8):1224–

30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24813547>

Karam-Hage M, Robinson JD, Lodhi A, and Brower KJ. Bupropion-sr for smoking reduction and cessation in alcohol-dependent outpatients: A naturalistic, open-label study. *Curr Clin Pharmacol*, 2014; 9(2):123–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24218993>

Jacques D, Manceaux P, de Timary P, Tecco J, Reynaert C, et al. Proposing concurrent alcohol and tobacco withdrawal. Analysis of the perceptions of belgian nurses in the alcoholism unit. *Psychiatr Danub*, 2014; 26 Suppl 1:41–2. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25413510>

Hunt JJ, Cupertino AP, Gajewski BJ, Jiang Y, Ronzani TM, et al. Staff commitment to providing tobacco dependence in drug treatment: Reliability, validity, and results of a national survey. *Psychol Addict Behav*, 2014; 28(2):389–95. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24128292>

Hendricks PS, Peters EN, Thorne CB, Delucchi KL, and Hall SM. Expectancies for smoking cessation among drug-involved smokers: Implications for clinical practice. *Journal of Substance Abuse Treatment*, 2014; 46(3):320–4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24314605>

Foster DW, Zvolensky MJ, Garey L, Ditre JW, and Schmidt NB. Relationships between drinking motives and smoking expectancies among daily smokers who are also problem drinkers. *Journal of Dual Diagnosis*, 2014; 10(3):118–29. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25392285>

Eby LT, Laschober TC, and Muilenburg JL. Sustained, new, never, and discontinued tobacco cessation services adopters. *Journal of Substance Abuse Treatment*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25178991>

de Dios MA, Anderson BJ, Caviness CM, and Stein MD. Early quit days among methadone-maintained smokers in a smoking cessation trial. *Nicotine & Tobacco Research*, 2014; 16(11):1463–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24951495>

Dahne J, Hoffman EM, and MacPherson L. The association between anxiety sensitivity and motivation to quit smoking among women and men in residential substance use treatment. *Substance Use and Misuse*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25265420>

Cookson C, Strang J, Ratschen E, Sutherland G, Finch E, et al. Smoking and its treatment in addiction services: Clients' and staff behaviour and attitudes. *BMC Health Services Research*, 2014; 14:304. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25017205>

Breland AB, Almond L, Kienzle J, Ondersma SJ, Hart A, Jr., et al. Targeting tobacco in a community-based addiction recovery cohort: Results from a computerized, brief, randomized intervention trial. *Contemporary Clinical Trials*, 2014; 38(1):113–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24721481>

Alessi SM and Petry NM. Smoking reductions and increased self-efficacy in a randomized controlled trial of smoking abstinence-contingent incentives in residential substance abuse treatment patients. *Nicotine & Tobacco Research*, 2014; 16(11):1436–45. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24935755>

Luhar RB, Sawyer KS, Gravitz Z, Ruiz SM, and Oscar-Berman M. Brain volumes and neuropsychological performance are related to current smoking and alcoholism history.

Neuropsychiatr Dis Treat, 2013; 9:1767–84. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/24273408>

Guydish J, Ziedonis D, Tajima B, Seward G, Passalacqua E, et al. Addressing tobacco through organizational change (attoc) in residential addiction treatment settings. *Drug and Alcohol Dependence*, 2012; 121(1-2):30–7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/21906892>

Rothrauff TC and Eby LT. Counselors' knowledge of the adoption of tobacco cessation medications in substance abuse treatment programs. *American Journal on Addictions*, 2011; 20(1):56–62. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21175921>

Rothrauff TC, Eby LT, and Public Health S. Substance abuse counselors' implementation of tobacco cessation guidelines. *J Psychoactive Drugs*, 2011; 43(1):6–13. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/21615002>

Reid M, Jiang H, Fallon B, Sonne S, Rinaldi P, et al. Smoking cessation treatment among patients in community-based substance abuse rehabilitation programs: Exploring predictors of outcome as clues toward treatment improvement. *The American Journal of Drug and Alcohol Abuse*, 2011; 37(5):472–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21854292>

Khara M and Okoli CT. The tobacco-dependence clinic: Intensive tobacco-dependence treatment in an addiction services outpatient setting. *American Journal on Addictions*, 2011; 20(1):45–55.

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/21175920>

Karam-Hage M, Strobbe S, Robinson J, and Brower K. Bupropion-sr for smoking cessation in early recovery from alcohol dependence: A placebo-controlled, double-blind pilot study. *The American Journal of Drug and Alcohol Abuse*, 2011; [Epub ahead of print]. Available from:

<http://informahealthcare.com/doi/full/10.3109/00952990.2011.598591>

Hunt J, Cupertino A, Garrett S, Friedmann P, and Richter K. How is tobacco treatment provided during drug treatment? *Journal of Substance Abuse Treatment*, 2011; 42(1):4–15. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/21831563>

Holbrook A and Kaltenbach K. Effectiveness of a smoking cessation intervention for methadone-maintained women: A comparison of pregnant and parenting women. *International Journal of Pediatrics*, 2011; 2011:567056. Available from:

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3147124/?tool=pubmed>

Guydish J, Passalacqua E, Tajima B, Chan M, Chun J, et al. Smoking prevalence in addiction treatment: A review. *Nicotine & Tobacco Research*, 2011; 13(6):401–11. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/21464202>

Fucito LM, Toll BA, Wu R, Romano DM, Tek E, et al. A preliminary investigation of varenicline for heavy drinking smokers. *Psychopharmacology (Berl)*, 2011; 215(4):655–63. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/21221531>

Boissoneault J, Gilbertson R, Prather R, and Nixon S. Contrasting behavioral effects of acute nicotine and chronic smoking in detoxified alcoholics. *Addictive Behaviors*, 2011; 36(12):1344–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21855223>

Teater B and Hammond G. Exploring smoking prevalence, quit attempts, and readiness to quit cigarette use among women in substance abuse treatment. *Social Work in Health Care*, 2010; 49(2):176–92. Available from: <http://www.informaworld.com/smpp/ftinterface~content=a919418592~fulltext=713240928>

Poling J, Rounsaville B, Gonsai K, Severino K, and Sofuoglu M. The safety and efficacy of varenicline in cocaine using smokers maintained on methadone: A pilot study. *American Journal on Addictions*, 2010; 19(5):401–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20716302>

Nieva G, Ortega L, Mondon S, Ballbe M, and Gual A. Simultaneous versus delayed treatment of tobacco dependence in alcohol-dependent outpatients. *European Addiction Research*, 2010; 17(1):1–9. Available from: <http://content.karger.com/ProdukteDB/produkte.asp?Aktion=ShowPDF&ArtikelNr=000321256&Ausgabe=254620&ProduktNr=224233&filename=000321256.pdf>

Hogue A, Dauber S, and Morgenstern J. Validation of a contemplation ladder in an adult substance use disorder sample. *Psychology of Addictive Behaviors*, 2010; 24(1):137–44. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2845310/?tool=pubmed>

Dunn KE, Sigmon SC, Reimann EF, Badger GJ, Heil SH, et al. A contingency-management intervention to promote initial smoking cessation among opioid-maintained patients. *Experimental and Clinical Psychopharmacology*, 2010; 18(1):37–50. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20158293>

Allsop S, Carter O, and Lenton S. Enhancing clinical research with alcohol, tobacco and cannabis problems and dependence. *Drug and Alcohol Review*, 2010; 29(5):483–90. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20887571>

Toussaint DW, VanDeMark NR, Silverstein M, and Stone E. Exploring factors related to readiness to change tobacco use for clients in substance abuse treatment. *Journal of Drug Issues*, 2009; 39(2). Available from: <https://journals.sagepub.com/doi/abs/10.1177/002204260903900203>

Saatcioglu O and Erim R. Aggression among male alcohol-dependent inpatients who smoke cigarettes. *The Journal of Psychology*, 2009; 143(6):615–24. Available from: <http://heldref-publications.metapress.com/app/home/contribution.asp?referrer=parent&backto=issue,3,5;journal,3,80;linkingpublicationresults,1:119938,1>

Roth T. Does effective management of sleep disorders reduce substance dependence? *Drugs*, 2009; 69(suppl. 2):S65–S75. Available from: <http://adisonline.com/drugs/pages/articleviewer.aspx?year=2009&issue=69002&article=00006&type=abstract>

McKee SA, Harrison ELR, O'Malley SS, Krishnan-Sarin S, Shi J, et al. Varenicline reduces alcohol self-administration in heavy-drinking smokers. *Journal of Biological Psychiatry*, 2009; In Press.

Jones H, Heil S, O'Grady K, Martin P, Kaltenbach K, et al. Smoking in pregnant women screened for an opioid agonist medication study compared to related pregnant and non-pregnant patient samples. *The American Journal of Drug and Alcohol Abuse*, 2009; 35(5):375–80. Available from: <http://informahealthcare.com/doi/full/10.1080/00952990903125235>

Hays JT, Hurt RD, Decker PA, Croghan IT, Offord KP, et al. A randomized, controlled trial of bupropion sustained-release for preventing tobacco relapse in recovering alcoholics. *Nicotine & Tobacco Research*, 2009; 11(7):859–67. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19483180>

Hall SM and Prochaska JJ. Treatment of smokers with co-occurring disorders: Emphasis on integration in mental health and addiction treatment settings. *Annu Rev Clin Psychol*, 2009; 5:409–31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19327035>

Ericson M, Löf E, Stomberg R, and Söderpalm B. The smoking cessation medication varenicline attenuates alcohol and nicotine interactions in the rat mesolimbic dopamine system. *Journal of Pharmacology And Experimental Therapeutics*, 2009; 329(1):225–30. Available from: <http://jpet.aspetjournals.org/cgi/content/full/329/1/225>

Elkader A, Brands B, Selby P, and Sproule B. Methadone-nicotine interactions in methadone maintenance treatment patients. *Journal of Clinical Psychopharmacology*, 2009; 29(3):231–8. Available from: <http://journals.lww.com/psychopharmacology/pages/articleviewer.aspx?year=2009&issue=06000&article=00007&type=abstract>

Diehl A, Nakovics H, Mutschler J, Hermann D, and Kiefer F. Rivastigmine reduces tobacco craving in alcohol-dependent smokers. *Pharmacopsychiatry*, 2009; 42(3):89–94. Available from: <http://www.thieme-connect.com/ejournals/html/pharmaco/doi/10.1055/s-0028-1103295>

Chisolm M, Tuten M, Brigham E, Strain E, and Jones H. Relationship between cigarette use and mood/anxiety disorders among pregnant methadone-maintained patients. *American Journal on Addictions*, 2009; 18(5):422–9. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=19874163

Saatcioglu O, Celikel F, and Cakmak D. Depression and anxiety in alcohol dependent inpatients who smoke. *The Israel Journal of Psychiatry and Related Sciences*, 2008; 45(1):33–8. Available from: <http://www.highbeam.com/doc/1P3-1502905851.html>

Ream G, Benoit E, Johnson B, and Dunlap E. Smoking tobacco along with marijuana increases symptoms of cannabis dependence. *Drug and Alcohol Dependence*, 2008; 95(3):199–208. Available from: [http://linkinghub.elsevier.com/retrieve/pii/S0376-8716\(08\)00048-3](http://linkinghub.elsevier.com/retrieve/pii/S0376-8716(08)00048-3)

Myers M and Prochaska J. Does smoking intervention influence adolescent substance use disorder treatment outcomes? *Substance Abuse*, 2008; 29(2):81–8. Available from: <https://www.haworthpress.com/store/ArticleAbstract.asp?sid=L5MP1KGNP29S9NURWS4WJEAMN5380CGC&ID=109812>

Mooney M, Poling J, Gonzalez G, Gonsai K, Kosten T, et al. Preliminary study of buprenorphine and bupropion for opioid-dependent smokers. *American Journal on Addictions* 2008; 17(4):287–92. Available from:

<http://www.informaworld.com/smpp/content~db=all?content=10.1080/10550490802138814>

McCambridge J and Jenkins RJ. Do brief interventions which target alcohol consumption also reduce cigarette smoking? Systematic review and meta-analysis. *Drug and Alcohol Dependence*, 2008; 96(3):263–70. Available from: [http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6T63-4SK4X94-](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6T63-4SK4X94-1&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_version=1&_urlVersion=0&_userid=10&md5=5b73c8994f7c8c299263581caf471a2c)

[1&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_version=1&_urlVersion=0&_userid=10&md5=5b73c8994f7c8c299263581caf471a2c](http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6T63-4SK4X94-1&_user=10&_rdoc=1&_fmt=&_orig=search&_sort=d&_view=c&_version=1&_urlVersion=0&_userid=10&md5=5b73c8994f7c8c299263581caf471a2c)

Jessup M and Song Y. Tobacco-related practices and policies in residential perinatal drug treatment programs. *Journal of Psychoactive Drugs*, 2008; (Suppl 5):357–64. Available from:

http://www.unboundmedicine.com/medline/ebm/record/19248393/full_citation/Tobacco_related_practices_and_policies_in_residential_perinatal_drug_treatment_programs

Finkelstein J, Lapshin O, and Cha E. Feasibility of promoting smoking cessation among methadone users using multimedia computer-assisted education. *Journal of Medical Internet Research*, 2008; 10(5):e33. Available from: <http://www.jmir.org/2008/5/e33/>

Durazzo T, Rothlind J, Gazdzinski S, and Meyerhoff D. The relationships of sociodemographic factors, medical, psychiatric, and substance-misuse co-morbidities to neurocognition in short-term abstinent alcohol-dependent individuals. *Alcohol*, 2008; 42(6):439–49. Available from:

http://www.sciencedirect.com/science?_ob=MImg&_imagekey=B6T40-4T9VGCM-1-1&_cdi=4960&_user=10&_orig=search&_coverDate=09%2F30%2F2008&_sk=999579993&_view=c&_wchp=dGLzVzz-zSkzS&md5=61a885664c0f6dba8d40045c70045c1f&ie=/sdarticle.pdf

Dunn K, Sigmon S, Thomas C, Heil S, and Higgins S. Voucher-based contingent reinforcement of smoking abstinence among methadone-maintained patients: A pilot study. *Journal of Applied Behavior Analysis*, 2008; 41(4):527–38. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/19192857>

Baca C and Yahne C. Smoking cessation during substance abuse treatment: What you need to know. *Journal of Substance Abuse Treatment*, 2008; 36(2):205–19. Available from:

[http://www.journalofsubstanceabusetreatment.com/article/S0740-5472\(08\)00097-4/abstract](http://www.journalofsubstanceabusetreatment.com/article/S0740-5472(08)00097-4/abstract)

Alessi SM, Petry NM, and Urso J. Contingency management promotes smoking reductions in residential substance abuse patients. *Journal of Applied Behavior Analysis*, 2008; 41(4):617–22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/19192865>

Steensland P, Simms JA, Holgate J, Richards JK, and E. Bartlett SE. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, selectively decreases ethanol consumption and seeking. *Proceedings of the National Academy of Sciences of the United States of America*, 2007; 104(30):12518–23. Available from:

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pubmed&pubmedid=17626178>

McKee SA, Falba T, O'Malley SS, Sindelar J, and O'Connor PG. Smoking status as a clinical indicator for alcohol misuse in US adults Archives of Internal Medicine 2007; 167:716–21.

Gershon Grand R, Hwang S, Han J, George T, and Brody A. Short-term naturalistic treatment outcomes in cigarette smokers with substance abuse and/or mental illness. The Journal of Clinical Psychiatry 2007; 68(6):892–8.

Durazzo T, Gazdzinski S, and Meyerhoff D. The neurobiological and neurocognitive consequences of chronic cigarette smoking in alcohol use disorders. Alcohol and Alcoholism, 2007; 42(3):174– 5. Available from: <http://alcalc.oxfordjournals.org/cgi/content/full/42/3/174>

Walsh RA, Bowman JA, Tzelepis F, and Lecathelinais C. Regulation of environmental tobacco smoke by Australian drug treatment agencies. Australian and New Zealand Journal of Public Health, 2005; 29(3):276–8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15991778>

Walsh RA, Bowman JA, Tzelepis F, and Lecathelinais C. Smoking cessation interventions in Australian drug treatment agencies: A national survey of attitudes and practices. Drug and Alcohol Review, 2005; 24(3):235–44. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16096127>

Penetar DM, Kouri EM, Gross MM, McCarthy EM, Rhee CK, et al. Transdermal nicotine alters some of marihuana's effects in male and female volunteers. Drug and Alcohol Dependence, 2005; 79(2):211– 23. Available from: <http://www.mrw.interscience.wiley.com/cochrane/clcentral/articles/851/CN-00560851/frame.html>

McCarthy WJ, Collins C, and Hser YI. Does cigarette smoking affect drug abuse treatment? Journal of Drug Issues, 2002; 32(2):61–79. Available from: <http://www.highbeam.com/doc/1P3-117543970.html>

McLellan T, Lewis D, O'Brien C, and Kleber H. Drug dependence, a chronic medical illness: Implications for treatment, insurance and outcomes evaluations. Journal of the American Medical Association, 2000; 284:1689–95. Available from: <http://jama.ama-assn.org/cgi/content/full/284/13/1689>

Richmond R, G-Novak K, Kehoe L, Calfas G, Mendelson C, et al. Effect of training on general practitioners' use of a brief intervention for excessive drinkers. Australian and New Zealand Journal of Public Health, 1998; 22(2):206–9. Available from: <http://www3.interscience.wiley.com/journal/120143837/abstract>

Roll J, Higgins S, and Tidey J. Cocaine use can increase cigarette smoking: Evidence from laboratory and naturalistic settings. Experimental and Clinical Psychopharmacology, 1997; 5(3):263–8. Available from: <http://psycnet.apa.org/index.cfm?fa=main.landing>

Bien TH and Burge R. Smoking and drinking: A review of the literature. Int J Addict, 1990; 25(12):1429–54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/2094682>

9A.3.5.6 Post-traumatic stress disorder (PTSD)

Young, JR, Polick, CS, Michael, AM, Dannhauer, M, Galla, JT, Evans, MK et al. (2024). Multimodal smoking cessation treatment combining repetitive transcranial magnetic stimulation, cognitive behavioral therapy, and nicotine replacement in veterans with posttraumatic stress disorder: A feasibility randomized controlled trial protocol. *PLoS One*, 19(9), e0291562. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39240791>

Shevorykin, A, Hyland, BM, Robles, D, Ji, M, Vantucci, D, Bensch, L et al. (2024). Tobacco use, trauma exposure and PTSD: a systematic review. *Health Psychol Rev*, 1-32. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38711288>

Zvolensky, MJ, Shepherd, JM, Clausen, BK, Robison, J, Cano, MA, de Dios, M, & Correa-Fernandez, V. (2024). Posttraumatic stress and probable post traumatic stress disorder as it relates to smoking behavior and beliefs among trauma exposed hispanic persons who smoke. *J Behav Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38409553>

Barrett, BW, Meanley, S, Brennan-Ing, M, Haberen, SA, Ware, D, Detels, R et al. (2023). The Relationship Between Posttraumatic Stress Disorder and Alcohol Misuse and Smoking Among Aging Men Who Have Sex With Men: No Evidence of Exercise or Volunteering Impact. *J Aging Health*, 8982643231215475. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37976419>

Jin, Y, Xu, S, Luo, X, Wang, Y, Li, J, Liang, B et al. (2023). A network approach to the symptom-level associations between smoking and posttraumatic stress disorder (PTSD) among young adults exposed to childhood sexual abuse. *J Glob Health*, 13, 04037. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37350563>

Tran, DD, Davis, JP, Tucker, JS, Bricker, J, Lee, DS, Fitzke, RE, & Pedersen, ER. (2023). Cigarette smoking and depression among US veterans: Longitudinal associations with posttraumatic stress disorder. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37094359>

Lilly, G, & Calvert, GM. (2023). The World Trade Center Health Program: Smoking cessation. *Arch Environ Occup Health*, 1-4. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36919568>

Nizio, P, Smit, T, Matoska, CT, Chavez, J, Tullos, EA, Garey, L et al. (2022). Trauma exposure and smoking outcomes: The indirect effects of anxious and depressive symptoms. *Addict Behav*, 134, 107409. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35717891>

Shevorykin A, Ruglass LM, Mancini AD, Carl E, Legg A, et al. Exploring the role of traumatic event exposure in tobacco dependence treatment outcomes among african americans. *J Psychoactive Drugs*, 2021:1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34694215>

Ibrahim C, Le Foll B, and Hassan AN. The effect of nicotine dependence on the risk of developing post-traumatic stress disorder: Results from the national epidemiologic survey on alcohol and related conditions (nesarc). *Nicotine & Tobacco Research*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34734244>

Hammett PJ, Japuntich SJ, Sherman SE, Rogers ES, Danan ER, et al. Proactive tobacco treatment for veterans with posttraumatic stress disorder. *Psychol Trauma*, 2021; 13(1):114–22. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32614201>

Gowarty MA, Aschbrenner KA, and Brunette MF. Acceptability and usability of mobile apps for smoking cessation among young adults with psychotic disorders and other serious mental illness. *Front Psychiatry*, 2021; 12:656538. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34025477>

Budenz A, Klein A, and Prutzman Y. The relationship between trauma exposure and adult tobacco use: Analysis of the national epidemiologic survey on alcohol and related conditions (iii). *Nicotine & Tobacco Research*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33848342>

Rosenblum MS, Engle JL, Piper ME, Kaye JT, and Cook JW. Motives for smoking in those with ptsd, depression, and no psychiatric disorder. *Journal of Dual Diagnosis*, 2020:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32393117>

Peltzer K and Pengpid S. Tobacco use and associated mental symptoms and health risk behaviours amongst individuals 15 years or older in south africa. *S Afr J Psychiatr*, 2020; 26:1499. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33240550>

Osaki Y, Maesato H, Minobe R, Kinjo A, Kuwabara Y, et al. Changes in smoking behavior among victims after the great east Japan earthquake and tsunami. *Environ Health Prev Med*, 2020; 25(1):19. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32527213>

Mu W, Narine K, Farris S, Lieblich S, Zang Y, et al. Trauma-related cognitions predict treatment response in smokers with ptsd: Evidence from cross-lagged panel analyses. *Addictive Behaviors*, 2020; 108:106376. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32413581>

Mathew AR, Yang E, Avery EF, Crane MM, Lange-Maia BS, et al. Trauma exposure, ptsd symptoms, and tobacco use: Does church attendance buffer negative effects? *J Community Psychol*, 2020; 48(7):2364-74. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32789875>

Koyama S, Tabuchi T, Aida J, Osaka K, and Miyashiro I. Determinants of increased tobacco consumption following a major disaster. *Disaster Med Public Health Prep*, 2020:1-5. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32183911>

Kaye JT, Baker TB, Beckham JC, and Cook JW. Tobacco withdrawal symptoms before and after nicotine deprivation in veteran smokers with posttraumatic stress disorder and with major depressive disorder. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33245346>

Gross GM, Bastian LA, Smith NB, Harpaz-Rotem I, and Hoff R. Sex differences in associations between depression and posttraumatic stress disorder symptoms and tobacco use among veterans of recent conflicts. *J Womens Health (Larchmt)*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31934813>

Estey D, Platt J, Goodwin RD, and Weinberger AH. Relationship of trauma exposure and ptsd to cigarette smoking prevalence, frequency, and quantity: Data from a nationally representative sample of u.s. Adults. *Psychol Trauma*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33252969>

Walsh K, Gilmore AK, Schumacher JA, Coffey SF, Frazier PA, et al. Post-sexual assault cigarette smoking: Findings from a randomized clinical trial of a video-based intervention. *Addictive Behaviors*, 2019; 100:106121. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31622944>

Kheirallah KA, Cobb CO, Alsulaiman JW, Alzoubi A, Hoetger C, et al. Trauma exposure, mental health and tobacco use among vulnerable syrian refugee youth in Jordan. *J Public Health (Oxf)*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31742341>

Fonda JR, Gregor KL, Fortier CB, Scioli-Salter ER, McGlinchey RE, et al. Tobacco dependence is associated with increased risk for multi-morbid clustering of posttraumatic stress disorder, depressive disorder, and pain among post-9/11 deployed veterans. *Psychopharmacology (Berl)*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30617565>

Erwin MC, Dennis PA, Coughlin LN, Calhoun PS, and Beckham JC. Examining the relationship between negative affect and posttraumatic stress disorder symptoms among smokers using ecological momentary assessment. *J Affect Disord*, 2019; 253:285-91. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31077971>

Dedert EA, Resick PA, Dennis PA, Wilson SM, Moore SD, et al. Pilot trial of a combined cognitive processing therapy and smoking cessation treatment. *Journal of Addiction Medicine*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30664539>

Asnaani A, Kaczkurkin AN, Fitzgerald HE, Jerud A, and Foa EB. The association between cognitive coping strategies and treatment outcomes in smokers with ptsd. *Psychol Trauma*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31120265>

Alexander AC, Ward KD, Forde DR, Stockton M, and Read MC. Do current smokers use more cigarettes and become more dependent on nicotine because of psychological distress after a natural disaster? *Addictive Behaviors*, 2019; 93:129-34. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30710806>

van den Berk-Clark C, Secrest S, Walls J, Hallberg E, Lustman PJ, et al. Association between posttraumatic stress disorder and lack of exercise, poor diet, obesity, and co-occurring smoking: A systematic review and meta-analysis. *Health Psychology*, 2018; 37(5):407–16. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29698016>

Seelig AD, Bensley KM, Williams EC, Armenta RF, Rivera AC, et al. Longitudinal examination of the influence of individual posttraumatic stress disorder symptoms and clusters of symptoms on the initiation of cigarette smoking. *Journal of Addiction Medicine*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29864086>

Pericot-Valverde I, Elliott RJ, Miller ME, Tidey JW, and Gaalema DE. Posttraumatic stress disorder and tobacco use: A systematic review and meta-analysis. *Addictive Behaviors*, 2018; 84:238–47. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29753221>

Komuro H, Shigemura J, Uchino S, Takahashi S, Nagamine M, et al. Longitudinal factors associated with increased alcohol and tobacco use in fukushima nuclear power plant workers 32 months after

the nuclear disaster: The Fukushima news project study. *J Occup Environ Med*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30335679>

Kearns NT, Carl E, Stein AT, Vujanovic AA, Zvolensky MJ, et al. Posttraumatic stress disorder and cigarette smoking: A systematic review. *Depress Anxiety*, 2018; 35(11):1056–72. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30192425>

Japuntich SJ, Lee LO, Pineles SL, Gregor K, Joos CM, et al. Contingency management and cognitive behavioral therapy for trauma-exposed smokers with and without posttraumatic stress disorder. *Addictive Behaviors*, 2018; 90:136-42. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30391774>

Dedert EA, Dennis PA, Calhoun PS, Dennis MF, and Beckham JC. A randomized clinical trial of nicotine preloading for smoking cessation in people with posttraumatic stress disorder. *Journal of Dual Diagnosis*, 2018:0. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29693495>

Brown ES and Green A. "A focus on topics related to smoking and PTSD". *Journal of Dual Diagnosis*, 2018:1-2. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29505335>

Alexander AC, Ward KD, Forde DR, and Stockton M. Are posttraumatic stress and depressive symptoms pathways to smoking relapse after a natural disaster? *Drug and Alcohol Dependence*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30455073>

9A.3.5.7 Severe mental illness, including bipolar disorder and schizophrenia

Asmita, Patil, PS, & Sahu, N. (2024). Evaluating the Impact of Motivational Enhancement Therapy on Tobacco Cessation in Schizophrenia: A Comprehensive Review. *Cureus*, 16(9), e70046. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39469376>

Johnstone, S, Hubbard, AN, Schenkel, A, Ashare, RL, & Hawk, LW, Jr. (2024). Effects of acute smoking abstinence among people with schizophrenia: A systematic review and meta-analysis of laboratory studies. *Schizophr Res*, 274, 220-236. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39353226>

Xie, H, Ma, R, Yu, M, Wang, T, Chen, J, Liang, J et al. (2024). History of tobacco smoking and alcohol use can predict the effectiveness of electroconvulsive therapy in individuals with schizophrenia: A multicenter clinical trial. *J Psychiatr Res*, 180, 1-7. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39366272>

Foo, CYS, Potter, K, Nielsen, L, Rohila, A, Maravic, MC, Schnitzer, K et al. (2024). Implementation of Community Health Worker Support for Tobacco Cessation: A Mixed-Methods Study. *Psychiatr Serv*, appips20240044. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39118574>

Leutwyler, H, Hubbard, E, Humfleet, G, Souza, R, Balestra, D, & Wallhagen, M. (2024). The Whole Package: A Multi-Component Smoking Cessation Intervention for Adults With Serious Mental Illness: A Qualitative Study. *Tob Use Insights*, 17, 1179173X241253229. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38779493>

Meza, BPL, Pollack, CE, Tilchin, C, Jennings, JM, Latkin, CA, Cather, C et al. (2024). Social networks of people with serious mental illness who smoke: potential role in a smoking cessation intervention. *J Ment Health*, 1-10. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38588708>

Fibbins, H, Ward, PB, Morell, R, Lederman, O, Teasdale, S, Davies, K et al. (2024). Evaluation of a smoking cessation program for adults with severe mental illness in a public mental health service. *J Psychiatr Ment Health Nurs*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38551076>

Kaasgaard, DM, Sorensen, MK, Christiansen, RB, Breum, UN, Asiamah, N, Friis, LBT, & Hjorth, P. (2024). Video consultation and treatment in the community smoking cessation therapy success rates in patients with mental illness: a randomized controlled trial. *Nord J Psychiatry*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38385357>

Cruvinel, E, Mussulman, L, Scheuermann, T, Shergina, E, He, J, Sherman, S et al. (2024). Hospital-Initiated Smoking Cessation Among Patients Admitted with Behavioral Health Conditions. *J Gen Intern Med*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38326585>

Adlard, RJ, Roos, T, & Temmingh, H. (2023). Alcohol, Smoking and Substance Involvement Screening Test validity in bipolar and psychotic disorders. *S Afr J Psychiatr*, 29, 2109. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38223305>

Brady, DJ, Phalen, PL, Roche, DJO, Cowan, T, & Bennett, ME. (2024). A reduction in cigarette smoking improves health-related quality of life and does not worsen psychiatric symptoms in individuals with serious mental illness. *Addict Behav*, 151, 107949. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38176326>

Rajan, S, Mitchell, A, Zavala, GA, Podmore, D, Khali, H, Chowdhury, AH et al. (2023). Tobacco use in people with severe mental illness: Findings from a multi-country survey of mental health institutions in South Asia. *Tob Induc Dis*, 21, 166. . Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38098747>

Lund, MK, Hojlund, M, & Lund, LC. (2023). Smoking and severe mental illness: Drawing inference on the directionality of the association requires careful consideration of the temporality of the data. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37933520>

Anthenelli, RM, McKenna, BS, Giannini, J, Attaluri, SV, Rubin, M, O'Crowley, E et al. (2023). Combining varenicline preloading with Acceptance and Commitment Therapy (ACT) in persons with serious mental illness who smoke: The randomized ACTSLOW pilot feasibility trial. *Drug Alcohol Depend*, 253, 111012. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37931328>

Liang, LS, Xiao, K, Yu, CH, Li, MY, & Xu, PS. (2023). [Progress in the pharmacological treatment of tobacco dependence in special populations]. *Zhonghua Jie He He Hu Xi Za Zhi*, 46(7), 725-729. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37402666>

Garcia-Jimenez, J, Gomez-Sierra, FJ, Martinez-Hortelano, A, Moreno-Merino, P, Girela-Serrano, B, Molero, P, & Gutierrez-Rojas, L. (2023). Cigarette smoking and risk of suicide in bipolar disorder: a systematic review. *Front Psychiatry*, 14, 1179733. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37275988>

- Mazumdar, P, Zavala, G, Aslam, F, Muliya, KP, Chaturvedi, SK, Kandasamy, A et al. (2023). IMPACT smoking cessation support for people with severe mental illness in South Asia (IMPACT 4S): A protocol for a randomised controlled feasibility trial of a combined behavioural and pharmacological support intervention. *PLoS One*, 18(6), e0287185. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37315070>
- Murphy, CM, Onu, MC, Goodwin, C, Williams, DM, & Tidey, JW. (2023). Acute effects of exercise among individuals with schizophrenia who smoke cigarettes. *Addict Behav*, 144, 107749. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37172547>
- Sanchez-Gutierrez, T, Rodriguez-Toscano, E, Roldan, L, Ferraro, L, Parellada, M, Calvo, A et al. (2023). Tobacco use in first-episode psychosis, a multinational EU-GEI study. *Psychol Med*, 1-12. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37185055>
- Dickerson, D, Cain, M, & Garcia, A. (2023). Primary Psychiatric Diagnoses, Commercialized Tobacco Use, and Homelessness: Comparisons Between Urban American Indian/ Alaska Native and Non-American Indian/Alaska Native Adult Clinical Samples. *Am Indian Alsk Native Ment Health Res*, 30(1), 14-34. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37027498>
- Han Almis, B, Tekin, A, Hakyemez Geylani, G, Baltaci, E, & Egeli, A. (2023). Is there a Relationship Between Caffeine Intake and Smoking and Positive and Negative Symptom Severity in Schizophrenia? *Psychiatr Danub*, 35(1), 56-61. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37060593>
- Kerkova, B, Knizkova, K, Sironova, A, Hruby, A, Vecerova, M, Sustova, P et al. (2023). Smoking and attention in schizophrenia spectrum disorders: What are we neglecting? *Front Psychol*, 14, 1114473. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37063581>
- Zoghbi, Y, Barazi, R, Abdelkader, MH, Eltorki, YH, & Ghuloum, S. (2023). First Episode of Psychosis in the Context of Varenicline Treatment for Smoking Cessation. *Cureus*, 15(3), e36677. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37102012>
- Al-Soufi, L, & Costas, J. (2023). Genetic susceptibility for schizophrenia after adjustment by genetic susceptibility for smoking: implications in identification of risk genes and genetic correlation with related traits. *Psychol Med*, 1-11. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36876478>
- Botsari, I, Marouli, G, Arvanitaki, A, & Peritogiannis, V. (2023). Smoking Habits and Attitudes toward Smoking in Patients with Severe Mental Illness in Residential Facilities in Insular Greece. *Healthcare (Basel)*, 11(5). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36900647>
- Caponnetto, P, Maglia, M, Gulizia, A, Prezzavento, GC, Polosa, R, Quattropani, MC, & Signorelli, M S. (2023). Psychological Experience of Smoking Addiction in Family and Friends of Schizophrenic Adults Who Smoke Daily: A Qualitative Study. *Healthcare (Basel)*, 11(5). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36900649>
- Chen, Y. (2023). Altered functional dynamics gradient in schizophrenia with cigarette smoking. *Cereb Cortex*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36864548>

- Han, B, Aung, TW, Volkow, ND, Silveira, ML, Kimmel, HL, Blanco, C, & Compton, WM. (2023). Tobacco Use, Nicotine Dependence, and Cessation Methods in US Adults With Psychosis. *JAMA Netw Open*, 6(3), e234995. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36976558>
- Lin, BD, Vermeulen, JM, Bolhuis, K, Chang, X, Schirmbeck, F, van Eijk, KR. Et al. (2023). Associations between genetic liabilities to smoking behavior and schizophrenia symptoms in patients with a psychotic disorder, their siblings and healthy controls. *Psychiatry Res*, 323, 115143. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36948018>
- Rajalu, BM, Jayarajan, D, Muliya, KP, Sharma, P, Gandhi, S, & Chand, PK. (2023). Development of a Personalized Tobacco Cessation Intervention Package (PTCIP) for Persons with Schizophrenia in India. *Indian J Psychol Med*, 45(2), 132-138. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36925498>
- Kucukaksu, MH, van Meijel, B, Jansen, L, Hoekstra, T, & Adriaanse, MC. (2023). A smoking cessation intervention for people with severe mental illness treated in ambulatory mental health care (KISMET): study protocol of a randomised controlled trial. *BMC Psychiatry*, 23(1), 108. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36797709>
- Correction: Time trends in access to smoking cessation support for people with depression or severe mental illness: a cohort study in English primary care. (2023). *BMJ Open*, 13(2), e048341corr048341. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36792340>
- Davies, K, Courtney, RJ, Summersby-Mitchell, M, Morell, R, Briggs, N, & Lappin, JM. (2023). A systematic review of factors associated with sustained smoking abstinence in people experiencing severe mental illness following participation in a smoking intervention. *Aust N Z J Psychiatry*, 48674221147206. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36744432>
- Donde, C, Pouchon, A, & Polosan, M. (2023). Recovery effect: Cigarette smoking acts on the neural dynamics of early auditory processing in schizophrenia. *Schizophr Res*, 254, 76-77. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36805235>
- Glastad, SH, Aminoff, SR, Hagen, R, Hoegh, MC, Buchmann, CB, Barrett, EA et al. (2023). Nicotine use and non-pathological alcohol use and their relationship to affective symptoms and sleep disturbances in bipolar disorder. *J Affect Disord*, 327, 236-243. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36746243>
- Mattock, R, Owen, L, & Taylor, M. (2023). The cost-effectiveness of tailored smoking cessation interventions for people with severe mental illness: a model-based economic evaluation. *EClinicalMedicine*, 57, 101828. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36798753>
- Nobile, B, Godin, O, Gard, S, Samalin, L, Brousse, G, Loftus, J et al. (2023). Physical and mental health status of former smokers and non-smokers patients with bipolar disorder. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36751870>
- Veldhuizen, S, Behal, A, Zawertailo, L, Melamed, O, Agarwal, M, & Selby, P. (2023). Outcomes Among People With Schizophrenia Participating in General-Population Smoking Cessation Treatment: An Observational Study. *Can J Psychiatry*, 7067437231155693. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36760089>

Wang, Q, Du, W, Wang, H, Geng, P, Sun, Y, Zhang, J et al. (2023). Nicotine's effect on cognition, a friend or foe? *Prog Neuropsychopharmacol Biol Psychiatry*, 124, 110723. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36736944>

Yeap, ZJS, Marsault, J, George, TP, Mizrahi, R, & Rabin, RA. (2023). Does tobacco dependence worsen cannabis withdrawal in people with and without schizophrenia-spectrum disorders? *Am J Addict*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36815595>

Grunze, A, Mosolov, S, Grunze, H, & Born, C. (2022). The detrimental effects of smoking on the course and outcome in adults with bipolar disorder-A narrative review. *Front Psychiatry*, 13, 1114432. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36699491>

Rajalu, BM, Jayarajan, D, Muliya, KP, Sharma, P, Gandhi, S, & Chand, PK. (2023). Effectiveness of personalized tobacco cessation intervention package among patients with schizophrenia and related psychotic disorders - A two-group experimental study. *Asian J Psychiatr*, 81, 103447. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36652840>

Zolezzi, M, Al-Rawi, S & Eltorki, Y. (2022). An Exploration of Smoking Patterns Among People with Serious Mental Illness Attending an Outpatient Clinic in Qatar. *Neuropsychiatr Dis Treat*, 18, 2847-2854. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36518535>

Caponnetto, P, Maglia, M, Mangione, M, Vergopia, C, Prezzavento, GC, Polosa, R et al. (2022). Smoking Addiction in Patients with Schizophrenia Spectrum Disorders and Its Perception and Intervention in Healthcare Personnel Assigned to Psycho-Rehabilitation Programs: A Qualitative Research. *Healthcare (Basel)*, 10(11). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36421599>

Conley, AC, Albert, KM, Armstrong, K, Johnson, JV, Kem, WR, Newhouse, PA, & Lewis, AS. (2022). A pilot study of transdermal nicotine effects on facial emotion responding in non-smoking healthy controls and individuals with schizophrenia: Feasibility and effect size estimates. *Schizophr Res*, 250, 89-91. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36372000>

Nagawa, CS, Lane, IA, McKay, CE, Kamberi, A, Shenette, LL, Kelly, MM et al. (2022). Use of a Rapid Qualitative Method to Inform the Development of a Text Messaging Intervention for People With Serious Mental Illness Who Smoke: Formative Research Study. *JMIR Form Res*, 6(11), e40907. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36342765>

Petty, D, Rae, J, & Aubry, T. (2022). An Examination of the Effectiveness of Smoking Cessation Treatment Interventions for Individuals with Severe Mental Illness: A Pilot Randomized Controlled Trial. *Community Ment Health J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36327043>

Steffens, EB, Park, ER, Muzikansky, A, Streck, JM, Ostroff, JS, Rasmussen, AW et al. (2022). Patients With Serious Mental Illness Can Engage in Tobacco Treatment Trials and Tobacco Cessation Interventions During Cancer Treatment. *JCO Oncol Pract*, OP2200505. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36351206>

Bolstad, I, Alakokkare, AE, Bramness, JG, Rognli, EB, Levola, J, Mustonen, A et al. (2022). The relationships between use of alcohol, tobacco and coffee in adolescence and mood disorders in adulthood. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36177725>

Chen, N, Liu, H, Yao, J, Chen, S, Xiu, M, Wu, F, & Zhang, X. (2022). Smoke, GPx activity and symptoms improvement in patients with drug-naïve first-episode schizophrenia: A large-scale 12-week follow-up study. *Asian J Psychiatr*, 77, 103267. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36202003>

Ferchiou, A, Szoke, A, Lajnef, M, Bran, M, Racof, R, Schurhoff, F, & Ladea, M. (2022). Schizotypal dimensions are associated with current but not former tobacco consumption. *Encephale*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36266103>

McCarter, K, McKinlay, ML, Cocks, N, Brasier, C, Hayes, L, Baker, AL et al. (2022). The value of compassionate support to address smoking: A qualitative study with people who experience severe mental illness. *Front Psychiatry*, 13, 868032. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36276321>

Zeng, L, Lv, H, Li, J, Xue, R, Liu, X, Zhou, C, & Yu, H. (2022). Cigarette smoking, coffee consumption, alcohol intake, and clozapine metabolism: A Mendelian randomization study. *Front Psychiatry*, 13, 1002235. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36245885>

Mallet, J, Godin, O, Dansou, Y, Mazer, N, Scognamiglio, C, Berna, F et al (2022). Current (but not ex) cigarette smoking is associated with worse cognitive performances in schizophrenia: results from the FACE-SZ cohort. *Psychol Med*, 1-12. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36073848>

Su, MH, Lai, RY, Lin, YF, Chen, CY, Feng, YA, Hsiao, PC, & Wang, SH. (2022). Evaluation of the causal relationship between smoking and schizophrenia in East Asia. *Schizophrenia (Heidelb)*, 8(1), 72. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36085329>

Sawyer, C, Hassan, L, Guinart, D, Agulleiro, LM, & Firth, J. (2022). Smoking Cessation Apps for People with Schizophrenia: How Feasible Are m-Health Approaches? *Behav Sci (Basel)*, 12(8). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36004836>

Kucukaksu, MH, Hoekstra, T, Jansen, L, Vermeulen, J, Adriaanse, MC, & van Meijel, B. (2022). Developing a Smoking Cessation Intervention for People With Severe Mental Illness Treated by Flexible Assertive Community Treatment Teams in the Netherlands: A Delphi Study. *Front Psychiatry*, 13, 866779. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35873255>

Rogers, RT, Marino, L, Scodes, J, Wall, M, Nossel, I, & Dixon, L. (2022). Prevalance and course of tobacco use among individuals receiving coordinated specialty care for first-episode psychosis. *Early Interv Psychiatry*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35751414>

9A.3.5.7.1 Bipolar disorder

Anthenelli, RM, McKenna, BS, Giannini, J, Attaluri, SV, Rubin, M, O'Crowley, E et al. (2023). Combining varenicline preloading with Acceptance and Commitment Therapy (ACT) in persons with serious mental illness who smoke: The randomized ACTS_{Low} pilot feasibility trial. *Drug Alcohol Depend*, 253, 111012. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37931328>

Kertes J, Stein Reisner O, Grunhaus L, Nezry R, Alcalay T, et al. Comparison of smoking cessation program registration, participation, smoking cessation medication utilization, and abstinence rates between smokers with and without schizophrenia, schizo-affective disorder, or bipolar disorder.

Nicotine & Tobacco Research, 2022; 24(5):670–8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34626108>

Vermeulen JM, Wootton RE, Treur JL, Sallis HM, Jones HJ, et al. Smoking and the risk for bipolar disorder: Evidence from a bidirectional mendelian randomisation study. *The British Journal of Psychiatry*, 2021; 218(2):88–94. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31526406>

Oliveira MEC, Almeida NL, Fernandes TP, and Santos NA. Relation between smoking and visual processing in bipolar disorder. *Journal of Addictive Diseases*, 2021:1-7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34075846>

Khobragade B, Sharma V, and Deshpande SN. A study on tobacco use in women with major mental illnesses- schizophrenia, bipolar disorder and recurrent depression. *Psychiatry Res*, 2020; 290:113121. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32505927>

Heffner JL, Kelly MM, Waxmonsky J, Mattocks K, Serfozo E, et al. Pilot randomized controlled trial of web-delivered acceptance and commitment therapy versus smokefree.Gov for smokers with bipolar disorder. *Nicotine & Tobacco Research*, 2020; 22(9):1543–52. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31883336>

Pearsall R, Smith DJ, and Geddes JR. Pharmacological and behavioural interventions to promote smoking cessation in adults with schizophrenia and bipolar disorders: A systematic review and meta-analysis of randomised trials. *BMJ Open*, 2019; 9(11):e027389. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31784428>

Icick R, Melle I, Etain B, Ringen PA, Aminoff SR, et al. Tobacco smoking and other substance use disorders associated with recurrent suicide attempts in bipolar disorder. *J Affect Disord*, 2019; 256:348–57. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31202989>

Heffner JL, Evins AE, Russ C, Lawrence D, Ayers CR, et al. Safety and efficacy of first-line smoking cessation pharmacotherapies in bipolar disorders: Subgroup analysis of a randomized clinical trial. *J Affect Disord*, 2019; 256:267–77. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31195244>

Sarramea F, Jaen-Moreno MJ, Balanza-Martinez V, Osuna MI, Alcala JA, et al. Setting the stage to quit smoking in bipolar disorder patients: Brief advice in clinical practice. *Adicciones*, 2018; 0(0):1006. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30059580>

Heffner JL, Watson NL, McClure JB, Anthenelli RM, Hohl S, et al. "I smoke like this to suppress these issues that are flaws of my character": Challenges and facilitators of cessation among smokers with bipolar disorder. *Journal of Dual Diagnosis*, 2018:1-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29351507>

Heffner JL, Mull KE, Watson NL, McClure JB, and Bricker JB. Smokers with bipolar disorder, other affective disorders, and no mental health conditions: Comparison of baseline characteristics and success at quitting in a large 12-month behavioral intervention randomized trial. *Drug and Alcohol Dependence*, 2018; 193:35–41. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30340143>

Ceylan D, Scola G, Tunca Z, Isaacs-Trepanier C, Can G, et al. DNA redox modulations and global DNA methylation in bipolar disorder: Effects of sex, smoking and illness state. *Psychiatry Res*, 2018; 261:589-96. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29407727>

Ter Meulen WG, van Zaane J, Draisma S, Beekman ATF, and Kupka RW. Does the number of previous mood episodes moderate the relationship between alcohol use, smoking and mood in bipolar outpatients? *BMC Psychiatry*, 2017; 17(1):185. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28506220>

Medeiros GC, Lafer B, Kapczinski F, Miranda-Scippa A, and Almeida KM. Bipolar disorder and tobacco smoking: Categorical and dimensional clinical correlates in subjects from the Brazilian bipolar research network. *Compr Psychiatry*, 2017; 82:14-21. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29367058>

Chu CW, Kao CH, and Chan CK. Could the nicotine receptor partial agonist varenicline moderate manic symptoms in heavy smokers with manic episodes? *Australian and New Zealand Journal of Psychiatry*, 2017;4867417747403. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29226691>

Wilens TE, Biederman J, Martelon M, Zulauf C, Anderson JP, et al. Further evidence for smoking and substance use disorders in youth with bipolar disorder and comorbid conduct disorder. *Journal of Clinical Psychiatry*, 2016; 77(10):1420-7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27574842>

Ospina LH, Russo M, Nitzburg GM, Cuesta-Diaz A, Shanahan M, et al. The effects of cigarette smoking behavior and psychosis history on general and social cognition in bipolar disorder. *Bipolar Disorders*, 2016; 18(6):528-38. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27650399>

Icick R, Gard S, Barde M, Carminati M, Desage A, et al. Physical and mental health burden in cases of bipolar disorder classified as current, former, or non-tobacco smokers. *J Affect Disord*, 2016; 208:406-13. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27810725>

9A.3.5.7.2 Schizophrenia

Flanagan, RJ, Obee, SJ, Kim, AHM, & Every-Palmer, S. (2024). Plasma Clozapine and N-Desmethylozapine (Norclozapine) Concentrations and the Clozapine/Norclozapine Ratio: Effect of Dose, Sex, and Cigarette Smoking. *J Clin Psychopharmacol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39173038>

Du, X, Choa, FS, Chiappelli, J, Bruce, H, Kvarta, M, Summerfelt, A et al. (2024). Combining neuroimaging and brain stimulation to test alternative causal pathways for nicotine addiction in schizophrenia. *Brain Stimul*, 17(2), 324-332. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38453003>

Tsukahara, M, So, R, Yoshimura, Y, Nakajima, S, & Takeuchi, H. (2024). Response to the letter to the Editor from Araujo and colleagues on the research article entitled, "Effect of smoking habits and concomitant valproic acid use on relapse in patients with treatment-resistant schizophrenia receiving clozapine: A 1-year retrospective cohort study". *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38359861>

Araujo, MB, Nogueira, GN, Silveira, P, de Souza Junior, SA, de Matos, ESFG, & Bisol, L W. (2024). Insights and improvements: A critical analysis of methodological limitations in the investigation of smoking and valproic acid in treatment-resistant schizophrenia. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38263735>

Lozano, R, & Bona, C. (2023). Influence of Cigarette Smoking Habit on Clozapine-to-Norclozapine Ratio in Male Patients. *J Res Pharm Pract*, 12(1), 29-31. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38213602>

Flanagan, RJ, Hunter, S, & Obee, SJ. (2023). Effect of Cigarette Smoking on Clozapine Dose and on Plasma Clozapine and N-Desmethylozapine (Norclozapine) Concentrations in Clinical Practice. *J Clin Psychopharmacol*, 43(6), 514-519. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37930204>

Hahn, B, & Holzel, ES. (2023). Nicotine insensitivity and cue-controlled smoking behavior in people with schizophrenia. *Schizophr Res*, 262, 51-52. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37922844>

Gao, Z, Xiu, M, Liu, J, Wu, F, & Zhang, X. (2023). Smoking, Symptoms Improvement, and Total Antioxidant Capacity in Patients with Drug-naïve First-episode Schizophrenia: A Prospective Cohort Study. *Curr Neuropsychopharmacol*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37859307>

Tsukahara, M, So, R, Yoshimura, Y, Yamashita, R, Yada, Y, Kodama, M et al. (2023). Effect of smoking habits and concomitant valproic acid use on relapse in patients with treatment-resistant schizophrenia receiving clozapine: A 1-year retrospective cohort study. *Acta Psychiatr Scand*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37681448>

Zhang, M, Tang, J, Li, W, Xue, K, Wang, Z, Chen, Y L. (2023). Schizophrenia mediating the effect of smoking phenotypes on antisocial behavior: A Mendelian randomization analysis. *CNS Neurosci Ther*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37650156>

Song W, Lin GN, Yu S, and Zhao M. Genome-wide identification of the shared genetic basis of cannabis and cigarette smoking and schizophrenia implicates *ncam1* and neuronal abnormality. *Psychiatry Res*, 2022; 310:114453. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35235886>

Moeller SJ, Gil R, Weinstein JJ, Baumvoll T, Wengler K, et al. Deep rtms of the insula and prefrontal cortex in smokers with schizophrenia: Proof-of-concept study. *NPJ Schizophr*, 2022; 8(1):6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35217662>

Johnstone S, Sorkhou M, Rabin RA, and George TP. Dose-dependent effects of varenicline on tobacco craving and withdrawal in tobacco smokers with and without schizophrenia. *Drug and Alcohol Dependence*, 2022; 234:109412. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/35395548>

Whitton AE, Lewandowski KE, and Hall MH. Smoking as a common modulator of sensory gating and reward learning in individuals with psychotic disorders. *Brain Sci*, 2021; 11(12). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34942883>

Shi J, Chen N, Wang Z, Wang F, Tan Y, et al. Cholinergic receptor nicotinic beta 3 subunit polymorphisms and smoking in male chinese patients with schizophrenia. *EC Psychol Psychiatr*, 2021; 10(7):11-23. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34368810>

Rajalu BM, Jayarajan D, Muliya KP, Sharma P, Gandhi S, et al. Non-pharmacological interventions for smoking in persons with schizophrenia spectrum disorders - a systematic review. *Asian J Psychiatr*, 2021; 56:102530. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33465747>

Pinho S, Rocha V, and Vieira-Coelho MA. Effectiveness of multimodal interventions focused on smoking cessation in patients with schizophrenia: A systematic review. *Schizophr Res*, 2021; 231:145–53. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33857662>

Peterson RE, Bigdeli TB, Ripke S, Bacanu SA, Gejman PV, et al. Genome-wide analyses of smoking behaviors in schizophrenia: Findings from the psychiatric genomics consortium. *Journal of Psychiatric Research*, 2021; 137:215–24. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33691233>

Moe AM, Keller-Hamilton B, Lee A, Breitborde NJK, and Ferketich AK. Perceptions of smoking cessation and a novel psychosocial intervention for quitting among individuals with psychosis: A qualitative study. *J Community Psychol*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34021909>

Mak YW, Loke AY, and Leung DYP. Acceptance and commitment therapy versus social support for smoking cessation for people with schizophrenia: A randomised controlled trial. *J Clin Med*, 2021; 10(19). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34640321>

Ding JB and Hu K. Cigarette smoking and schizophrenia: Etiology, clinical, pharmacological, and treatment implications. *Schizophr Res Treatment*, 2021; 2021:7698030. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34938579>

Desalegn D, Abdu Z, and Hajure M. Prevalence of tobacco dependence and associated factors among patients with schizophrenia attending their treatments at southwest ethiopia; hospital-based cross-sectional study. *PLoS One*, 2021; 16(12):e0261154. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34910737>

Dekker TEG, van der Heijden HS, Schirmbeck F, van Amelsvoort T, Bartels-Velthuis AA, et al. The association between smoking behaviour, social cognition and social functioning in patients with a non-affective psychotic disorder: A prospective follow-up study. *Schizophr Res Cogn*, 2021; 26:100206. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34258239>

Breitborde NJK, Keller-Hamilton B, Moe AM, Pine JG, Nelson N, et al. An open trial of metacognitive remediation therapy and pharmacotherapy to promote smoking cessation among individuals with psychotic-spectrum disorders. *J Smok Cessat*, 2021; 2021:6617716. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34429783>

An H, Fan H, Chen S, Qi S, Ma B, et al. Effects of dose, age, sex, body weight, and smoking on plasma concentrations of olanzapine and n-desmethyl olanzapine in inpatients with schizophrenia. *J Clin*

Psychopharmacol, 2021; 41(3):255-9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33814547>

Al-Soufi L, Martorell L, Molto MD, Gonzalez-Penas J, Garcia-Portilla MP, et al. A polygenic approach to the association between smoking and schizophrenia. Addict Biol, 2021:e13104. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34779080>

Zeng LN, Zong QQ, Zhang L, Feng Y, Ng CH, et al. Worldwide prevalence of smoking cessation in schizophrenia patients: A meta-analysis of comparative and observational studies. Asian J Psychiatr, 2020; 54:102190. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32622029>

Wagner E, Oviedo-Salcedo T, Pelzer N, Strube W, Maurus I, et al. Effects of smoking status on remission and metabolic and cognitive outcomes in schizophrenia patients treated with clozapine. Pharmacopsychiatry, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32757178>

Siskind DJ, Wu BT, Wong TT, Firth J, and Kisely S. Pharmacological interventions for smoking cessation among people with schizophrenia spectrum disorders: A systematic review, meta-analysis, and network meta-analysis. Lancet Psychiatry, 2020; 7(9):762–74. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32828166>

Roche DJO, Bennett M, and Weiner E. Pharmacological interventions to address cigarette smoking in people with schizophrenia spectrum disorders. Lancet Psychiatry, 2020; 7(9):723-4. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32828151>

Nguyen AS, Grossman-Kahn R, and Wichser L. Anticipatory clozapine dosing after discharge from inpatient treatment for the patient with comorbid schizophrenia and tobacco use. J Clin Psychopharmacol, 2020; 40(6):618-9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33065717>

Medenblik AM, Mann AM, Beaver TA, Dedert EA, Wilson SM, et al. Treatment outcomes of a multi-component mobile health smoking cessation pilot intervention for people with schizophrenia. Journal of Dual Diagnosis, 2020:1-9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32735514>

Mak YW, Chiang VCL, and Loke AY. Experiences of tobacco use among chinese individuals with schizophrenia in community-based residential settings: A qualitative study. International Journal of Environmental Research and Public Health, 2020; 17(1). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/31906522>

Ma Y, Li J, Xu Y, Wang Y, Yao Y, et al. Identification of 34 genes conferring genetic and pharmacological risk for the comorbidity of schizophrenia and smoking behaviors. Aging (Albany NY), 2020; 12(3):2169-225. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32012119>

Langlois S, Zern A, Kelley ME, and Compton MT. Adversity in childhood/adolescence and premorbid tobacco, alcohol, and cannabis use among first-episode psychosis patients. Early Interv Psychiatry, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33289325>

Kozak K and George TP. Pharmacotherapy for smoking cessation in schizophrenia: A systematic review. *Expert Opin Pharmacother*, 2020; 21(5):581–90. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32011186>

King M, Jones R, Petersen I, Hamilton F, and Nazareth I. Cigarette smoking as a risk factor for schizophrenia or all non-affective psychoses. *Psychological Medicine*, 2020:1-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32148211>

Kanter Bax O, Hakim N, Jeggo M, Phelan D, Stevens T, et al. Improving smoking cessation in first episode psychosis: A quality improvement project by the city & hackney early and quick intervention psychosis (equip). *BMJ Open Qual*, 2020; 9(4). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33303490>

Hunter A, Murray R, Asher L, and Leonardi-Bee J. The effects of tobacco smoking, and prenatal tobacco smoke exposure, on risk of schizophrenia: A systematic review and meta-analysis. *Nicotine & Tobacco Research*, 2020; 22(1):3–10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30102383>

Fan YS, Yang S, Li Z, Li J, Guo X, et al. A temporal chronnectomic framework: Cigarette smoking preserved the prefrontal dysfunction in schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry*, 2020; 99:109860. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31927054>

Evins AE, West R, Benowitz NL, Russ C, Lawrence D, et al. Efficacy and safety of pharmacotherapeutic smoking cessation aids in schizophrenia spectrum disorders: Subgroup analysis of eagles. *Psychiatric Services*, 2020:appips202000032. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33138708>

Coustals N, Martelli C, Brunet-Lecomte M, Petillion A, Romeo B, et al. Chronic smoking and cognition in patients with schizophrenia: A meta-analysis. *Schizophr Res*, 2020; 222:113–21. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32507373>

Caponnetto P, Polosa R, Bauld L, and Robson D. A qualitative study of the views about smoking, licensed cessation aids and e-cigarettes in people with schizophrenia spectrum disorders. *Journal of Addictive Diseases*, 2020:1-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33043838>

Caponnetto P and Polosa R. Erratum: Tobacco smoking, related harm and motivation to quit smoking in people with schizophrenia spectrum disorders. *Health Psychol Res*, 2020; 8(3):9607. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33553796>

Caponnetto P and Polosa R. Approved and emerging smoking cessation treatments for people with schizophrenia spectrum disorders: A narrative review. *Health Psychol Res*, 2020; 8(2):9237. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33123649>

Brunette MF, Ferron JC, McGurk SR, Williams JM, Harrington A, et al. Brief, web-based interventions to motivate smokers with schizophrenia: Randomized trial. *JMIR Ment Health*, 2020; 7(2):e16524. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32039811>

Yang S, Meng Y, Li J, Fan YS, Du L, et al. Temporal dynamic changes of intrinsic brain activity in schizophrenia with cigarette smoking. *Schizophr Res*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31239219>

Whitton AE, Green AI, Pizzagalli DA, Roth RM, Williams JM, et al. Potent dopamine d2 antagonists block the reward-enhancing effects of nicotine in smokers with schizophrenia. *Schizophrenia Bulletin*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30690638>

Wang YY, Wang S, Zheng W, Zhong BL, Ng CH, et al. Cognitive functions in smoking and non-smoking patients with schizophrenia: A systematic review and meta-analysis of comparative studies. *Psychiatry Res*, 2019; 272:155–63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30583258>

Vermeulen J, Schirmbeck F, Blankers M, van Tricht M, van den Brink W, et al. Smoking, symptoms, and quality of life in patients with psychosis, siblings, and healthy controls: A prospective, longitudinal cohort study. *Lancet Psychiatry*, 2019; 6(1):25-34. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30527763>

Stolz PA, Wehring HJ, Liu F, Love RC, Ellis M, et al. Effects of cigarette smoking and clozapine treatment on 20-year all-cause & cardiovascular mortality in schizophrenia. *Psychiatr Q*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30632082>

Sagud M, Mihaljevic Peles A, and Pivac N. Smoking in schizophrenia: Recent findings about an old problem. *Curr Opin Psychiatry*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31135490>

Ryan JE, Veliz P, McCabe SE, Stoddard SA, and Boyd CJ. Association of early onset of cannabis, cigarette, other drug use and schizophrenia or psychosis. *Schizophr Res*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31623965>

Miller AP, Gizer IR, Fleming Iii WA, Otto JM, Deak JD, et al. Polygenic liability for schizophrenia predicts shifting-specific executive function deficits and tobacco use in a moderate drinking community sample. *Psychiatry Res*, 2019; 279:47-54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31299563>

Mak YW, Loke AY, and Chiang VC. Co-occurrence of schizophrenia and smoking: A qualitative study. *Hong Kong Med J*, 2019; 25 Suppl 2:21-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30674704>

Lally J, Spaducci G, Gardner-Sood P, Atakan Z, Greenwood K, et al. Tobacco smoking and nicotine dependence in first episode and established psychosis. *Asian J Psychiatr*, 2019; 43:125-31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31132542>

Krieger I, Tzur Bitan D, Comaneshter D, Cohen A, and Feingold D. Increased risk of smoking-related illnesses in schizophrenia patients: A nationwide cohort study. *Schizophr Res*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31395488>

Kozak K, Lowe DJE, and George TP. Effects of tobacco smoking status on verbal learning and memory in patients with schizophrenia and non-psychiatric controls. *American Journal on Addictions*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31291035>

Inami R and Kirino E. Nicotine effect on mismatch negativity in smoking and nonsmoking patients with schizophrenia. *Schizophr Res*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30630703>

Huang H, Dong M, Zhang L, Zhong BL, Ng CH, et al. Psychopathology and extrapyramidal side effects in smoking and non-smoking patients with schizophrenia: Systematic review and meta-analysis of comparative studies. *Prog Neuropsychopharmacol Biol Psychiatry*, 2019; 92:476–82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30794823>

Higuchi Y, Fujiwara M, Nakaya N, Fujimori M, Hayashibara C, et al. Change in smoking cessation stage over 1 year in patients with schizophrenia: A follow up study in Japan. *BMC Psychiatry*, 2019; 19(1):367. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31752799>

Hidese S, Matsuo J, Ishida I, Hiraishi M, Teraishi T, et al. Association between lower estimated premorbid intelligence quotient and smoking behavior in patients with schizophrenia. *Schizophr Res Cogn*, 2019; 15:7-13. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30310770>

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6176847/pdf/main.pdf>

Hassanzadah M, Bitar AH, Khanfar NM, Khasawneh FT, Lutfy K, et al. A retrospective cohort study of the prevalence of anxiety and agitation in schizophrenic smokers and the unmet needs of smoking cessation programs. *Medicine (Baltimore)*, 2019; 98(40):e17375. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31577741>

Fang Y, Wang W, Zhu C, Lin GN, Cheng Y, et al. Use of tobacco in schizophrenia: A double-edged sword. *Brain Behav*, 2019:e01433. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31605440>

Cocks N, Brophy L, Segan C, Stratford A, Jones S, et al. Psychosocial factors affecting smoking cessation among people living with schizophrenia: A lived experience lens. *Front Psychiatry*, 2019; 10:565. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31474884>

Castle D, Baker AL, and Bonevski B. Editorial: Smoking and schizophrenia. *Front Psychiatry*, 2019; 10:738. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31681044>

Barkhuizen W, Taylor MJ, Freeman D, and Ronald A. A twin study on the association between psychotic experiences and tobacco use during adolescence. *J Am Acad Child Adolesc Psychiatry*, 2019; 58(2):267-76 e8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30738553>

Erratum: Influence of dose, gender, and cigarette smoking on clozapine plasma concentrations [corrigendum]. *Neuropsychiatr Dis Treat*, 2019; 15:655. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30880990>

Ward HB, Lawson MT, Addington J, Bearden CE, Cadenhead KS, et al. Tobacco use and psychosis risk in persons at clinical high risk. *Early Interv Psychiatry*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30362261>

<https://onlinelibrary.wiley.com/doi/abs/10.1111/eip.12751>

Vermeulen JM, Schirmbeck F, Blankers M, van Tricht M, Bruggeman R, et al. Association between smoking behavior and cognitive functioning in patients with psychosis, siblings, and healthy control subjects: Results from a prospective 6-year follow-up study. *American Journal of Psychiatry*, 2018:appiajp201818010069. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30138044>

Vanderhoof TS, Gurvits TV, Baker-Nolan JE, Borsook D, and Elman I. Visuospatial and sensory integration tasks in patients with schizophrenia or schizoaffective disorder: Relationship to body mass index and smoking. *Front Psychiatry*, 2018; 9:473. Available from:

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6198087/pdf/fpsyt-09-00473.pdf>

Shawen AE and Drayton SJ. Review of pharmacotherapy for smoking cessation in patients with schizophrenia. *Ment Health Clin*, 2018; 8(2):78–85. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29955550>

Scott JG, Matuschka L, Niemela S, Miettunen J, Emmerson B, et al. Evidence of a causal relationship between smoking tobacco and schizophrenia spectrum disorders. *Front Psychiatry*, 2018; 9:607.

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30515111>

Sagud M, Vuksan-Cusa B, Jaksic N, Mihaljevic-Peles A, Rojnic Kuzman M, et al. Smoking in schizophrenia: An updated review. *Psychiatr Danub*, 2018; 30(Suppl 4):216–23. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29864763>

Reeves LE, Gaudiano BA, Metrik J, Guzman Holst C, Morena A, et al. Comorbid cannabis and tobacco use disorders in hospitalized patients with psychotic-spectrum disorders. *Journal of Dual Diagnosis*, 2018;1-10. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30265850>

Rabin RA, Dermody SS, and George TP. Changes in tobacco consumption in cannabis dependent patients with schizophrenia versus non-psychiatric controls during 28-days of cannabis abstinence. *Drug and Alcohol Dependence*, 2018; 185:181-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29459327>

Potvin S, Dugre JR, Fahim C, and Dumais A. Increased connectivity between the nucleus accumbens and the default mode network in patients with schizophrenia during cigarette cravings. *Journal of Dual Diagnosis*, 2018;1-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30445892>

Okoli CTC, El-Mallakh P, and Seng S. Which types of tobacco treatment interventions work for people with schizophrenia? Provider and mental health consumer perspectives. *Issues Ment Health Nurs*, 2018;1-10. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30388915>

Ohi K, Shimada T, Kuwata A, Kataoka Y, Okubo H, et al. Smoking rates and number of cigarettes smoked per day in schizophrenia: A large cohort meta-analysis in a Japanese population. *Int J Neuropsychopharmacol*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30239793>

Ohi K, Kuwata A, Shimada T, Kataoka Y, Yasuyama T, et al. Genome-wide variants shared between smoking quantity and schizophrenia on 15q25 are associated with chrna5 expression in the brain. *Schizophrenia Bulletin*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30202994>

Mallet J, Mazer N, Dubertret C, and Le Strat Y. Tobacco smoking and psychotic-like experiences in a general population sample. *Journal of Clinical Psychiatry*, 2018; 79(6). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30289630>

<http://www.psychiatrist.com/JCP/article/Pages/2018/v79/17m11994.aspx>

Mallet J, Le Strat Y, Schurhoff F, Mazer N, Portalier C, et al. Tobacco smoking is associated with antipsychotic medication, physical aggressiveness, and alcohol use disorder in schizophrenia: Results from the face-sz national cohort. *European Archives of Psychiatry and Clinical Neuroscience*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29396753>

Lyu X, Du J, Zhan G, Wu Y, Su H, et al. Naltrexone and bupropion combination treatment for smoking cessation and weight loss in patients with schizophrenia. *Front Pharmacol*, 2018; 9:181. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29563871>

Lum A, Skelton E, Wynne O, and Bonevski B. A systematic review of psychosocial barriers and facilitators to smoking cessation in people living with schizophrenia. *Front Psychiatry*, 2018; 9:565. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30459658>

Lucatch AM, Lowe DJE, Clark RC, Kozak K, and George TP. Neurobiological determinants of tobacco smoking in schizophrenia. *Front Psychiatry*, 2018; 9:672. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30574101>

Liao W, Yang S, Li J, Fan YS, Duan X, et al. Nicotine in action: Cigarette smoking modulated homotopic functional connectivity in schizophrenia. *Brain Imaging Behav*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30511111>

Liao W, Fan YS, Yang S, Li J, Duan X, et al. Preservation effect: Cigarette smoking acts on the dynamic of influences among unifying neuropsychiatric triple networks in schizophrenia. *Schizophrenia Bulletin*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30561724>

Jones HJ, Gage SH, Heron J, Hickman M, Lewis G, et al. Association of combined patterns of tobacco and cannabis use in adolescence with psychotic experiences. *JAMA Psychiatry*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29344610>

Elkholy H, Nagy N, Taha GRA, Elhabiby M, Yosef M, et al. Stop turning a blind eye: Tobacco smoking among egyptian patients with schizophrenia. *Front Psychiatry*, 2018; 9:703. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30687137>

Djordjevic N, Radmanovic B, Cukic J, Baskic D, Djukic-Dejanovic S, et al. Cigarette smoking and heavy coffee consumption affecting response to olanzapine: The role of genetic polymorphism. *World J Biol Psychiatry*, 2018;1-53. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30513034>

Boggs DL, Surti TS, Esterlis I, Pittman B, Cosgrove K, et al. Minimal effects of prolonged smoking abstinence or resumption on cognitive performance challenge the "self-medication" hypothesis in schizophrenia. *Schizophr Res*, 2018; 194:62–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28392208>

Baker AL, Robson D, Lawn S, Steinberg ML, Bucci S, et al. Reducing smoking among people with schizophrenia: Perspectives on priorities for advancing research. *Front Psychiatry*, 2018; 9:711. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30618881>

Baker AL, Richmond R, Kay-Lambkin FJ, Fila SL, Castle D, et al. Randomised controlled trial of a healthy lifestyle intervention among smokers with psychotic disorders: Outcomes to 36 months.

Australian and New Zealand Journal of Psychiatry, 2018; 52(3):239–52. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28610482>

Ahmed S, Virani S, Kotapati VP, Bachu R, Adnan M, et al. Efficacy and safety of varenicline for smoking cessation in schizophrenia: A meta-analysis. Front Psychiatry, 2018; 9:428. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30283363>

Aguocha C, Uwakwe R, Olose E, Amadi K, Onyeama G, et al. Clinical implication of smoking among patients with schizophrenia at a tertiary institution in south east Nigeria. Afr Health Sci, 2018; 18(1):102-10. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29977263>

Wehring HJ, Heishman SJ, McMahon RP, Liu F, Feldman S, et al. Antipsychotic treatment and tobacco craving in people with schizophrenia. Journal of Dual Diagnosis, 2017:0. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28166471>

Schuster RM, Cather C, Pachas GN, Zhang H, Cieslak KM, et al. Predictors of tobacco abstinence in outpatient smokers with schizophrenia or bipolar disorder treated with varenicline and cognitive behavioral smoking cessation therapy. Addictive Behaviors, 2017; 71:89-95. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28285208>

Rey R, D'Amato T, Boyer L, Brunel L, Aouizerate B, et al. Nicotine dependence is associated with depression and childhood trauma in smokers with schizophrenia: Results from the face-sz dataset. European Archives of Psychiatry and Clinical Neuroscience, 2017. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28389889>

Potvin S, Tikasz A, Lungu O, Stip E, Zaharieva V, et al. Impaired coupling between the dorsomedial prefrontal cortex and the amygdala in schizophrenia smokers viewing anti-smoking images. Front Psychiatry, 2017; 8:109. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28674507>

Perkins KA, Roy Chengappa KN, Karelitz JL, Boldry MC, Michael V, et al. Initial cross-over test of a positive allosteric modulator of alpha-7 nicotinic receptors to aid cessation in smokers with or without schizophrenia. Neuropsychopharmacology, 2017. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29185480>

Moran LV, Betts JM, Ongur D, and Janes AC. Neural responses to smoking cues in schizophrenia. Schizophrenia Bulletin, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29106683>

Miyauchi M, Kishida I, Suda A, Shiraishi Y, Fujibayashi M, et al. Long term effects of smoking cessation in hospitalized schizophrenia patients. BMC Psychiatry, 2017; 17(1):87. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28270120>

Mallet J, Le Strat Y, Schurhoff F, Mazer N, Portalier C, et al. Cigarette smoking and schizophrenia: A specific clinical and therapeutic profile? Results from the face-schizophrenia cohort. Prog Neuropsychopharmacol Biol Psychiatry, 2017; 79(Pt B):332-9. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28663115>

Liu XA and Kenny PJ. Alpha5 nicotinic receptors link smoking to schizophrenia. Nat Med, 2017; 23(3):277-8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28267713>

Kozak K, Dermody SS, Rabin RA, Zack M, Barr MS, et al. Effects of varenicline on cognitive function in non-smokers with schizophrenia. *Schizophr Res*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28291692>

Kamp D, Engelke C, Wobrock T, Kunze B, Wolwer W, et al. Letter to the editor: Influence of rtms on smoking in patients with schizophrenia. *Schizophr Res*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28578924>

Hickling LM, Perez-Iglesias R, Ortiz-Garcia de la Foz V, Balanza-Martinez V, McGuire P, et al. Tobacco smoking and its association with cognition in first episode psychosis patients. *Schizophr Res*, 2017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28412088>

Haig GM, Wang D, Zhao J, Othman AA, and Bain EE. Efficacy and safety of the alpha7-nicotinic acetylcholine receptor agonist abt-126 in the treatment of cognitive impairment associated with schizophrenia: Results from a phase 2b randomized controlled study in smokers. *Journal of Clinical Psychiatry*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28922590>

Gee SH, Taylor DM, Shergill SS, Flanagan R, and MacCabe JH. Effects of a smoking ban on clozapine plasma concentrations in a nonsecure psychiatric unit. *Ther Adv Psychopharmacol*, 2017; 7(2):79-83. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28255437>

Ferron JC, Brunette MF, Geiger P, Marsch LA, Adachi-Mejia AM, et al. Mobile phone apps for smoking cessation: Quality and usability among smokers with psychosis. *JMIR Hum Factors*, 2017; 4(1):e7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28258047>

Cather C, Pachas GN, Cieslak KM, and Evins AE. Achieving smoking cessation in individuals with schizophrenia: Special considerations. *CNS Drugs*, 2017; 31(6):471–81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28550660>

Cather C, Hoepfner S, Pachas G, Pratt S, Achtyes E, et al. Improved depressive symptoms in adults with schizophrenia during a smoking cessation attempt with varenicline and behavioral therapy. *Journal of Dual Diagnosis*, 2017:1-11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28414583>

Li Y, Hou CL, Ma XR, Zang Y, Jia FJ, et al. Smoking and its associations with sociodemographic and clinical characteristics and quality of life in patients with schizophrenia treated in primary care in China. *General Hospital Psychiatry*, 2016; 38:79–83. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26633862>

Kowalczyk WJ, Wehring HJ, Burton G, Raley H, Feldman S, et al. Predictors of the perception of smoking health risks in smokers with or without schizophrenia. *Journal of Dual Diagnosis*, 2016:0. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27858591>

Huang W, Shen F, Zhang J, and Xing B. Effect of repetitive transcranial magnetic stimulation on cigarette smoking in patients with schizophrenia. *Shanghai Arch Psychiatry*, 2016; 28(6):309-17. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28638206>

Grossman M, Bowie CR, Lepage M, Malla AK, Joober R, et al. Smoking status and its relationship to demographic and clinical characteristics in first episode psychosis. *Journal of Psychiatric Research*, 2016; 85:83-90. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27863280>

Mortimer J and Kulkarni J. A balancing act: Clozapine adjustment in smokers. *Australian and New Zealand Journal of Psychiatry*, 2015. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26063698>

Kishi T and Iwata N. Varenicline for smoking cessation in people with schizophrenia: Systematic review and meta-analysis. *European Archives of Psychiatry and Clinical Neuroscience*, 2015; 265(3):259–68. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25283510>

Schuck RK, Dahl A, Hall SM, Delucchi K, Fromont SC, et al. Smokers with serious mental illness and requests for nicotine replacement therapy post-hospitalisation. *Tobacco Control*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25209524>

Rabin RA, Giddens JL, and George TP. Relationship between tobacco and cannabis use status in outpatients with schizophrenia. *American Journal on Addictions*, 2014; 23(2):170–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25187052>

Prochaska JJ, Fromont SC, Delucchi K, Young-Wolff KC, Benowitz NL, et al. Multiple risk-behavior profiles of smokers with serious mental illness and motivation for change. *Health Psychology*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24467257>

Prikryl R, Ustohal L, Kucerovala HP, Kasperek T, Jarkovsky J, et al. Repetitive transcranial magnetic stimulation reduces cigarette consumption in schizophrenia patients. *Prog Neuropsychopharmacol Biol Psychiatry*, 2014; 49:30–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24211840>

Mendelsohn C. Interaction between clopidogrel and smoking. *Australian Family Physician*, 2014; 43(8):503. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25252366>

Lee J, Green MF, Calkins ME, Greenwood TA, Gur RE, et al. Verbal working memory in schizophrenia from the consortium on the genetics of schizophrenia (cogs) study: The moderating role of smoking status and antipsychotic medications. *Schizophr Res*, 2014. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25248939>

Leal-Micharet A, Vaz-Leal FJ, Mendez-Salguero MJ, Mendez-Sanchez F, Baltasar-Tello I, et al. Relationship between smoking and psychotic symptoms in a patient treated with oral olanzapine. A case report. *Actas Esp Psiquiatr*, 2014; 42(5):263–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25179099>

Diehl A, Reinhard I, Schmitt A, Mann K, and Gattaz W. Does the degree of smoking effect the severity of tardive dyskinesia? A longitudinal clinical trial. *European Psychiatry : The Journal of the Association of European Psychiatrists*, 2008; 24(1):33–40. Available from: http://www.sciencedirect.com/science?_ob=MIimg&_imagekey=B6VM1-4TCPN2K-3-1&_cdi=6137&_user=10&_orig=search&_coverDate=09%2F06%2F2008&_sk=999999999&view=c&wchp=dGLbVzz-zSkWA&md5=78934785f594a0f50715c80bd6624372&ie=/sdarticle.pdf

Hirasawa-Fujita M, Bly MJ, Ellingrod VL, Dalack GW, and Domino EF. Genetic variation of the mu opioid receptor (oprm1) and dopamine d2 receptor (drd2) is related to smoking differences in patients with schizophrenia but not bipolar disorder. *Clin Schizophr Relat Psychoses*; 11(1):39-48. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28548579>

9A.3.6 *The role of health professionals and health settings*

Boonstra, N, & Spoelstra, SK. (2024). Addressing Tobacco Use Disorder in Individuals With Mental Health Disorders: The Critical Role of Nurses. *J Adv Nurs*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39584775>

Pleaver, S, Kisely, SR, Bonevski, B, Siskind, D, Guillaumier, A, McCarter, K, & Gartner, C E. (2024). Interventions for smoking cessation in inpatient psychiatry settings. *Cochrane Database Syst Rev*, 9(9), CD015934. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39229858>

Kleinman, RA, & Barnett, BS. (2024). Smoking Cessation as a Priority for Psychiatrists. *JAMA Psychiatry*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/39083329>

Okoli, CTC, Abufarsakh, B, Seng, S, Robertson, H, & Almogheer, Z. (2024). Assessing Tobacco Treatment Engagement in Assertive Community Treatment Programs. *J Am Psychiatr Nurses Assoc*, 10783903241252804. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38747141>

Foo, CYS, Potter, K, Nielsen, L, Rohila, A, Maravic, MC, Schnitzer, K et al. (2024). Implementation of Community Health Worker Support for Tobacco Cessation: A Mixed-Methods Study. *medRxiv*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/38343842>

Britton, M, Rogova, A, Chen, TA, Martinez Leal, I, Kyburz, B, Williams, T et al. (2023). Texas tobacco quitline knowledge, attitudes, and practices within healthcare agencies serving individuals with behavioral health needs: A multimethod study. *Prev Med Rep*, 35, 102256. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37752980>

Tildy, BE, McNeill, A, East, K, Gravely, S, Fong, GT, Cummings, KM et al. (2023). Self-reported depression and anxiety and healthcare professional interactions regarding smoking cessation and nicotine vaping: Findings from 2018 International Tobacco Control Four Country Smoking and Vaping (ITC 4CV) Survey. *Tob Prev Cessat*, 9, 26. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37533461>

Jafry, MZ, Martinez, J, Chen, T A, Britton, M, Martinez Leal, I, Rogova, A et al. (2023). Behavioral health care provider's beliefs, confidence, and knowledge in treating cigarette smoking in relation to their use of the 5A's intervention. *Addict Behav Rep*, 17, 100493. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37347047>

Martinez, C, Lisha, N, McCuistian, C, Straus, E, Delucchi, K, & Guydish, J. (2023). Corrigendum: Comparing client and staff reports on tobacco-related knowledge, attitudes, beliefs and services provided in substance use treatment. *Tob Induc Dis*, 21, 59. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37181461>

Pleaver, S, Kisely, S, Bonevski, B, McCarthy, I, Emmerson, B, Ballard, E et al. (2023). Can improvement in delivery of smoking cessation care be sustained in psychiatry inpatient settings through a system

change intervention? An analysis of statewide administrative health data. *Aust N Z J Psychiatry*, 48674231164566. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37038343>

Schnoll, RA, Leone, FT, Quinn, MH, Stevens, N, Flitter, A, Wileyto, P et al. (2023). A randomized clinical trial testing two implementation strategies to promote the treatment of tobacco dependence in community mental healthcare. *Drug Alcohol Depend*, 247, 109873. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/37084508>

Lobbe, C, Bahnisch, J, Lin, C, Demirkol, A, & Murnion, B. (2023). How do patients and staff in an opioid agonist treatment service view smoking cessation medications and e-cigarettes? *Drug Alcohol Rev*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36877583>

Carter, BJ, Siddiqi, AD, Chen, TA, Britton, M, Martinez Leal, I, Correa-Fernandez, V et al. (2023). Educating Substance Use Treatment Center Providers on Tobacco Use Treatments Is Associated with Increased Provision of Counseling and Medication to Patients Who Use Tobacco. *Int J Environ Res Public Health*, 20(5). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36901024>

Mondon, S, Raich, A, Marti, C, Fernandez, E, Ballbe, M, & La Red Catalana de Hospitales Sin Humo, G. (2023). Training mental health residents in tobacco and alcohol: Relationship with their clinical intervention. *Adicciones*, 0(0), 1837. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36975072>

Siddiqi, AD, Chen, TA, Britton, M, Martinez Leal, I, Carter, BJ, Correa-Fernandez, V et al. (2023). Changes in Substance Use Treatment Providers' Delivery of the 5A's for Non-Cigarette Tobacco Use in the Context of a Comprehensive Tobacco-Free Workplace Program Implementation. *Int J Environ Res Public Health*, 20(3). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36768097>

Jafry, MZ, Reuven, SM, Britton, M, Chen, TA, Martinez Leal, I, Rogova, A et al. (2022). Providers' Non-Cigarette Tobacco Use Intervention Practices in Relation to Beliefs about Patients, Prioritization of and Skills for Intervention, and Referral Knowledge in Texas Healthcare Centers Providing Care to Persons with Behavioral Health Needs. [MS Top Pick]. *Int J Environ Res Public Health*, 19(21). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36361149>

Bellali, T, Karagkounis, C, Liamopoulou, P, Minasidou, E, & Manomenidis, G. (2022). Comparative cross-sectional study of knowledge, attitudes and perceptions among mental health and ward nursing staff towards smoking. *Int J Nurs Pract*, e13108. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36176179>

de Oliveira, RM, Santos, JLF, & Furegato, ARF. (2022). Stages of the Smoke-Free Policy Implementation in a Psychiatric Hospital: Evolution, Effects, and Complications. *Issues Ment Health Nurs*, 1-9. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36227892>

Demir, AC, Donmez, YE, Kartalci, G, Bingol, ME, Temelli, G, & Ozcan, O. (2022). The relationship between smoking, alcohol, and substance abuse and psychiatric diseases among adolescents treated in a child and adolescent psychiatry inpatient unit. *Turk J Pediatr*, 64(5), 816-824. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36305431>

- Sawyer, K, Fredman Stein, K, Jacobsen, P, Freeman, TP, Blackwell, AKM, Metcalfe, C et al. (2022). Acceptability of integrating smoking cessation treatment into routine care for people with mental illness: A qualitative study. *Health Expect*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36222067>
- Shoesmith, E, Huddleston, L, Pervin, J, Shahab, L, Coventry, P, Coleman, T et al. (2022). Promoting and maintaining changes in smoking behaviour for patients following discharge from a smokefree mental health inpatient stay: Development of a complex intervention using the Behaviour Change Wheel. *Nicotine Tob Res*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36250614>
- Cerci, D. (2022). Attitudes of staff towards smoke-free environments in psychiatric hospitals in Germany. *Tob Induc Dis*, 20, 76. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36118555>
- Chawlowiska, E, Karasiewicz, M, Marcinkowska, K, Giernas, B, Jozwiak, P, & Lipiak, A. (2022). Nurses' Perspectives on Smoking Policies, Safety and Cessation Support in Psychiatric Wards: A Cross-Sectional Survey. *Healthcare (Basel)*, 10(9). Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36141347>
- Sourry, RJ, Hyslop, F, Butler, TG, & Richmond, RL. (2022). Impact of smoking bans and other smoking cessation interventions in prisons, mental health and substance use treatment settings: A systematic review of the evidence. *Drug Alcohol Rev*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/36097413>
- Oliveira, RM, Santos, JLF, & Furegato, ARF. (2022). Hospital indicators and inpatient behavior in a psychiatric hospital that implemented the smoking ban. *Rev Lat Am Enfermagem*, 30, e3611. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35920539>
- Japuntich, SJ, Adkins-Hempel, M, Lundtvedt, C, Becker, S J, Helseth, SA, Fu, SS et al. (2022). Implementing Chronic Care Model Treatments for Cigarette Dependence in Community Mental Health Clinics. *J Dual Diagn*, 1-12. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35763554>
- Nestor, S, Carlos, P, Cristina, P, Jose, MR, Ignacio, B, & Pilar, S. (2022). TOBACCO USE DISORDER AND DUAL DISORDERS Joint statement by the Spanish Psychiatry Society and the Spanish Dual Disorders Society. *Actas Esp Psiquiatr*, 50(Supplement), 77-138. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35731182>
- Otachi, JK, Otis, M, Flaherty, C, & Okoli, CTC. (2022). Enhancing Provider Delivery of Tobacco Treatment Within the Inpatient Psychiatric Setting. *Soc Work Public Health*, 1-12. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35762032>
- Young, WJ, Delnevo, CD, Singh, B, Steinberg, ML, Williams, JM, & Steinberg, MB. (2022). Tobacco Treatment Knowledge and Practices Among US Psychiatrists. *Community Ment Health J*. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/35768703>
- Taylor GMJ, Baker AL, Fox N, Kessler DS, Aveyard P, et al. Addressing concerns about smoking cessation and mental health: Theoretical review and practical guide for healthcare professionals. *BJPsych Adv*, 2021; 27(2):85–95. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34513007>

Short B, Giles L, Karageorge A, and Bauer L. Exploring and reorienting psychiatrists' attitudes regarding smoking cessation and its potential to improve mental health outcomes: A pilot study. *Australas Psychiatry*, 2021; 29(6):663–7. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34488489>

Pedersen ALW, Lindekilde CR, Andersen K, Hjorth P, and Gildberg FA. Health behaviours of forensic mental health service users, in relation to smoking, alcohol consumption, dietary behaviours and physical activity-a mixed methods systematic review. *J Psychiatr Ment Health Nurs*, 2021; 28(3):444–61. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32916759>

Pleaver S, McCarthy I, Anzolin M, Emmerson B, Allan J, et al. Queensland smoking care in adult acute mental health inpatient units: Supporting practice change. *Australian and New Zealand Journal of Psychiatry*, 2020; 54(9):919–27. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32375495>

Lappin JM, Thomas D, Curtis J, Blowfield S, Gatsi M, et al. Targeted intervention to reduce smoking among people with severe mental illness: Implementation of a smoking cessation intervention in an inpatient mental health setting. *Medicina (Kaunas)*, 2020; 56(4). Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32344790>

Kagabo R, Gordon AJ, and Okuyemi K. Smoking cessation in inpatient psychiatry treatment facilities: A review. *Addictive Behaviors Reports*, 2020; 11:100255. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32467844>

Johnson SE, Mitrou F, Lawrence D, Zubrick SR, Wolstencroft K, et al. Feasibility of a consumer centred tobacco management intervention in community mental health services in Australia. *Community Ment Health J*, 2020; 56(7):1354–65. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32036516>

Twyman L, Cowles C, Walsberger SC, Baker AL, Bonevski B, et al. 'They're going to smoke anyway': A qualitative study of community mental health staff and consumer perspectives on the role of social and living environments in tobacco use and cessation. *Front Psychiatry*, 2019; 10:503. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31379622>

Skelton E, Tzelepis F, Shakeshaft A, Guillaumier A, Wood W, et al. Integrating smoking cessation care into a medically supervised injecting facility using an organizational change intervention: A qualitative study of staff and client views. *International Journal of Environmental Research and Public Health*, 2019; 16(11). Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31185619>

Davis A, Ngo H, and Coleman M. An evaluation of a pilot specialist smoking cessation clinic in a mental health setting. *Australas Psychiatry*, 2019; 27(3):275–8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30507301>

Skelton E, Tzelepis F, Shakeshaft A, Guillaumier A, McCrabb S, et al. Integrating smoking cessation care in alcohol and other drug treatment settings using an organizational change intervention: A systematic review. *Addiction*, 2018; 113(12):2158–72. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29920839>

Sharma-Kumar R, Meurk C, Ford P, Beere D, and Gartner C. Are Australian smokers with mental illness receiving adequate smoking cessation and harm reduction information? *International Journal of Mental Health Nursing*, 2018; 27(6):1673–88. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29718549>

Sharma R, Meurk C, Bell S, Ford P, and Gartner C. Australian mental health care practitioners' practices and attitudes for encouraging smoking cessation and tobacco harm reduction in smokers with severe mental illness. *International Journal of Mental Health Nursing*, 2018; 27(1):247–57. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/28160384>

Metse AP, Wiggers J, Wye P, and Bowman JA. Patient receipt of smoking cessation care in four Australian acute psychiatric facilities. *International Journal of Mental Health Nursing*, 2018; 27(5):1556–63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29573164>

Malone V, Harrison R, and Daker-White G. Mental health service user and staff perspectives on tobacco addiction and smoking cessation: A meta-synthesis of published qualitative studies. *J Psychiatr Ment Health Nurs*, 2018; 25(4):270–82. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29498459>

Dean TD, Cross W, and Munro I. An exploration of the perspectives of associate nurse unit managers regarding the implementation of smoke-free policies in adult mental health inpatient units. *Issues Ment Health Nurs*, 2018; 39(4):328–36. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29436879>

Magor-Blatch LE and Rugendyke AR. Going smoke-free: Attitudes of mental health professionals to policy change. *J Psychiatr Ment Health Nurs*, 2016; 23(5):290–302. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/27278902>

de Oliveira RM and Furegato AR. The decreasing number of cigarettes during psychiatric hospitalization: Intervention or punishment? *Revista Brasileira de Enfermagem*, 2015; 68(1):69–76, – 83. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25946498>

Schwindt RG, McNelis AM, and Sharp D. Evaluation of a theory-based education program to motivate nursing students to intervene with their seriously mentally ill clients who use tobacco. *Archives of Psychiatric Nursing*, 2014; 28(4):277–83. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/25017562>

Rogers E and Sherman S. Tobacco use screening and treatment by outpatient psychiatrists before and after release of the American psychiatric association treatment guidelines for nicotine dependence. *American Journal of Public Health*, 2014; 104(1):90–5. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/24228666>

Abufarsakh B, Otachi JK, Wang T, Al-Mrayat Y, and Okoli CTC. The impact of a nurse-led service on tobacco treatment provision within a psychiatric hospital: A time series study. *Journal of the American Psychiatric Nurses Association*, 2022:10783903221093582. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/35549464>

Martinez C, Feliu A, Torres N, Nieva G, Pinet C, et al. Acceptability and participation predictors for a pragmatic randomized controlled trial to test a smoking cessation intervention after discharge from mental health wards. *Drug and Alcohol Dependence*, 2022; 234:109390. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/35278807>

Schnitzer K, AhnAllen CG, Beck S, Oliveira Y, Fromson JA, et al. Multidisciplinary barriers to addressing tobacco cessation during an inpatient psychiatric hospitalization. *Addictive Behaviors*, 2021; 120:106988. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34051645>

Olson R, McLay M, Hamm J, and Callaghan RC. Identification of tobacco-related cancer diagnoses among individuals with psychiatric disorders: A population-based matched cohort study using a competing risks approach from British Columbia. *Curr Oncol*, 2021; 28(6):4953-60. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34898588>

McLennan JD. Wondering about the psychiatrist's role in smoking cessation from inpatients to whole populations. *J Can Acad Child Adolesc Psychiatry*, 2021; 30(1):2. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33552166>

McCuistion C, Kapiteni K, Le T, Safier J, Delucchi K, et al. Reducing tobacco use in substance use treatment: An intervention to promote tobacco-free grounds. *Journal of Substance Abuse Treatment*, 2021:108640. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34743925>

Madden BJ and Fogger SA. Implementing clinical practice guidelines for tobacco cessation in a women's residential substance use treatment facility. *Journal of Addictions Nursing*, 2021; 32(2):159-64. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34060768>

Lopez-Lazcano AI, Lopez-Pelayo H, Balcells-Olivero M, Segura L, and Gual Sole A. Validation of the alcohol smoking and substance involvement screening test (assist) in acute psychiatric inpatients. *Adicciones*, 2021; 0(0):1496. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34171103>

Korchia T, Faugere M, Suc N, Garosi A, Andrieu-Haller C, et al. Recommendations of the treatment-resistant depression expert center network for promoting tobacco smoking cessation based on the results from the real-world face-trd national cohort. *Prog Neuropsychopharmacol Biol Psychiatry*, 2021; 114:110479. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34826559>

Kneeland ET, Peckham AD, McHugh RK, Weiss RD, Beard C, et al. Tobacco use predicts treatment dropout and outcome in an acute, transdiagnostic psychiatric treatment setting. *Behavior Therapy*, 2021; 52(4):897-906. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34134829>

Kertes J, Stein Reisner O, Grunhaus L, and Neumark Y. Adherence to smoking cessation guidelines (5as) by psychiatrists and primary care physicians for persons with serious mental illness. *Journal of Addictive Diseases*, 2021:1-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34931584>

Kazan AS, Perry LM, Atiya WF, Voss HM, Kim S, et al. Development and acceptability of an educational video about a smoking cessation quitline for use in adult outpatient mental healthcare. *Transl Behav Med*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33647990>

Jenkin G, McIntosh J, Hoek J, Mala K, Paap H, et al. There's no smoke without fire: Smoking in smoke-free acute mental health wards. *PLoS One*, 2021; 16(11):e0259984. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34780542>

Guydish J, Le T, Hosakote S, Straus E, Wong J, et al. Tobacco use among substance use disorder (sud) treatment staff is associated with tobacco-related services received by clients. *Journal of Substance Abuse Treatment*, 2021; 132:108496. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34111773>

Gucuk S, Erim BR, and Kayhan M. A pre-evaluation of psychological factors may positively affect the outcomes of smoking cessation treatments: A comparison in terms of smoking behavior. *Tob Prev Cessat*, 2021; 7:12. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33598591>

Freiburghaus T, Raffing R, Ballbe M, Gual A, and Tonnesen H. The right to smoke and the right to smoke-free surroundings: International comparison of smoke-free psychiatric clinic implementation experiences. *BJPsych Open*, 2021; 7(3):e81. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33858559>

Fond G, Trouve M, Andrieu-Haller C, Sunhary de Verville PL, and Boyer L. Barriers in psychiatrists' mind to active smoking cessation promotion in severe psychiatric disorders. *Encephale*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34862009>

Falcaro M, Osborn D, Hayes J, Coyle G, Couperthwaite L, et al. Time trends in access to smoking cessation support for people with depression or severe mental illness: A cohort study in english primary care. *BMJ Open*, 2021; 11(12):e048341. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34862277>

Christiansen B, Smith SS, and Fiore MC. Measuring therapeutic alliance for tobacco cessation counseling for behavioral health clinicians. *J Smok Cessat*, 2021; 2021:6671899. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34306231>

Castaldelli-Maia JM, Silva ASM, Goncalves PD, Sanches MV, Lassi DLS, et al. Smoking cessation for individuals living with mental disorders in Brazil: Increasing providers' commitment and impact. *General Hospital Psychiatry*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34961631>

Castaldelli-Maia JM, Goncalves PD, Lima DR, Moura HF, and Apter G. Quality of service and commitment to tobacco dependence treatment for individuals living with mental disorders in france: A pilot study. *Int J Soc Psychiatry*, 2021:207640211042914. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/34465236>

Brown RA, Minami H, Hecht J, Kahler CW, Price LH, et al. Sustained care smoking cessation intervention for individuals hospitalized for psychiatric disorders: The helping hand 3 randomized clinical trial. *JAMA Psychiatry*, 2021. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33950156>

Abufarsakh B and Okoli CTC. Nursing staff attitude, subjective norms, perceived behavior control, and intention to provide tobacco treatment in a psychiatric hospital. *Journal of the American*

Psychiatric Nurses Association, 2021:10783903211045737. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/34538112>

Zern A, Seserman M, Dacus H, Wallace B, Friedlander S, et al. Screening and treatment of tobacco use disorder in mental health clinics in new york state: Current status and potential next steps.

Community Ment Health J, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33083939>

Vermeulen JM and van den Brink W. [smoking in psychiatry: From smoking as the norm to 'just stop it']. Tijdschr Psychiatr, 2020; 62(11):946-8. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33443744>

Vermeulen JM, Doedens P, Schirmbeck F, van den Brink W, and de Haan L. [implementation of a stop-smoking support program for patients and employees in a psychiatric hospital: First results].

Tijdschr Psychiatr, 2020; 62(4):266-73. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32388848>

Taylor GMJ, Sawyer K, Kessler D, Munafo MR, Aveyard P, et al. Views about integrating smoking cessation treatment within psychological services for patients with common mental illness: A multi-perspective qualitative study. Health Expect, 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33368996>

Tancredi N and Kohli N. Smoking cessation and duloxetine toxicity: A case report. J Am Pharm Assoc (2003), 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32094037>

Spaducci G, Richardson S, McNeill A, Pritchard M, Sanyal J, et al. An observational study of system-level changes to improve the recording of very brief advice for smoking cessation in an inpatient mental health setting. BMC Public Health, 2020; 20(1):559. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32334547>

Ratier-Cruz A, Smith JG, Firn M, and Rinaldi M. Staff attitudes to completely smoke-free policies and smoking cessation practices in a mental health setting. J Public Health (Oxf), 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32128592>

Mohiuddin SM, McDermott B, and Dillon J. Disconnect between psychiatric and addiction services: A review of patients' attendances at alcohol tobacco and other drug services after a psychiatric admission. Australas Psychiatry, 2020:1039856220956470. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/33103458>

Le K, Correa-Fernandez V, Leal IM, Kyburz B, Chen TA, et al. Tobacco-free workplace program at a substance use treatment center. American Journal of Health Behavior, 2020; 44(5):652-65. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33121583>

Kuzin M, Schoretsanis G, Haen E, Ridders F, Hiemke C, et al. Pharmacokinetic interactions between clozapine and sertraline in smokers and non-smokers. Basic Clin Pharmacol Toxicol, 2020. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/32307853>

Kelly JF, Greene MC, Hoffman LA, Hoepfner BB, and Bergman BG. On providing smoking cessation services in alcohol and other drug treatment settings: Results from a u.S. National survey of attitudes

among recovering persons. *Journal of Substance Abuse Treatment*, 2020; 117:108057. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32811636>

Jones SE, Mulrine S, Clements H, and Hamilton S. Supporting mental health service users to stop smoking: Findings from a process evaluation of the implementation of smokefree policies into two mental health trusts. *BMC Public Health*, 2020; 20(1):1619. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33115443>

Guydish J, Wahleithner J, Williams D, and Yip D. Tobacco-free grounds implementation in california residential substance use disorder (sud) treatment programs. *Journal of Addictive Diseases*, 2020; 38(1):55-63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/32186480>

Ainscough TS, Mitchell A, Hewitt C, Horspool M, Stewart P, et al. Investigating changes in patients' smoking behaviour, tobacco dependence and motivation to stop smoking following a 'smoke-free' mental health inpatient stay: Results from a longitudinal survey in england. *Nicotine & Tobacco Research*, 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33277655>

Action on Smoking and Health. Smokefree skills: Training needs of mental health nurses and psychiatrists - executive summary. 2020. Available from: https://ash.org.uk/wp-content/uploads/2020/12/ExecSummary_MHTraining_F.pdf.

Williams JM, Steinberg ML, Wang H, Chaguturu V, Poulsen R, et al. Practice change after training psychiatry residents in tobacco use disorder. *Psychiatric Services*, 2019:appips201900272. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31690223>

Taylor G, Aveyard P, Bartlem K, Shaw A, Player J, et al. Integrating smoking cessation treatment as part of usual psychological care for depression and anxiety (escape): Protocol for a randomised and controlled, multicentre, acceptability, feasibility and implementation trial. *Pilot Feasibility Stud*, 2019; 5:16. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30693096>

Spaducci G, McNeill A, Hubbard K, Stewart D, Yates M, et al. Smoking-related violence in a mental health setting following the implementation of a comprehensive smoke-free policy: A content analysis of incident reports. *International Journal of Mental Health Nursing*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31513336>

Smith CA, McNeill A, Kock L, and Shahab L. Exploring mental health professionals' practice in relation to smoke-free policy within a mental health trust: A qualitative study using the com-b model of behaviour. *BMC Psychiatry*, 2019; 19(1):54. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30717722>

Scheeres A, Xhezo R, Julius R, Coffman R, Frisby J, et al. Changes in voluntary admission and restraint use after a comprehensive tobacco-free policy in inpatient psychiatric health facilities. *Substance Abuse*, 2019:1-7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31295085>

Romano I, Costello MJ, Ropp C, Li Y, Sousa S, et al. Evaluating the short-term impact of a tobacco-free policy in an inpatient addiction treatment setting. *Journal of Substance Abuse Treatment*, 2019; 107:50-9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31757265>

Rogers ES and Wysota CN. Tobacco screening and treatment of patients with a psychiatric diagnosis, 2012-2015. American Journal of Preventive Medicine, 2019; 57(5):687-94. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31564608>

Okoli CTC and Seng S. Correlates of tobacco use and consumption among hospitalized psychiatric patients. West J Nurs Res, 2019:193945918823483. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30658562>

Okoli CTC, Otachi JK, and Seng S. Assessing opinions and barriers to providing evidence-based tobacco treatment among health care providers within an in-patient psychiatric facility. J Ment Health, 2019:1-11. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30862266>

No authors listed. Progress towards smokefree mental health services: Findings from a survey of mental health trusts in england. ASH, 2019. Available from: https://ash.org.uk/wp-content/uploads/2019/10/PHE-mental-health-trust-2019-survey-full-report_v12.pdf.

Kleinman RA, Kendra MS, and De Golia SG. Smoking cessation prescribing and referral practices among psychiatry residents. Acad Psychiatry, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31435900>

Kathuria H, Seibert RG, Cobb V, Weinstein ZM, Gowarty M, et al. Patient and physician perspectives on treating tobacco dependence in hospitalized smokers with substance use disorders: A mixed methods study. Journal of Addiction Medicine, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31232740>

Kagabo R, Kim J, Zubietta JK, Kleinschmit K, and Okuyemi K. Association between smoking, and hospital readmission among inpatients with psychiatric illness at an academic inpatient psychiatric facility, 2000-2015. Addictive Behaviors Reports, 2019; 9:100181. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31193775>

Japuntich SJ, Dunne EM, Krieger NH, Ryan PM, Rogers E, et al. Proactive tobacco treatment in a behavioral health home. Community Ment Health J, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31520255>

Holler JT, Borowsky H, and Peterson A. Smoking cessation counseling on inpatient psychiatric units. Psychiatric Services, 2019; 70(12):1180. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31787060>

Hecht J, Rigotti NA, Minami H, Kjome KL, Bloom EL, et al. Adaptation of a sustained care cessation intervention for smokers hospitalized for psychiatric disorders: Study protocol for a randomized controlled trial. Contemporary Clinical Trials, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31212100>

Gubner NR, Williams DD, Le T, Garcia W, Vijayaraghavan M, et al. Smoking related outcomes before and after implementation of tobacco-free grounds in residential substance use disorder treatment programs. Drug and Alcohol Dependence, 2019; 197:8-14. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30743196>

de Oliveira RM, Santos JLF, and Furegato ARF. Opinions of hospitalized patients with mental disorders and patients in basic health units in Brazil regarding tobacco smoking. *Issues Ment Health Nurs*, 2019;1-8. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30620628>

Chavarria J, Liu M, Kast L, Salem E, and King AC. A pilot study of counsel to quit(r): Evaluating an ask advise refer (aar)-based tobacco cessation training for medical and mental healthcare providers. *Journal of Substance Abuse Treatment*, 2019; 99:163-70. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30797390>

Allen AM, Muramoto ML, Campbell J, Connolly TE, McGuffin BA, et al. Multimethod formative research to improve the training and delivery of tobacco-cessation interventions in behavioral health settings. *Journal of Addiction Medicine*, 2019. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30908345>

Action on Smoking and Health. Smokefree skills: Community mental health. ASH, 2019. Available from: <https://ash.org.uk/wp-content/uploads/2019/10/SmokefreeSkillsCommunityMentalHealth.pdf>.

Van de Graaf RC, Veldman K, and Anthonio GG. Reaction to "intergrating smoking cessation care in alcohol and other drug treatment settings using an organisational change intervention: A systematic review" by skelton, et al. *Addiction*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30091225>

Sarramea F, Jaen-Moreno MJ, Feu N, Redondo-Ecija J, Balanza-Martinez V, et al. Prepare the smoking cessation in severe mental illness: Early diagnosis and prevention opportunities. *Rev Psiquiatr Salud Ment*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30274862>

Samples H, Bandara S, Olfson M, and Saloner B. Tobacco screening and counseling in the u.s.: Smokers with mental health and substance use problems. *American Journal of Preventive Medicine*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30135038>

Okoli CTC, Al-Mrayat YD, and Stead B. The effect of implementing a tobacco treatment service on adherence to evidence-based practice in an inpatient state-owned psychiatric hospital. *American Journal on Addictions*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29741217>

Noblett S, Beenstock J, Noblett J, Ireland J, and Ormiston S. Implementing national institute for health and care excellence smoke-free guidance in a secure facility: An evaluation of the prescribing costs in clozapine users. *BJPsych Bull*, 2018; 42(2):59-62. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29544555>

Muladore E, Brown JA, Haefner J, and Kupferschmid B. Improving patient education about tobacco withdrawal and nicotine gum use by registered nurses in inpatient psychiatry: A feasibility study. *J Psychiatr Ment Health Nurs*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30129262>

Metse AP, Hizam NAN, Wiggers J, Wye P, and Bowman JA. Factors associated with retention in a smoking cessation trial for persons with a mental illness: A descriptive study. *BMC Med Res Methodol*, 2018; 18(1):177. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30587149>

Marynak K, VanFrank B, Tetlow S, Mahoney M, Phillips E, et al. Tobacco cessation interventions and smoke-free policies in mental health and substance abuse treatment facilities - United States, 2016. *Morbidity and Mortality Weekly Report*, 2018; 67(18):519-23. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29746451>

Marynak K, VanFrank B, Tetlow S, Mahoney M, Phillips E, et al. Tobacco cessation interventions and smoke-free policies in mental health and substance abuse treatment facilities — United States, 2016. *Morbidity and Mortality Weekly Report (MMWR)*, 67(18);519–523 2018. Available from:

<http://dx.doi.org/10.15585/mmwr.mm6718a3>.

Jegade O, Ojo O, Ahmed S, Kodjo K, Virk I, et al. Tobacco and substance use among psychiatric inpatients in a community hospital: Cessation counseling, correlates, and patterns of use. *J Addict*, 2018; 2018:7919704. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30662786>

Huddlestone L, Sohal H, Paul C, and Ratschen E. Complete smokefree policies in mental health inpatient settings: Results from a mixed-methods evaluation before and after implementing national guidance. *BMC Health Services Research*, 2018; 18(1):542. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/29996855>

Hedges E and Facer-Irwin E. Implementation of a smoke-free policy appears to reduce physical violence in inpatient settings. *Evid Based Ment Health*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30580254>

Graydon MM, Corno CM, Schacht RL, Knoblach DJ, Wiprovnick AE, et al. A statewide initiative to train behavioral health providers in smoking cessation. *Transl Behav Med*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30202855>

Flitter A, Lubitz SF, Ziedonis D, Stevens N, Leone F, et al. A cluster-randomized clinical trial testing the effectiveness of the addressing tobacco through organizational change (attoc) model for improving the treatment of tobacco use in community mental healthcare: Preliminary study feasibility and baseline findings. *Nicotine & Tobacco Research*, 2018. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30388273>

Conrad M, Bolte T, Gaines L, Avery Z, and Bodie L. The untreated addiction: Going tobacco-free in a va substance abuse residential rehabilitation treatment program (sarrrtp). *J Behav Health Serv Res*, 2018. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29721729>

Collins A, Ajayi O, Diamond S, Diamond W, and Holroyd S. Tobacco use and associated factors in patients presenting to a psychiatric emergency room. *J Addict*, 2018; 2018:8102165. Available from:

<https://www.ncbi.nlm.nih.gov/pubmed/30009080>

Chen LS, Baker TB, Korpecki JM, Johnson KE, Hook JP, et al. Low-burden strategies to promote smoking cessation treatment among patients with serious mental illness. *Psychiatric Services*, 2018:appips201700399. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29852824>

Burns A, Webb M, Styne G, O'Brien T, Rohde D, et al. Implementation of a quit smoking programme in community adult mental health services-a qualitative study. *Front Psychiatry*, 2018; 9:670.

Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30622485>

Burns A, Lucey JV, Strawbridge J, Clancy L, and Doyle F. Prospective study of provided smoking cessation care in an inpatient psychiatric setting. *Journal of Psychosomatic Research*, 2018; 115:24-31. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30470313>

Beyraghi N, Mazaheri Meybodi A, and Jafarian Bahri RS. Smoking ban in psychiatric inpatient unit: An Iranian study on the views and attitudes of the mental health staff and psychiatric patients. *Psychiatry J*, 2018; 2018:2450939. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/30310814>
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6166367/pdf/PSYCHIATRY2018-2450939.pdf>

van der Eijk Y. The convention on the rights of persons with disabilities as a tobacco control tool in the mental health setting. *Tobacco Control*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29123024>

Samaha HL, Correa-Fernandez V, Lam C, Wilson WT, Kyburz B, et al. Addressing tobacco use among consumers and staff at behavioral health treatment facilities through comprehensive workplace programming. *Health Promotion Practice*, 2017; 18(4):561-70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28629277>

Rogers ES, Gillespie C, Smelson D, and Sherman SE. A qualitative evaluation of mental health clinic staff perceptions of barriers and facilitators to treating tobacco use. *Nicotine & Tobacco Research*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29059344>

Perry BI, Meehan K, and Jainer AK. Assessing the second-hand effects of a new no-smoking policy in an acute mental health trust. *BJPsych Bull*, 2017; 41(6):325-30. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29234509>

Okoli CTC, Otachi JK, Manuel A, and Woods M. A cross-sectional analysis of factors associated with the intention to engage in tobacco treatment among inpatients in a state psychiatric hospital. *J Psychiatr Ment Health Nurs*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28976063>

Okoli CTC, Otachi JK, Kaewbua S, Woods M, and Robertson H. Factors associated with staff engagement in patients' tobacco treatment in a state psychiatric facility. *Journal of the American Psychiatric Nurses Association*, 2017; 23(4):268-78. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28398833>

Okoli CT, Mason DA, Brumley-Shelton A, and Robertson H. Providing tobacco treatment in a community mental health setting: A pilot study. *Journal of Addictions Nursing*, 2017; 28(1):34-41. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28252509>

Okoli CT, Anand V, and Khara M. A retrospective analysis of the outcomes of smoking cessation pharmacotherapy among persons with mental health and substance use disorders. *Journal of Dual Diagnosis*, 2017;0. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28166473>

Fallin-Bennett A, Parker KA, Miller A, Ashford K, and Hahn EJ. Smoking and tobacco-free policies in women's residential substance use disorder treatment facilities: A community engaged approach. *Nicotine & Tobacco Research*, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29059449>

Castaldelli-Maia JM, da Silva NR, Campos MR, Moura HF, Zabert G, et al. Implementing evidence-based smoking cessation treatment in psychosocial care units (caps) in Brazil. *Int J Soc Psychiatry*, 2017;20764017731462. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28920521>

Aschbrenner KA, Dixon LB, Naslund JA, Bienvenida JCM, McManus KL, et al. An online survey of family members' beliefs and attitudes about smoking and mental illness. *Journal of Dual Diagnosis*, 2017;1-5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28481179>

Alderman CP. Pharmacological considerations related to smoke-free inpatient environments. *Australas Psychiatry*, 2017; 25(5):523. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28992788>

Viron M. Smoke free program at massachusetts mental health center: A peer-supported approach. *Psychiatric Services*, 2016; 67(6):696–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27245084>

Soyster P, Anzai NE, Fromont SC, and Prochaska JJ. Correlates of nicotine withdrawal severity in smokers during a smoke-free psychiatric hospitalization. *Preventive Medicine*, 2016; 92:176-82. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26892910>

Sohal H, Huddleston L, and Ratschen E. Preparing for completely smoke-free mental health settings: Findings on patient smoking, resources spent facilitating smoking breaks, and the role of smoking in reported incidents from a large mental health trust in england. *International Journal of Environmental Research and Public Health*, 2016; 13(3). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26927143>

Simioni N and Cottencin O. Screening for eating disorders in outpatient smoking cessation: Feasibility, pertinence, and acceptance of referral to specific treatment. *Int J Eat Disord*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27218668>

Schroeder SA. Smoking cessation should be an integral part of serious mental illness treatment. *World Psychiatry*, 2016; 15(2):175–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27265708>

Oliveira RM, Duarte AF, Alves D, and Furegato AR. Development of the tabacoquest app for computerization of data collection on smoking in psychiatric nursing. *Rev Lat Am Enfermagem*, 2016; 24:e2726. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27579930>

Lawn S and Lucas T. Addressing smoking in supported residential facilities for people with severe mental illness: Has any progress been achieved? *International Journal of Environmental Research and Public Health*, 2016; 13(10). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27735881>

Du Plooy JL, Macharia M, and Verster C. Cigarette smoking, nicotine dependence, and motivation to quit smoking in south african male psychiatric inpatients. *BMC Psychiatry*, 2016; 16(1):403. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/27852297>

Dixon S and Francisco I. Smoke-free acute psychiatric settings are a safe and ethical public health intervention. *Australas Psychiatry*, 2016; 24(5):511. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27683860>

Christiansen BA, Macmaster DR, Heiligenstein EL, Glysch RL, Riemer DM, et al. Measuring the integration of tobacco policy and treatment into the behavioral health care delivery system: How are we doing? *Journal of Health Care for the Poor and Underserved*, 2016; 27(2):510–26. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27180692>

Blankers M, Buisman R, Hopman P, van Gool R, and van Laar M. Modelling intentions to provide smoking cessation support among mental health professionals in the netherlands. *Tobacco Induced Diseases*, 2016; 14(1):32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27570503>

Bizzarri JV, Casetti V, Panzani P, Unterhauser J, Mulas S, et al. Risky use and misuse of alcohol and cigarettes in psychiatric inpatients: A screening questionnaire study. *Compr Psychiatry*, 2016; 70:9–16. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27624418>

Ballbe M, Gual A, Nieva G, Salto E, Fernandez E, et al. Deconstructing myths, building alliances: A networking model to enhance tobacco control in hospital mental health settings. *Gac Sanit*, 2016. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27325123>

Muilenburg JL, Laschober TC, and Eby LT. Substance use disorder counselors' reports of tobacco cessation services availability, implementation, and tobacco-related knowledge. *The Journal of Adolescent Health*, 2015; 57(3):327–33. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26299559>

Marshall LL, Kuiper NM, and Lavinghouze SR. Strategies to support tobacco cessation and tobacco-free environments in mental health and substance abuse facilities. *Preventing Chronic Disease*, 2015; 12:E167. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26425871>

Liu K and Creamer A. Improving smoking cessation policy by assessing user demand for an inpatient smoking cessation service in adult psychiatric wards. *BMJ Qual Improv Rep*, 2015; 4(1). Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26734337>

Christofferson DE, Hamlett-Berry K, and Augustson E. Suicide prevention referrals in a mobile health smoking cessation intervention. *American Journal of Public Health*, 2015; 105(8):e7–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26066949>

Britton J. Treating smoking in mental health settings. *Lancet Psychiatry*, 2015; 2(5):364–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26360261>

Williams JM, Willett JG, and Miller G. Partnership between tobacco control programs and offices of mental health needed to reduce smoking rates in the United States. *JAMA Psychiatry*, 2013; 70(12):1261–2. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24172829>

Ashare R and Hawk L, Jr. Effects of smoking abstinence on impulsive behavior among smokers high and low in adhd-like symptoms. *Psychopharmacology* 2011; [Epub ahead of print]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21559802>

Allen M, Debanne M, Lazignac C, Adam E, Dickinson L, et al. Effect of nicotine replacement therapy on agitation in smokers with schizophrenia: A double-blind, randomized, placebo-controlled study.

The American Journal of Psychiatry, 2011; 168(4):395–9. Available from:

<http://ajp.psychiatryonline.org/cgi/reprint/appi.ajp.2010.10040569v1>

Ashton M, Lawn S, and Hosking J. Mental health workers' views on addressing tobacco use. The Australian and New Zealand Journal of Psychiatry, 2010; 44(9):846–51. Available from:

<http://informahealthcare.com/doi/full/10.3109/00048674.2010.488637>

Prochaska J, Fromont S, Leek D, Suchanek Hudmon K, Louie A, et al. Evaluation of an evidence-based tobacco treatment curriculum for psychiatry residency training programs. Academic Psychiatry, 2008; 32(6):484–92. Available from: <http://ap.psychiatryonline.org/cgi/content/abstract/32/6/484>

News:

9A.3 People with substance use and mental disorders

Taking Action on Smoking and Health. (2022). *Closing the Inequality Gap: Smoking and Mental Health*. Retrieved from <https://www.ashscotland.org.uk/media/869884/closing-the-inequality-gap-smoking-and-mental-health.pdf>

Harris B, Beswick H, Bowman J, and Bartlem K. No butts – it's time to help people with mental health conditions quit smoking The Conversation, 2018. Available from: <https://theconversation.com/no-butts-its-time-to-help-people-with-mental-health-conditions-quit-smoking-104571>

No authors listed. Giving up cigarettes linked with recovery from illicit substance use disorders. ScienceDaily, 2017. Available from: <https://www.sciencedaily.com/releases/2017/03/170308081136.htm>

No authors listed. 3 myths about mental health and quitting smoking. Truth Initiative (American Legacy Foundation), 2017. Available from: <https://truthinitiative.org/news/3-myths-about-mental-health-and-quitting-smoking>

No authors listed. Cigarette smokers are 10 times more likely to be daily marijuana users. Medical Xpress, 2017. Available from: <https://medicalxpress.com/news/2017-11-cigarette-smokers-daily-marijuana-users.html>

9A.3.1 Prevalence of smoking among people with mental health problems

Kotecha S. Mental health patients 'smoke three times as much'. BBC News, 2015. Available from: <http://www.bbc.com/news/health-32982669>

9A.3.1.2 Smoking prevalence over time

listed Na. Incidence of psychiatric disorders has increased in a shrinking population of smokers. Medical News Today 2016. Available from: <http://www.medicalnewstoday.com/releases/305692.php?tw>

9A.3.2 Contribution of smoking to health outcomes and social inequality

Wootton, R, Sallis, H, & Munafo, M. (2022). *Is there a causal effect of smoking on mental health?* Retrieved from <https://ash.org.uk/wp-content/uploads/2022/06/Causal-effect-smoking-and-mental-health.pdf>

9A.3.3 Explanations for higher smoking prevalence

9A.3.4 Barriers to be overcome when quitting

9A.3.5 Interventions for reducing smoking

Action on Smoking and Health, & Public Mental Health Implementation Centre. (2022). *Public mental health and smoking: a framework for action*. Retrieved from <https://ash.org.uk/wp-content/uploads/2022/06/Public-mental-health-and-smoking.pdf>

O'Dowd A. Call to help smokers with mental health issues quit habit. OnMedica, 2020. Available from: <http://www.onmedica.com/NewsArticle.aspx?id=65636957-e1fa-4bf0-b580-cc709f2ec790>

listed Na. Smoking cessation benefits persist in spite of weight gain in patients with mental illness. Medical News Today 2016. Available from: <http://www.medicalnewstoday.com/releases/308300.php>

listed Na. The stolen years - the mental health and smoking action report. ASH UK, 2016. Available from: http://www.ash.org.uk/files/documents/ASH_1018.pdf

listed Na. No link between anti-smoking drugs, mental health issues: Study. HealthDay News, 2016. Available from: <https://consumer.healthday.com/cancer-information-5/smoking-cessation-news-628/study-doesn-t-find-link-between-anti-smoking-drugs-and-mental-health-side-effects-710238.html>

White V. Trial supports safety and efficacy profile of pfizer's varenicline. European Pharmaceutical Review, 2016. Available from: <http://www.europeanpharmaceuticalreview.com/40620/news/industry-news/trial-pfizer-varenicline/>

9A.3.5.1 Depression

listed Na. Smokers with depression try to quit more often but find it harder. Medical Xpress, 2016. Available from: <http://medicalxpress.com/news/2016-02-smokers-depression-harder.html>

9A.3.5.2 Anxiety

9A.3.5.3 Suicidal ideation

9A.3.5.4 Attention-deficit/hyperactivity disorder (ADHD)

9A.3.5.5 Substance use disorders

Campbell D. Addiction treatment harder to get since councils took over, says tory MP. The Guardian, 2021. Available from: <https://www.theguardian.com/society/2021/jun/10/addiction-treatment-harder-to-get-since-councils-took-over-care-says-tory-mp>

No authors listed. More people in recovery from substance use problems are quitting smoking than ever before. ScienMag, 2019. Available from: <https://scienmag.com/more-people-in-recovery-from-substance-use-problems-are-quitting-smoking-than-ever-before/>

Stern J. Study shows alcohol relapse and cigarette smoking correlation. Yale Daily News, 2015. Available from: <http://yaledailynews.com/blog/2015/09/29/study-shows-alcohol-relapse-and-cigarette-smoking-correlation/>

9A.3.5.6 Post-traumatic stress disorder (PTSD)

9A.3.5.7 Severe mental illness, including bipolar disorder and schizophrenia

9A.3.6 The role of health professionals and health settings

Bryan M. Specialist mental health support available to help lanarkshire smokers quit. Daily Record, 2021. Available from: <https://www.dailyrecord.co.uk/in-your-area/lanarkshire/specialist-mental-health-support-available-24775442>

No authors listed. A change in the air results of a study of smokefree policy and practice in mental health trusts in England. Smoke Free Action, 2019. Available from:

<http://smokefreeaction.org.uk/wp-content/uploads/2019/01/A-Change-In-The-Air.pdf>

No authors listed. Little 'quit-smoking' help at U.S. mental health centers. Medical Xpress, 2018.

Available from: <http://medicalxpress.com/news/2018-05-quit-smoking-mental-health-centers.html>

No authors listed. Tobacco is a social justice issue: Mental health. Truth Initiative, 2018. Available

from: <https://truthinitiative.org/news/tobacco-social-justice-issue-mental-health>

No authors listed. Hospitals use armed police to fine mental health inpatients for smoking. Courier

Mail 2017. Available from: <http://www.couriermail.com.au/news/queensland/hospitals-use-armed-police-to-fine-mental-health-inpatients-for-smoking/news-story/e282c24827d1a3079db5db80c08a00fc>

No authors listed. More mental health facilities are going smoke-free. Truth Initiative, 2017.

Available from: https://truthinitiative.org/news/more-mental-health-facilities-are-going-smoke-free?utm_source=Truth+Initiative+Mailing+List&utm_campaign=261a107b70-Newsletter_072_2017_10_12&utm_medium=email&utm_term=0_c91fd8a5c5-261a107b70-86454907

Borland S. Lose weight and give up smoking or you can't have surgery, hospital tells patients Daily

Mail, 2014. Available from: <http://www.dailymail.co.uk/health/article-2809866/Lose-weight-smoking-t-surgery-hospital-tells-patients.html>

McArthur G. Smokers urged to come clean ahead of surgery Herald Sun, 2014. Available from:

<http://www.heraldsun.com.au/news/smokers-urged-to-come-clean-ahead-of-surgery/story-fni0fiyv-1227091881165?nk=2c142e0bee33c50cbcc60a580404fadc>